

Neotropical Diptera

Neotropical Diptera 17: 1-178 (April 15, 2009)
ISSN 1982-7121
www.neotropicaldiptera.org

Depto. de Biologia - FFCLRP
Universidade de São Paulo
Ribeirão Preto, SP, Brazil

Catalogue of Neotropical Diptera. Asilidae¹

Nelson Papavero

Museu de Zoologia, Universidade de São Paulo, São Paulo, SP, Brasil
Pesquisador Visitante do Departamento de Biologia, Faculdade de Filosofia, Ciências e Letras,
Universidade de São Paulo, Ribeirão Preto, SP, Brasil

In the area covered by this catalogue, the Asilidae are distributed in 8 subfamilies, in a total of 217 genera and 1576 species, as shown in the table below. The catalogue includes 659 references.

Three new generic names are proposed herein: *Amorimius* (for *Macrocolus* Engel, 1930), *Scarboroughia* (for *Furcilla* Martin, 1975) and *Systologaster* (for *Systellogaster* Hermann, 1926). The new name *Cophura gellergrimi* is proposed to replace *Cophura wilcoxi* Kaletta, 1983. *Caenarolia* Thomson, 1869 is revalidated, as its senior synonym, *Allopogon* Schiner, 1886, is preoccupied.

Subfamily	Tribe or Generic group	Genus	Number of species
ASILINAE			599
	<i>Asilus</i> -group		83
		<i>Anarmostus</i>	1
		<i>Chilesus</i>	1
		<i>Cnodalomyia</i>	2
		<i>Cratopoda</i>	2
		<i>Epipamponeurus</i>	1
		<i>Leinendera</i>	1
		<i>Lestophonax</i>	1
		<i>Lochyrous</i>	3
		<i>Lycomya</i>	1
		<i>Machimus</i>	21
		<i>Menexenus</i>	3
		<i>Neomochtherus</i>	1
		<i>Philonicus</i>	6
		<i>Polacantha</i>	7
		<i>Prolatiforceps</i>	8
		<i>Pteralbis</i>	3
		<i>Stenasilus</i>	5
		<i>Stizolestes</i>	8
		<i>Threnia</i>	6
		<i>Wyliea</i>	2
	<i>Efferia</i> -group		141
		<i>Albibarbefferia</i>	16
		<i>Carinefferia</i>	9
		<i>Cratolestes</i>	2
		<i>Diplosynapsis</i>	4

¹ This project was supported by FAPESP grants # 2003/10.274-9 and 2007/50878-1.

	<i>Efferia</i>	31
	<i>Nerax</i>	50
	<i>Pogonioefferia</i>	12
	<i>Porasilus</i>	7
	<i>Triorla</i>	10
	<i>Eicherax</i>-group	18
	<i>Eicherax</i>	8
	<i>Eraxasilus</i>	10
	<i>Eichoichemus</i>-group	15
	<i>Eichoichemus</i>	12
	<i>Proctophoroides</i>	2
	<i>Wygodasilus</i>	1
	<i>Glaphyropyga</i>-group	22
	<i>Glaphyropyga</i>	12
	<i>Leptoharpacticus</i>	1
	<i>Megalometopon</i>	2
	<i>Neotes</i>	1
	<i>Nomomyia</i>	2
	<i>Regasilus</i>	1
	<i>Tsacasia</i>	1
	<i>Zoticus</i>	2
	<i>Lecania</i>-group	26
	<i>Cerozodus</i>	1
	<i>Ctenodontina</i>	4
	<i>Lecania</i>	21
	<i>Lochmorhynchus</i>-group	15
	<i>Apotinocerus</i>	1
	<i>Lochmorhynchus</i>	14
	<i>Mallophora</i>-group	102
	<i>Amblyonychus</i>	16
	<i>Carreraomyia</i>	2
	<i>Mallophora</i>	59
	<i>Megaphorus</i>	7
	<i>Promachella</i>	1
	<i>Promachus</i>	17
	<i>Myaptex</i>-group	21
	<i>Atractocoma</i>	1
	<i>Martintella</i>	1
	<i>Myaptex</i>	2
	<i>Myaptexaria</i>	3
	<i>Rhadinosoma</i>	1
	<i>Scarboroughia</i>	2
	<i>Wilcoxius</i>	11
	<i>Ommatius</i>-group	113
	<i>Ommatius</i>	113
	<i>Proctacanthus</i>-group	43
	<i>Eccritosia</i>	4
	<i>Proctacanthella</i>	4
	<i>Proctacanthus</i>	15
	<i>Taurhynchus</i>	20
	Unplaced or unrecognized species of Asilinae	9
	Unrecognized genera of Asilinae	
	<i>Labromyia</i>	1
	<i>Pediophoneus</i>	(no species included)
DASYPOGONINAE		231
	Dasypogonini	48
	<i>Aczelia</i>	3
	<i>Amorimius</i>	4
	<i>Araucopogon</i>	1
	<i>Austenmyia</i>	1
	<i>Cleptomyia</i>	1

Papavero

<i>Neoderomyia</i>	1
<i>Parataracticus</i>	2
<i>Saropogon</i>	24
<i>Taracticus</i>	10
<i>Tocantinia</i>	
Isopogonini	43
<i>Alvarenga</i>	2
<i>Annamyia</i>	1
<i>Aphamartania</i>	7
<i>Aspidopyga</i>	1
<i>Cophura</i>	25
<i>Hodophylax</i>	2
<i>Nicocles</i>	2
<i>Theromyia</i>	3
Lastaurini	100
<i>Apolastaurooides</i>	1
<i>Blepharepium</i>	7
<i>Caenarolia</i>	10
<i>Diogmites</i>	57
<i>Lastaurina</i>	3
<i>Lastaurus</i>	9
<i>Neodiogmites</i>	11
<i>Phonicocleptes</i>	2
Megapodini	40
<i>Cyrtophrys</i>	4
<i>Deromyia</i>	2
<i>Megapoda</i>	1
<i>Pronomopsis</i>	4
<i>Pseudorus</i>	11
<i>Senobasis</i>	18
LAPHRIINAE	276
Andresosomini	51
<i>Andrenosoma</i>	29
<i>Cerotainiops</i>	2
<i>Dasyllis</i>	6
<i>Pilica</i>	13
<i>Pogonosoma</i>	1
Atomosiini	154
<i>Aphestia</i>	2
<i>Aphractia</i>	3
<i>Atomosia</i>	49
<i>Atomosiella</i>	1
<i>Atoniomya</i>	13
<i>Atractia</i>	2
<i>Bathropsis</i>	4
<i>Cerotainia</i>	35
<i>Cyphomyiactia</i>	1
<i>Dissmyringodes</i>	5
<i>Eumecosoma</i>	13
<i>Hodites</i>	1
<i>Hybozelodes</i>	14
<i>Lamprozona</i>	2
<i>Lycosimyia</i>	2
<i>Oidardis</i>	7
<i>Strombocodia</i>	1
<i>Rhatimomyia</i>	1
Dasytrichini	11
<i>Cryptomyinx</i>	2
<i>Dasythrix</i>	2
<i>Smeryngolaphria</i>	7
Laphriini	41

Catalogue of the Neotropical Diptera. Asilidae

<i>Ichneumolaphria</i>	3
<i>Joartigasia</i>	3
<i>Lampria</i>	19
<i>Laphria</i>	6
<i>Phellopteron</i>	8
<i>Rhopalogaster</i>	
Neophoneini	2
<i>Neophoneus</i>	2
Unplaced species of Laphriinae	17
LAPHYSTIINAE	33
<i>Apoxyrria</i>	2
<i>Asicya</i>	1
<i>Chrysotriclisis</i>	1
<i>Cochleariocera</i>	1
<i>Cymbipyga</i>	1
<i>Gymnotriclisis</i>	1
<i>Helolaphyctis</i>	2
<i>Hexameritia</i>	1
<i>Laphygmolestes</i>	1
<i>Laphystia</i>	3
<i>Macahyba</i>	1
<i>Martinomyia</i>	3
<i>Perasis</i>	1
<i>Protometer</i>	2
<i>Psilocurus</i>	4
<i>Triclioscelis</i>	4
<i>Zabrops</i>	3
Unplaced species of Laphystiinae	2
LEPTOGASTRINAE	89
<i>Apachekulos</i>	8
<i>Beameromyia</i>	12
<i>Eurhabdus</i>	4
<i>Leptogaster</i>	37
<i>Leptopteromyia</i>	7
<i>Psilonyx</i>	6
<i>Schildia</i>	8
<i>Systologaster</i>	6
<i>Tipulogaster</i>	1
STENOPOGONINAE	229
Acronymchini	10
<i>Acronymches</i>	10
Bathypogonini	1
<i>Carebaricus</i>	1
Cyrtopogonini	54
<i>Ablautus</i>	7
<i>Callinicus</i>	1
<i>Cyrtopogon</i>	2
<i>Dasycyrtion</i>	5
<i>Dasypeucus</i>	2
<i>Graptostylus</i>	1
<i>Hadrokolos</i>	1
<i>Heteropogon</i>	9
<i>Holopogon</i>	5
<i>Itolia</i>	5
<i>Ivettea</i>	1
<i>Lonquimayus</i>	3
<i>Metapogon</i>	5
<i>Nannocyrtopogon</i>	1
<i>Nothropogon</i>	1
<i>Raulcortesia</i>	1
<i>Sintoria</i>	3

	<i>Wilcoxia</i>	(unnamed sp.)
Unrecognized species of Cyrtopogonini		1
Dioctriini	1	
	<i>Nannodioctria</i>	1
Echthodopini	1	
	<i>Bohartia</i>	1
Enigmomorphini	55	
	<i>Alyssomyia</i>	5
	<i>Araujoa</i>	1
	<i>Archilestris</i>	5
	<i>Archilestroides</i>	1
	<i>Aymarasilus</i>	1
	<i>Creolestes</i>	5
	<i>Cylicomera</i>	2
	<i>Dicranus</i>	5
	<i>Enigmormorphus</i>	1
	<i>Leptochelina</i>	1
	<i>Microstylum</i>	5
	<i>Pritchardia</i>	5
	<i>Prolepsis</i>	17
	<i>Taperigna</i>	1
Phellini	2	
	<i>Obelophorus</i>	2
Plesiommatini	18	
	<i>Cystoprosopa</i>	2
	<i>Dapsilochetus</i>	1
	<i>Plesiomma</i>	15
Stenopogonini	53	
	<i>Ospriocerus</i>	16
	<i>Scleropogon</i>	9
	<i>Stenopogon</i>	28
Tillobromatini	34	
	<i>Euthrixius</i>	3
	<i>Grajahua</i>	1
	<i>Scylaticina</i>	1
	<i>Scylaticodes</i>	4
	<i>Tillobroma</i>	21
	<i>Zabrotica</i>	1
	Unrecognized sp.	3
STICHOPOGONINAE	28	
	<i>Argyropogon</i>	1
	<i>Lasiopogon</i>	3
	<i>Lissoteles</i>	9
	<i>Stichopogon</i>	5
	<i>Townsendia</i>	10
TRIGONOMIMINAE	43	
	<i>Haplopogon</i>	2
	<i>Holcocephala</i>	37
	<i>Meliponomima</i>	1
	<i>Orrhodops</i>	2
	<i>Seabramyia</i>	1
Unplaced genus of Asilidae		
	<i>Araripogon</i>	1
Unplaced species of Asilidae		47

List of acronyms

- AMNH – American Museum of Natural History, New York
BMNH – National Museum of Natural History, London
BUD – Magyar Természettudományi Múzeum, Budapest, Hungary
BZM – Berliner Zoologische Museum, Humboldt-Universität, Berlin, Germany
CAS – California Academy of Sciences, San Francisco, California, U. S. A.
CIS – California Insect Survey
COPN – Universitetets Zoologiske Museum, Copenhagen, Denmark
CSCLB – California State College, Long Beach, California, U. S. A.
CU – Cornell University, Ithaca, New York, U. S. A.
DEI – Deutsches Entomologisches Institut, Berlin, Germany
DZUC – Departamento de Zoología, Universidad de Concepción, Concepción, Chile
FUSAGembloix – Faculté Universitaire des Sciences Agronomiques de Gembloux, Gembloux, Namur, Belgium
FAUCV – Facultad de Agronomía, Universidad Central de Venezuela
FSCA – Florida State Collection of Arthropods, Gainesville, Florida, U. S. A.
IOC – Instituto Oswaldo Cruz, Rio de Janeiro, Brasil
IZAUCV – Instituto de Zoología Agrícola, Universidad Central de Venezuela
IZPAN – Instytut Zoologiczny Polskiej Akademii Nauk, Warsaw, Poland
KIEL – Zoologisches Museum, Christian-Albrechts Universität, Kiel, Germany
KANS – Kansas State University, Manhattan, Kansas, U.S.A.
KU – Snow Collection, University of Kansas, Lawrence, Kansas, U. S. A.
MCSNM – Museo Civico di Storia Naturale, Milan, Italy
MCZ – Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, U. S. A.
MNHNP – Muséum National d'Histoire Naturelle, Paris, France
MNHNS – Museo Nacional de Historia Natural, Santiago, Chile
MNRJ – Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brasil
MZUSP – Museu de Zoologia, Universidade de São Paulo, São Paulo, Brasil
NMW – Naturhistorisches Museum, Vienna, Austria
NCDA – North Carolina Department of Agriculture
OSU – Ohio State University, Columbus, Ohio, U. S. A.
OXF – Hope Museum, Oxford University, Oxford, England
SENCK – Senckenbergische Naturforschende Gesellschaft. Natur-Museum und Forschingsinstitut, Frankfurt a. M., Germany
SIKJ – Science Institute, Kingston, Jamaica
SMTD – Staatliches Museum für Tierkunde, Dresden, Germany
TORO – Istituto e Museo di Zoologia, Università di Torino, Turin, Italy
UCB – University of California, Berkeley, California, U. S. A.
UCD – University of California, Davis, California, U. S. A.
USNM – United States National Museum of Natural History, Smithsonian Institution, Washington, D. C.
UUEC – University of Utah Entomological Collection
ZMA – Zoölogisch Museum, Amsterdam, Holland
ZSBS – Zoologische Sammlung des Bayerischen Staates, Munich, Germany

Acknowledgments – Drs. Marcia Souto Couri (Museu Nacional, Universidade Federal do Rio de Janeiro), Carlos J. E. Lamas (Museu de Zoologia, Universidade de São Paulo), Aubrey G. Scarbrough (Department of Entomology, University of Arizona, Tucson) and Fritz Geller-Grimm (Wiesbaden Museum) have been extremely kind in providing literature; to them my best thanks.

P.S.: While this paper was in press, Dikow (2009) has proposed a new phylogeny for the Asilidae.

Family ASILIDAE Latreille, 1802

Asilici Latreille, 1802: 432. Type-genus, *Asilus* Linnaeus, 1758. Ref. – Dikow, 2004: 461.
Asilides Leach, 1815: 161.
Asilidae Kirby & Spence, 1817: 360, 379.
Asilimi Schinz, 1823: 861.
Asilica Burmeister, 1829: 24.
Asiliadae Harris, 1833: 594.
Asilina Newman, 1834: 379, 391.
Asilites Newman, 1834: 379, 392.
Asilitae Macquart, 1838: 79bis (1839: 195bis).
Asilopsinae Cockerell, 1920: 251.

Subfamily Asilinae Latreille, 1802

Apocliina Lehr, 1969: 532, *nomen nudum*. Ref. – Dikow, 2004: 461.
Apocleinae Papavero, 1973a: 233, 236.

Asilus-group

Lycomyiaria Lynch Arribálzaga, 1883a: 145. Type-genus, *Lycomya* Bigot, 1857.
Machimina Lehr, 1969: 532 (341), *nomen nudum*.
Machimini Lehr, 1977: 50. Type-genus, *Machimus* Loew, 1849. Ref. – Dikow, 2004: 262.

Genus **Anarmostus** Loew, 1860

Anarmostus Loew, 1860: 142. Type-species, *Asilus iopterus* Wiedemann, 1828 (orig. des.).

iopterus (Wiedemann), 1828: 438 (*Asilus*). Type-locality: “Brazil”. Distr. – Guyana, Peru (Loreto), Brazil (Pará), Bolivia (Santa Cruz). Refs. – Walker, 1849: 434 (*Asilus*), 1854: 701 (*Asilus*); Schiner, 1866: 684; Osten Sacken, 1887: 211; Williston, 1891: 89 (cat.); Aldrich, 1905: 275 (cat.); Kertész, 1909: 272 (cat.); Bromley, 1946: 117 (cat.); Carrera & d’Andretta, 1953: 74; Carrera, 1960a: 165; Hull, 1962 (2): 544, figs. 393, 759, 1617-1618, 2254, 2306, 2362, 2444; Martin & Papavero, 1970: 84 (cat.); Papavero, 1971d: 27; Lamas M., 1972: 314 (cat.); Fisher, 1985: 33. Syntypes, SENCK [cf. Geller-Grimm, 1999: 210], BZM [cf. Schumann, 1973: 121].
albolineata Macquart, 1850: 378 (1850: 74) (*Laphria*) Type-locality: Bolivia, Mojos [= Depto. Santa Cruz]. Refs. – Hull, 1962 (1): 323 (*Laphria*); Papavero, 1971d: 27 (syn.). Type, MNHNP.

Genus **Chilesus** Bromley, 1932

Chilesus Bromley, 1932: 272. Type-species, *Chilesus geminatus* Bromley, 1932 (orig. des.).

geminatus Bromley, 1932: 272. Type-locality: Chile, Magallanes. Distr. – Chile (Aysén, Magallanes), Argentina (Chubut). Ref. – Hull, 1962 (2): 538, figs. 210, 755, 1404, 1413, 2194, 2298, 2438; Artigas, 1970a: 262, figs. 223-226, 495. Type, BMNH.

Genus **Cnodalomyia** Hull, 1962

Cnodalomyia Hull, 1962 (2): 516. Type-species, *Cnodalomyia obtusa* Hull, 1962 (orig. des.).

catarinensis Lamas & Mellinger, 2008: 38, figs. 1-11. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (São Paulo, Paraná, Santa Catarina). Type, MZUSP.
obtusa Hull, 1962: 518, figs. 787, 1546, 1555, 2195, 2271. Type-locality: Brazil, Rio de Janeiro, Alto do Itatiaia. Distr. – Brazil (Rio de Janeiro). Refs. – Martin & Papavero, 1970: 83 (cat.); Lamas & Mellinger, 2008: 38, fig. 2B, 39, fig. 3B. Type, USNM.

Genus **Cratopoda** Hull, 1962

Cratopoda Hull, 1962 (2): 575. Type-species, *Asilus gayi* Macquart, 1838 (orig. des., misident.) = *Cratopoda helix* Bromley, 1935).

emarginata Artigas, 1970a: 271, figs. 236-241. Type-locality: Chile, Ñuble, Cordillera Talca, El Radal. Distr. – Chile (Biobío, Linares, Malleco, Ñuble), Argentina (Neuquén). Ref. – Artigas, 1971: 33, figs. 79-80. Type, DZUC.
helix (Bromley), 1935b: 110, fig. 1 (*Asilus*). Type-locality: “Central Chile”. Distr. – Chile (Aconcagua, Arauco, Biobío, Cautín, Concepción, Coquimbo, Curicó, Linares, Malleco, Maule, Ñuble, O’Higgins, Santiago, Talca, Valparaíso). Refs. – Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962: 545 (*Asilus*); Artigas, 1970a: 273, figs. 1-12, 242-247, 483 (n. comb.); Martin & Papavero, 1970: 97 (cat.); Artigas, 1971: 34, figs. 81-82. Type, DEI.
gayi Macquart of Hull, 1962 (2): 576, figs. 396, 737, 1527, 1536, 2227, 2430, 2454, misident.

Genus **Epipamponeurus** Becker, 1919

Epipamponeurus Becker, 1919: 166. Type-species, *Epipamponeurus americanus* Becker, 1919 (mon.).

americanus Becker, 1919: 166. Type-locality: Ecuador, Chimborazo, Riobamba, 2754 m, and Pichincha, Casitagua, 3512 m. Distr. – Ecuador. Ref. – Martin & Papavero, 1970: 84 (cat.). Syntypes, MNHNP.

Genus **Leinendera** Carrera, 1945

Leinendera Carrera, 1945c: 184. Type-species, *Leinendera rubra* Carrera (orig. des.).

rubra Carrera, 1945c: 185, figs. 13, 19, 22-28. Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Distr. – Brazil (Rio de Janeiro, São Paulo). Refs. – Carrera, 1958a: 150; Hull, 1962 (2): 512, figs. 379, 797, 1400, 1409, 2419-2420; Martin & Papavero, 1970: 83 (cat.). Type, MZUSP.

Genus **Lestophonax** Hull, 1962

Lestophonax Hull, 1962 (2): 568. Type-species, *Lestophonax mallophoroides* Hull, 1962 (orig. des.).

mallophoroides Hull, 1962 (2): 569, figs. 336, 783, 1549, 1558, 2218, 2498-2499. Type-locality: Ecuador, Azuay, Silván, 3400 m. Distr. – Ecuador. Ref. – Martin & Papavero, 1970: 87 (cat.). Type, AMNH.

Genus **Lochyrus** Artigas, 1970

Lochyrus Artigas, 1970a: 296. Type-species, *Asilus crassus* Bromley, 1932 (orig. des.).

Ref. – Artigas, 1970a: 297 (key to spp.).

balmacedensis Artigas, 1970a: 297, figs. 274-277, 279, 416. Type-locality: Chile, Aysén, Balmaceda. Distr. – Chile (Aysén, Magallanes). Type, DZUC.
crassus (Bromley), 1932: 279, figs. 27-28 (*Asilus*). Type-locality: Chile, Concepción. Distr. – Chile (Arauco, Biobío, Cautín, Concepción, Colchagua, Curicó, Linares, Malleco, Ñuble, O’Higgins, Santiago, Talca, Valparaíso). Refs. – Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 299, figs. 278-284, 286, 425 (n. comb.); Martin

& Papavero, 1970: 88 (cat; *Asilus*, as unplaced sp.); Artigas, 1971: 38, figs. 93-94. Type, BMNH.
frezieri Artigas, 1970a: 303, figs. 285, 287-290, 485. Type-locality: Chile, Arauco, Caramávida. Distr. – Chile (Aconcagua, Arauco, Biobío, Cautín, Concepción, Coquimbo, Curicó, Linares, Malleco, Ñuble, Santiago, Talca, Valdivia, Valparaíso). Ref. – Artigas, 1971: 39, figs. 95-96. Type, DZUC.

Genus **Lycomya** Bigot, 1857

Lycomya Bigot, 1857c: 288. Type-species, *Lycomya germainii* Bigot, 1857 (mon.).
Lycomyia, emend.

germainii Bigot, 1857c: 290, pl. 6, figs. 3a-b. Type-locality: “Chile”. Distr. – Chile (Arauco, Cautín, Concepción, Curicó, Malleco, Osorno, Santiago, Valdivia, Valparaíso). Refs. – Philippi, 1865: 694; Schiner, 1866: 706, 1868: 169; Osten Sacken, 1891: 419; Williston, 1891: 419 (cat.); Kertész, 1909: 208 (cat.); Bromley, 1932: 269; Stuardo Ortiz, 1946: 84 (cat.); Carrera, 1948b: 424, figs. 1-2; Carrera & Vulcano, 1961: 76 (preys); Hull, 1962 (2): 536, figs. 367, 754, 1402, 1411, 1566, 2197, 2289, 2358, 2504 (*Lycomya*); Artigas, 1970a: 307, figs. 291-295, 297, 409; Martin & Papavero, 1970: 84 (cat.); Artigas, 1971: 40, figs. 97-98. Type, OXF.

Genus **Machimus** Loew, 1849

Machimus Loew, 1849: 1. Type-species, *Asilus chrysitis* Meigen, 1820 (sub. des., Coquillett, 1910: 564).

Ref. – Martin, 1975c: 9-49 (rev.).

acutus Martin, 1975c: 35, figs. 8, 23. Type-locality: Mexico, Michoacán, Carapán. Distr. – Mexico (Michoacán). Type, UCD.
alterus (Williston), 1901: 330 (*Asilus (Tolmerus)*). Type-locality: Mexico, Guerrero, Omiltemi, 3000 ft. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 281 (cat.; *Tolmerus*); Kertész, 1909: 297 (cat.; *Tolmerus*); Hull, 1962 (2): 572 (*Tolmerus*); Martin & Papavero, 1970: 85 (cat.); Martin, 1975c: 44; Papavero & Ibáñez Bernal, 2003: 177 (hist. Mexican dipteral). Syntypes, BMNH, AMNH.

tenebrosus Williston of Williston, 1901: 328 (1 ST), misident. Ref. – Martin, 1975c: 33.

avidus (Wulp), 1869: 82 (*Asilus*). Type-locality: U. S. A., Wisconsin. Distr. – U. S. A. (Colorado, New Mexico), Mexico (Coahuila). Ref. – Hine, 1909: 147, fig. 6 (*Asilus*); Baker, 1939: 170 (*Asilus*); Hull, 1962 (2): 562 (n. comb.). Type, ZMA. [= *sadyates*?].

brevis Martin, 1975c: 36, fig. 16. Type-locality: Mexico, Durango, 10 miles west of El Salto, 9000 ft. Distr. – Mexico (Durango). Type, CNM.

bromleyanus (Carrera & d'Andretta), 1950: 179, figs. 7, 16 (*Asilus*). Type-locality: Mexico, Michoacán, Cerro Tancítaro. Distr. – Mexico (Michoacán). Refs. – Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 86 (cat.); Martin, 1975c: 36, fig. 34. Type, CMNH.

cancerae Martin, 1975c: 46. Type-locality: Mexico, Zacatecas, ½ mile south of Tropic of Cancer, highway 45, km post 758. Distr. – Mexico (Morelos, Zacatecas). Type, CAS.

cerasinus Martin, 1975c: 48, figs. 31-32. Type-locality: Mexico, Puebla, 20 miles north of Tehuacán, 6000 ft. Distr. – Mexico (Puebla). Type, CAS.

coleus Martin, 1975c: 17 (in key), 26, fig. 70. Type-locality: U. S. A., California, Los Angeles County, Azusa. Distr. – U. S. A. (California), Mexico. Type, CAS.

gertschi (Bromley), 1951: 33 (*Asilus*). Type-locality: Panama, Chiriquí, El Volcán. Distr. – Panama. Refs. – Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 86 (cat.); Martin, 1975c: 28, fig. 39. Type, AMNH.

guttatus Martin, 1975c: 42, figs. 7, 19. Type-locality: Mexico, Chiapas, 6 miles southeast of San Cristóbal. Distr. – Mexico (Chiapas). Type, CAS.

humilis (Bellardi), 1861: 151 (51) (*Asilus*). Type-locality: “Mexico”. Distr. – Mexico (Chihuahua, Distrito Federal, Guanajuato, Jalisco, México, Michoacán, Morelia, Puebla). Refs. – Aldrich, 1905: 283 (cat.; *Lophonotus*); Kertész, 1909: 276 (cat.; *Dysmachus*); Hull, 1962 (2): 514 (*Cerdistus*); Martin & Papavero, 1970: 86 (cat.); Martin, 1975c: 49, figs. 37, 77; Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipteral). Syntypes, TORO.

lurettae Lavigne, 1978: 55 (nom. nov. for *truncatus* Martin, 1975). Type-locality: Mexico, Morelos, Cuernavaca. Distr. – Mexico (Morelos).

truncatus Martin, 1975c: 41, fig. 18 (preocc. Oldroyd, 1972). Type-locality: Mexico, Morelos, Cuernavaca, 550 ft. Type, COR.

mealpinei Martin, 1975c: 38, fig. 10. Type-locality: Mexico, Durango, 3 miles east of El Salto, 8200 ft. Distr. – Mexico (Durango, Sinaloa). Type, CMN.

?mexicanus (Macquart), 1846: 222 (1846: 94) (*Asilus*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Walker, 1855: 695

- (*Asilus*); Aldrich, 1905: 282 (car.; *Asilus*); Kertész, 1909: 260 (cat.; *Asilus*); Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*); Papavero & Ibáñez Bernal, 2001: 85 (hist. Mexican dipterol.). Type, ?
- niveibarbus** (Bellardi), 1861: 153 (53) (*Asilus, Epitreptus* [sic]). Type-locality: Mexico, Veracruz, Córdoba. Distr. – Mexico (Veracruz, Distrito Federal). Refs. – Kertész, 1909: 296 (cat.; *Epitreptus*); Carrera & d'Andretta, 1950: 178, figs. 49-52 (*Asilus*); Hull, 1962 (2): 573 (*Epitreptus*); Martin & Papavero, 1970: 86 (cat.); Martin, 1975c: 37, figs. 9, 25, 33; Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Syntypes, TORO.
- niveibarba* Aldrich, 1905: 281 (cat.; *Epitreptus*), error.
- occidentalis** (Hine), 1909: 147, fig. 7 (*Asilus*). Lectotype-locality: U. S. A., California, Los Angeles County. Distr. – Canada, U. S. A., Mexico (Baja California). Refs. – Hull, 1962 (2): 562; Martin & Wilcox, 1965: 396 (cat.); Martin & Papavero, 1970: 85 (cat.); Martin, 1975c: 19, figs. 21, 72; O'Neill, 1992: 34 (biol.), 1995: 248 (predator of grasshoppers); O'Neill & Kemp, 1992: 323 (behavioral thermoregulation). Lectotype OSU [des. by Martin, 1975: 20].
- oriens** Martin, 1975c: 39, fig. 22. Type-locality: Mexico, México, highway 190, km post 49. Didtr. – Mexico (México). Type, CAS.
- painteri** Martin, 1975c: 40, fig. 20. Type-locality: Mexico, Morelos, Cuernavaca. Distr. – Mexico (Chiapas, Durango, Guerrero, Hidalgo, Michoacán, Morelos, Oaxaca, Puebla, Zacatecas). Type, CAS.
- submaculus** Martin, 1975c: 43, figs. 4, 27, 36. Type-locality: Mexico, west slope of Popocatépetl, 10,000 ft. Distr. – Mexico (Michoacán, Morelos). Type,
- tenebrosus** (Williston), 1901: 328 (*Asilus (Asilus)*). Type-locality: Mexico, Guerrero, Omiltemi, 8000 ft. Distr. – U. S. A. (Arizona, Colorado, New Mexico, Utah), Mexico (Durango, Guerrero). Refs. – Aldrich, 1905: 283 (cat.; *Asilus*); Hine, 1909: 146, fig. 5 (*Asilus*); Kertész, 1909: 260 (cat.; *Asilus*); Brown, 1929: 317, pl. 30, figs. 56-57 (*Asilus*); Bromley, 1935c: 2 (*Asilus*); Baker, 1939: 169 (*Asilus*); Carrera & d'Andretta, 1950: 178, fig. 15 (*Asilus*); Hull, 1962 (2): 562; Martin & Wilcox, 1965: 396 (cat.); Martin & Papavero, 1970: 86 (cat.); Martin, 1975c: 33, fig. 14; Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Syntypes, BMNH, AMNH.
- ventralis** Martin, 1975: 34, figs. 28, 42. Type-locality: Honduras, El Zamorano, Francisco Morazán, Mt. Uyuca, 6000 ft. Distr. – Honduras. Type, OSU.

Genus Menexenus Artigas, 1970

Menexenus Artigas, 1970a: 317. Type-species, *Asilus concepcionensis* Bromley, 1932 (orig. des.).

Ref. – Artigas, 1970a: 318 (key to spp.).

- concepcionensis** (Bromley), 1932: 281 (*Asilus*). Type-locality: Chile, Concepción. Distr. – Chile (Aconcagua, Concepción, Coquimbo, Linares, Maule, Malleco, Ñuble, Santiago, Valparaíso), Argentina (Río Negro). Refs. – Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 319, figs. 334-340, 408 (n. comb.); Martin & Papavero, 1970: 88 (cat.; *Asilus*, as unplaced sp.); Artigas, 1971: 42, figs. 103-104. Type, BMNH.
- schlengeri** Artigas, 1982: 19, figs. 1-11. Type-locality: Argentina, Chubut, 13 km north of Puerto Madryn, 120 m. Distr. – Argentina (Chubut). Type, DZUC.
- veredus** Artigas, 1970a: 321, figs. 341-344, 424. Type-locality: Chile, Santiago, El Canelo. Distr. – Chile (Santiago). Type, DZUC.

Genus Neomochtherus Osten Sacken, 1878

Mochtherus Loew, 1849: 58 (preocc. Schmidt-Goebel, 1846). Type-species, *Asilus pallipes* Meigen, 1820 (sub. des., Coquillett, 1910: 575).

Neomochtherus Osten Sacken, 1878: 82 (cat.; *nom. nov.* for *Mochtherus* Loew, 1849). Type-species, *Asilus pallipes* Meigen, 1820 (aut.).

californicus (Hine), 1909: 164, fig. 36 (*Asilus*). Type-locality: U. S. A., California, Cisco, 5000 ft. Distr. – U. S. A. (Oregon, Idaho, Nevada, California), Mexico (Baja California). Refs. – Hull, 1962 (2): 545 (*Asilus*); Martin & Wilcox, 1965: 392 (cat.; *Asilus*); Wilcox, 1965: 392 (*Asilus*).

oregonae Martin, 1975c: 92. Type-locality: U. S. A., Oregon, 2 miles west of Sisters. Type, CAS.

Genus Philonicus Loew, 1849

Philonicus Loew, 1849: 144. Type-species, *Asilus albiceps* Meigen, 1820 (mon.).

Philonotus Neuhaus, 1886: 64 (unnecessary *nom. nov.* for *Philonicus* Loew, 1849). Type-species, *Asilus albiceps* Meigen, 1820 (aut.).

One species known as fossil: *Philonicus saxorum* James (1939: 44) (Oligocene; U. S. A., Colorado).

- albospinosus** (Bellardi), 1861: 154 (54) (*Asilus*, *?Epitreptus*). Type-locality: Mexico, surroundings of Mexico City. Distr. – Mexico. Refs. – Aldrich, 1905: 281 (cat.; *?Epitreptus*); Kertész, 1909: 294 (cat.; *Epitreptus*); Hull, 1962 (2): 573; Martin & Papavero, 1970: 85 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, (male) TORO.
- taeniatus** Bellardi, 1861: 155 (55) (*Asilus*, *Philonicus*). Type-locality: “Mexico”. Refs. – Kertész, 1909: 289 (cat.; *Heligmoneura*, as syn. of *truquii* Bellardi, 1861); Hull, 1962 (2): 590 (*Neomochtherus*); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, (female) TORO.
- arizonensis** (Williston), 1893a: 73 (*Stenoprosopus*). Type-locality: U. S. A., Arizona. Distr. – U. S. A. (Arizona, New Mexico), Mexico (Chihuahua). Refs. – Hull, 1962 (2): 562; Martin & Wilcox, 1965: 398 (cat.); Martin & Papavero, 1970: 85 (cat.).
- fuliginosus** (Bellardi), 1861: 152 (52) (*Asilus*, *?Mochtherus*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Aldrich, 1905: 281 (cat.; *Heligmoneura*); Kertész, 1909: 281 (cat.; *Heligmoneura*); Hull, 1962 (2): 590 (*Neomochtherus*); Martin & Papavero, 1970: 85 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- plebeius** (Osten Sacken), 1887: 209 (*Neomochtherus*). Type-locality: Mexico, northern Sonora. Distr. – Mexico (Sonora). Refs. – Hull, 1962 (2): 590 (*Neomochtherus*; as syn. of *truquii* Bellardi, 1861); Martin & Papavero, 1970: 85 (cat.); Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.
- plebius* Martin, 1965b: 131, error.
- truquii** (Bellardi), 1861: 152 (52) (*Asilus*, *Mochtherus*). Type-locality: “Mexico”. Distr. – U. S. A. (Arizona), Mexico. Refs. – Wulp, 1882b: 116 (*Mochtherus*); Williston, 1901: 328 (*Heligmoneura*); Aldrich, 1905: 281 (cat.; *Heligmoneura*); Hine, 1909: 169, figs. 42-43 (*Asilus*); Kertész, 1909: 289 (cat.; *Heligmoneura*); Bromley, 1935c: 4 (n. comb.); Hull, 1962 (2): 590 (*Neomochtherus*); Martin & Wilcox, 1965: 398 (cat.); Martin & Papavero, 1970: 85 (cat.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- tuxpanganus** (Bellardi), 1862: 119 (22) (*Asilus*). Type-locality: Mexico, Tuxpango, near Orizaba. Distr. – Mexico (Michoacán, San Luís Potosí, Veracruz), Guatemala. Refs. – Aldrich, 1905: 282 (cat.; *Philoducus*); Hine, 1907: 118, 1909: 169; Kertész, 1909: 270 (cat.); Carrera & d’Andretta, 1950: 177 (*Asilus*); Hull, 1962 (2): 562; Martin & Papavero, 1970: 85 (cat.); Papavero & Ibáñez Bernal, 2001: 110 (hist. Mexican dipterol). Type, TORO.

Genus *Polacantha* Martin, 1975

Polacantha Martin, 1975c: 49. Type-species, *Asilus compositus* Hine, 1918 (orig. des.).

Polacantha, subg. *Echinotropis* Martin, 1975c: 60. Type-species, *Asilus (Heligmoneura) xanthocerus* Williston, 1901 (orig. des.). **n.syn.**

Ref. – Martin, 1975c: 49-61 (rev.).

- badia** Martin, 1975c: 53 [*Polacantha (Polacantha)*]. Type-locality: Mexico, Hidalgo, La Zorra, highway 85, km post 276. Distr. – Mexico (Hidalgo). Type, CAS.
- brevis** Martin, 1975c: 54 [*Polacantha (Polacantha)*]. Type-locality: Mexico, Chiapas, Tuxtla Gutiérrez, 1000 ft. Distr. – Mexico (Chiapas). Type, COR.
- composita** (Hine), 1918: 321 (*Asilus*). Type-locality: U. S. A., California, San Diego. Distr. – U. S. A. (Arizona, California, Nevada, New Mexico, Oregon, Texas, Utah), Mexico (Baja California and Chihuahua, Coahuila to San Luís Potosí). Refs. – Distr. – Bromley, 1935c: 4 (*Asilus*); Baker, 1939: 169 (*Erax*); Hull, 1962 (2): 545 (*Asilus*); Martin, 1975c: 55, figs. 43, 45, 58 ([*Polacantha (Polacantha)*], n. comb.). Type, OSU.
- petila** Martin, 1975c: 57 [*Polacantha (Polacantha)*]. Type-locality: Mexico, Durango, 24 miles west of La Ciudad. Distr. – Mexico (Durango). Type, CNM.
- sinuosa** Martin, 1975c: 58 [*Polacantha (Polacantha)*]. Type-locality: U. S. A., Texas, Chisos Mountains, Rim trail to Juniper Flats. Distr. – U. S. A. (Texas), Mexico (Durango, Nuevo León, Tamaulipas). Type, CAS.
- tridens** Martin, 1975c: 59, fig. 44. Type-locality: Mexico, Hidalgo, Zimapán. Distr. – Mexico (Durango, Hidalgo). Type, CAS.
- xanthocera** (Williston), 1901: 329 (*Asilus (Heligmoneura)*). Type-locality: Mexico, Guerrero: Dos Arroyos, río Papagaio, Venta de Zopilote, and Tierra Colorada. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 281 (cat.; *Heligmoneura*); Kertész, 1909: 289 (cat.; *Heligmoneura*); Hull, 1962 (2): 514 (*Cerdistus*); Martin & Papavero, 1970: 91 (cat.; unplaced); Martin, 1975c: 60, fig. 60 ([*Polacantha (Echinotropis)*]; n. comb.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Syntypes, BMNH, TP SMTD [cf. Kallweit, 1992: 198].

Genus **Prolatiforceps** Martin, 1975

Prolatiforceps Martin, 1975c: 61. Type-species, *Asilus melanocerus* Williston, 1901 (orig. des.).

Ref. – Martin, 1975c: 61-71 (rev.).

anonymus (Williston), 1901: 330 (*Asilus (Neoitamus)*). Type-locality: Mexico, Guerrero: Omiltemi and Sierra de las Aguas Escondidas, 8000-9500 ft. Distr. – Mexico (Guerrero, Chiapas, Michoacán, San Luis Potosí, Sinaloa). Refs. – Aldrich, 1905: 282 (cat.; *Asilus*); Kertész, 1909: 299 (cat.; *Machimus*); Carrera & d'Andretta, 1950: 178, figs. 3, 17, 30, 34 (*Asilus*); Hull, 1962 (2): 514 (*Cerdistus*); Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*); Martin, 1975c: 64, figs. 57, 68 (n. comb.); Papavero & Ibáñez Bernal, 2003: 177 (hist. Mexican dipterol). Syntypes, BMNH, AMNH, SMTD [cf. Kallweit, 1992: 194].

capillatus (Williston), 1901: 330 (*Asilus (Neoitamus)*). Type-locality: Mexico, Guerrero, Omiltemi, 8000 ft. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 281 (cat.; *Neoitamus*); Kertész, 1909: 291 (cat.; *Neoitamus*); Hull, 1962 (2): 557 (*Neoitamus*); Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*); Martin, 1975c: 64 (n. comb.); Papavero & Ibáñez Bernal, 2003: 177 (hist. Mexican dipterol). Syntypes, BMNH, AMNH.

dolichomerus (Williston), 1901: 329 (*Asilus (Heligmoneura)*). Type-locality: Mexico, Guerrero: Omiltemi, Sierra de las Aguas Escondidas, and Amula, 6000-9500 ft. Distr. – Mexico (Distrito Federal, Durango, Guerrero, Jalisco, Michoacán, Morelos, Oaxaca, Sinaloa). Refs. – Aldrich, 1905: 281 (cat.; *Heligmoneura*); Kertész, 1909: 286 (cat.; *Heligmoneura*); Carrera & d'Andretta, 1950: 178, figs. 45-48 (*Asilus*); Hull, 1962 (2): 514 (*Cerdistus*); Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*); Martin, 1975c: 65, figs. 55, 64 (n. comb.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Syntypes, BMNH, AMNH, SMTD [cf. Kallweit, 1992: 195].

fenestellus Martin, 1975c: 67, figs. 56, 61. Type-locality: Mexico, Guerrero, Omiltemi, 8000 ft. Distr. – Mexico (Guerrero). Type, CAS.

infuscatus (Bellardi), 1861: 156 (56), pl. 2, fig. 15 (*Asilus*). Type-locality: “Mexico”. Disrtr. – Mexico (Nuevo León, Veracruz). Refs. – Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*); Martin, 1975c: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Syntypes, TORO.

melanocerus (Williston), 1901: 329 (*Asilus (Heligmoneura)*). Type-locality: Mexico, Guerrero, Omiltemi, 8000 ft. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 281 (cat.; *Heligmoneura*); Kertész, 1909: 281 (cat.; *Heligmoneura*); Hull, 1962 (2): 514 (*Cerdistus*); Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*); Martin, 1975c: 69, figs. 51, 62 (n. comb.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Syntypes, BMNH, AMNH.

neriacensis (Bromley), 1951: 34 (*Asilus*). Type-locality: Mexico, Neriaco, 10,500 ft. Distr. – Mexico. Refs. – Hull, 1962 (2): 545 (*Asilus*); Martin, 1975c: 70, figs. 54, 63 (n. comb.). Type, AMNH.

neariacensis Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*), error.

nigrocaudatus (Williston), 1901: 329 (*Asilus (Neoitamus)*). Type-locality: Mexico, Guerrero: Omiltemi, and Xucumanatlán, 7000-8000 ft. Distr. – Mexico (Guerrero, Jalisco, Michoacán). Refs. – Aldrich, 1905: 281 (cat.; *Neoitamus*); Kertész, 1909: 293 (cat.; *Neoitamus*); Hull, 1962 (2): 557 (*Neoitamus*); Martin & Papavero, 1970: 86 (cat.; *Machimus (Tolmerus)*); Martin, 1975c: 70 (n. comb.); Papavero & Ibáñez Bernal, 2003: 177 (hist. Mexican dipterol). Syntypes, BMNH, AMNH.

Genus **Pteralbis** Ayala, 1981

Pteralbis Ayala, 1981: 51. Type-species, *Neoitamus peregrinus* Carrera & Machado-Allison, 1963 (orig. des.).

borburatensis Ayala, 1981: 55, figs. 12-13. Type-locality: Venezuela, Carabobo, río Borburata, 250 m. Distr. – Venezuela (Carabobo). Type, IZAUCV.

jolyi Ayala, 1981: 54, figs. 1-5, 8-9. Type-locality: Venezuela, Miranda, La Silla de Caracas, vía al Paraíso, 1300 m. Distr. – Venezuela (Aragua, Miranda). Type, IZAUCV.

peregrinus (Carrera & Machado-Allison), 1963: 258, figs. 4-5 (*Neoitamus*). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. – Venezuela (Aragua). Refs. – Martin & Papavero, 1970: 90 (cat.; unplaced); Ayala, 1981: 53, figs. 6-7, 10-11 (n. comb.). Type, FAUCV.

Genus **Stenasilus** Carrera, 1960

Stenasilus Carrera, 1960a: 166. Type-species, *Asilus tenuis* Wiedemann, 1828 (orig. des.).
Senoprosopis Macquart of authors, misident.

- brasiliensis** (Schiner), 1867: 404 (*Senoprosopis*). Type-locality: “Brazil”. Distr. – Brazil (Rio de Janeiro, São Paulo). Refs. – Williston, 1891: 89 (cat.; *Senoprosopis*); Kertész, 1909: 284 (cat.; *Stenopropsis*); Bromley, 1946: 117 (cat.; *Senoprosopis*); Carrera, 1945c: 189, figs. 29-35 (*Senoprosopis*); Hull, 1962 (2): 500, fig. 703 (*Senoprosopis*); Martin & Papavero, 1970: 61 (cat.). Type, NMW.
- impedens** (Wiedemann), 1828: 587 (*Asilus*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Schiner, 1866: 690 (*Erax*); Jaennicke, 1867: 366 (*Senoprosopis*); Williston, 1891: 85 (cat.; *Senoprosopis*); Kertész, 1909: 284 (cat.; *Stenopropsis*); Bromley, 1946: 117 (cat.; *Senoprosopis*); Hull, 1962 (2): 500, figs. 701, 1492, 1501, 2312 (*Senoprosopis*); Martin & Papavero, 1970: 61 (cat.). Type, SENCK [cf. Geller-Grimm, 1999: 209].
- lutipes** (Wiedemann), 1828: 454 (*Asilus*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1855: 707 (*Asilus*); Schiner, 1866: 685 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 259 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; unplaced); Geller-Grimm, 1999: 210 (n. comb.). Type, SENCK [cf. Geller-Grimm, 1999: 210].
- tenuis** (Wiedemann), 1828: 448 (*Asilus*). Type-locality: “Brazil”. Distr. – Guyana, Brazil (Rio de Janeiro). Refs. – Walker, 1855: 706 (*Asilus*); Schiner, 1866: 685 (*Senoprosopis*); Williston, 1891: 89 (cat.; *Senoprosopis*); Kertész, 1909: 284 (cat.; *Stenopropsis*); Bromley, 1946: 117 (cat.; *Senoprosopis*); Carrera, 1960a: 167 (n. comb.); Hull, 1962 (2): 500, figs. 334, 700, 1479, 2356 (*Senoprosopis*); Martin & Papavero, 1970: 61 (cat.).
- varipes** (Schiner), 1867: 405 (*Senoprosopis*). Type-locality: “Brazil”. Refs. – Williston, 1891: 89 (cat.; *Senoprosopis*); Kertész, 1909: 284 (cat.; *Stenopropsis*); Bromley, 1946: 117 (cat.; *Senoprosopis*); Hull, 1962 (2): 500, fig. 702 (*Senoprosopis*); Martin & Papavero, 1970: 61 (cat.). Type, NMW.

Genus *Stizolestes* Hull, 1962

Stizolestes Hull, 1962 (2): 560. Type-species, *Asilus nigriventris* Philippi, 1865 (orig. des.).

Ref. – Artigas, 1970a: 350 (key to spp.).

- atribarbis** Artigas, 1970a: 352, figs. 389, 459. Type-locality: Chile, Arauco, Pichinahuel, Cordillera Nahuelbuta. Distr. – Chile (Arauco, Malleco). Type, DZUC.
- aureomaculatus** (Bromley), 1932: 274, fig. 25 (*Asilus*). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. – Chile (Arauco, Cautín, Chiloé, Concepción, Linares, Llanquihue, Malleco, Osorno, Valdivia). Refs. – Stuardo Ortiz, 1946: 84 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 353, figs. 385, 386, 461 (n. comb.), 1971: 49, figs. 122-123. Type, BMNH.
- eritrichus** (Philippi), 1865: 698 (*Asilus*). Type-locality: Chile. Neotype-locality: Chile, Linares, Fundo Malcho, Cordillera Parral. Distr. – Chile (Linares). Refs. – Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 256 (cat.; *Asilus*); Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 355, figs. 387-388, 468 (n. comb.). Type, lost. NT DZUC (des. by Artigas, 1970a: 355).
- gayi** (Macquart), 1838: 148 (1839: 264) (*Asilus*). Type-locality: “Chile”. Distr. – Chile (Arauco, Cautín, Concepción, Curicó, Linares, Llanquihue, Malleco, Maule, Ñuble, Santiago, Talca, Valdivia). Refs. – Macquart, 1849: 95 (*Asilus*); Blanchard, 1852: 370 [misprinted as 270], pl. 2, figs. 12, 12a (*Asilus*); Walker, 1855: 704 (*Asilus*); Philippi, 1865: 695 (*Asilus*); Williston, 1891: 89 (cat.; *Asilus*); Kertész, 1909: 277 (cat.; *Dysmachus*); Bromley, 1932: 273 (*Asilus*); Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Artigas, 1970a: 356, figs. 391, 394, 501 (n. comb.), 1971: 49, figs. 126-127. Type, MNHNP.
- valdivianus** Philippi, 1865: 696 (*Asilus*). Type-locality: Chile, Valdivia. Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 265 (cat.; *Asilus*); Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 546 (*Asilus*). Type, lost.
- incomptus** Philippi, 1865: 696 (*Asilus*). Type-locality: Chile, Valdivia. Refs. – Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 257 (cat.; *Asilus*); Bromley, 1932: 281 (*Asilus*); Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*). Type, lost.
- mayi** (Edwards), 1932: 275 (*Asilus*). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. – Chile (Arauco, Aysén), Argentina (Río Negro). Refs. – Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 359, figs. 393, 444 (n. comb.), 1971: 50, figs. 124-125. Type, BMNH.
- modellus** (Bromley), 1932: 275 (*Asilus*). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. – Chile (Arauco, Cautín, Concepción, Curicó, Linares, Llanquihue, Malleco, Ñuble, Osorno, Valdivia), Argentina (Río Negro). Refs. – Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 360, figs. 390, 499 (n. comb.), 1971: 50, figs. 128-129. Type, BMNH.
- nigriventris** (Philippi), 1865: 966 (*Asilus*). Type-locality: Chile. Neotype-locality: Chile, Cautín, Temuco. Cerro Nielol. Distr. – Chile (Aconcagua, Arauco, Aysén, Biobío, Cautín, Chiloé, Concepción, Coquimbo, Curicó, Linares, Llanquihue, Malleco, Ñuble, O’ Higgins, Osorno, Santiago, Talca, Valdivia, Valparaíso). Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 261 (cat.; *Asilus*); Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 561, figs. 310, 773, 1528, 1537, 2177, 2211, 2351, 2407, 2410 (n. comb.); Artigas, 1970a: 362, figs. 392, 395-398, 503, 1971: 51, figs. 130-131; Artigas & Reyes, 1995: 201, figs. 1-2, 14-15. Type, lost. NT DZUC (des. by Artigas, 1970a: 362).

pamponeroides (Edwards), 1932: 277, fig. 26 (*Asilus*). Type-locality: Chile, Chiloé, Ancud. Distr. – Chile (Arauco, Chiloé, Concepción, Valdivia, Valparaíso). Ref. – Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 367, figs. 399, 502 (n. comb.), 1971: 51, figs. 132–133. Type, BMNH.
nigriventris Philippi of Hull, 1962 (2): 560 (*Asilus*), 561, figs. 210, 773, 1528, 1537, 2177, 2211, 2351, 2407, 2410, misident.

Genus **Threnia** Schiner, 1866

Threnia Schiner, 1866: 667 (in key), 674 (1868: 194; second erection of genus). Type-species, *Asilus carbonarius* Wiedemann, 1828 (orig. des.).

carbonaria (Wiedemann), 1828: 454 (*Asilus*). Type-locality: “Brazil” [Sello; cf. Papavero, 1971e: 56–60, 69–80, 99–102]. Distr. – Bolivia (Sara), Brazil (Rio de Janeiro, São Paulo). Refs. – Walker, 1855: 705 (*Asilus*); Schiner, 1866: 685, 1868: 74, pl. 2, fig. 11; Williston, 1891: 88 (cat.); Kertész, 1909: 272 (cat.); Carrera, 1945a: 165 (prey); Bromley, 1946: 116 (cat.); Carrera, 1952b: 240, figs. 1–6, 1958a: 150, 1960a: 165; Carrera & Vulcano, 1961: 76 (prey); Hull, 1962 (2): 578, figs. 311, 777, 1476, 1485, 2220, 2450, 2452, 2502; Martin & Papavero, 1970: 87 (cat.). Type, BZM [no. 373; cf. Schumann, 1973: 95].

kelleri Carrera, 1952b: 243, figs. 11–15. Type-locality: Brazil, São Paulo, Porto Cabral, rio Paraná. Distr. – Brazil (Minas Gerais, São Paulo). Refs. – Carrera & Vulcano, 1961: 76 (prey); Hull, 1962 (2): 578; Martin & Papavero, 1970: 61 (cat.). Type, MZUSP.

longipennis Schiner, 1868: 185. Type-locality: “South America”. Refs. – Williston, 1891: 88 (cat.); Kertész, 1909: 272 (cat.); Carrera, 1952b: 250; Hull, 1962 (2): 578; Martin & Papavero, 1970: 61 (cat.). Type, NMW.

lugens Schiner, 1868: 185. Type-locality: “South America”. Refs. – Williston, 1891: 88 (cat.); Kertész, 1909: 272 (cat.); Carrera, 1952b: 250; Hull, 1962 (2): 578; Martin & Papavero, 1970: 61 (cat.). Type, NMW.

rabellooi Carrera, 1952b: 242, figs. 7–10. Type-locality: Brazil, São Paulo, São Bento do Sapucaí. Distr. – Brazil (Rio de Janeiro, São Paulo, Santa Catarina). Refs. – Carrera, 1960a: 165; Carrera & Vulcano, 1961: 76 (prey); Hull, 1962 (2): 578; Martin & Papavero, 1970: 61 (cat.). Type, MZUSP.

therimachus (Walker), 1851: 137 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 705 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 264 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 546 (*Asilus*).

Genus **Wyliea** Martin, 1975

Wyliea Martin, 1975c: 78. Type-species, *Asilus mydas* Brauer, 1885 (orig. des.).

Ref. – Martin, 1975c: 79 (key to spp.).

chrysaugea (Osten Sacken), 1887: 208 (*Asilus*). Type-locality: Guatemala, Baja Verapaz, Panima. Distr. – Mexico (Oaxaca), Guatemala (Bava Verapaz). Refs. – Aldrich, 1905: 282 (cat.; *Asilus*); Kertész, 1909: 253 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 88 (cat.; unplaced); Martin, 1975c: 79 (in key; n. comb.); Fisher & Hespenheide, 1992: 630 (note 8). Type, BMNH.

mydas (Brauer), 1885: 387, pl., figs. 1, 1a (*Asilus*). Type-locality: “Mexico”. Distr. – U. S. A. (Arizona, New Mexico), Mexico (Durango, Morelos). Refs. – Kertész, 1909: 261 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Wilcox, 1965: 392 (cat.; *Asilus*); Martin & Papavero, 1970: 90 (cat.; unplaced); Martin, 1975c: 79 (in key), figs. 74–76 (n. comb.); Fisher & Hespenheide, 1992: 630 (note 8). Type, NMW.

midas Osten Sacken, 1887: 209 (*Asilus*); Williston, 1893a: 69 (*Asilus*); Aldrich, 1905: 282 (cat.; *Asilus*), Hine, 1909: 165 (*Asilus*), error or emend.

Efferia-group

Genus **Albibarbefferia** Artigas & Papavero, 1997

Albibarbefferia Artigas & Papavero, 1997b: 68. Type-species, *Erax abibarbis* Macquart, 1838 (orig. des.).

Erax, barbatus-group of Hine, 1919: 108.

Efferia, albobarbis-group of Wilcox, 1966c: 104.

- albibarbis** (Macquart), 1838: 118 (1839: 234) (*Erax*). Type-locality: "North America". Neotype-locality: U. S. A., Ohio, Paint Creek, Bainbridge. Distr. – North America south into Guatemala. Refs. – Walker, 1855: 618 (*Erax*); Schiner, 1867: 395 (*Erax*); Kertész, 1909: 235 (cat.; *Erax*; as syn. of *cinerascens* Bellardi, 1861); Hine, 1918: 3 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 393 (cat.; *Efferia*); Wilcox, 1966c: 108 (*Efferia*); Martin & Papavero, 1970: 62 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.). Type, MNHNP (only legs and wings remaining). NT USNM [des. by Martin, 1962b: 252].
- pumilus** Walker, 1855: 640 (*Erax*; preocc. Macquart, 1849). Type-locality: Mexico, Veracruz. Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*).
- cinerascens** Bellardi, 1861: 139 (39), pl. 2, fig. 10 (*Erax*). Type-locality: Mexico, Veracruz: Tuxpango, near Orizaba. Refs. – Schiner, 1867: 394 (*Erax*); Osten Sacken, 1887: 202 (*Erax*); Cockerell, 1894: 173 (*Erax*); Townsend, 1895: 599 (*Erax*); Williston, 1901: 323 (*Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*). Type, TORO.
- tricolor** Bellardi, 1861: 140 (40), pl. 2, fig. 12 (*Erax*). Type-locality: Mexico, surroundings of Mexico City. Refs. – Townsend, 1895: 599 (*Erax*); Aldrich, 1905: 278 (cat.; *Erax*); Kertész, 1909: 243 (cat.; *Erax*); Engel, 1930: 460 (*Erax*); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- furax** Williston, 1885: 67 (*Erax*). Type-locality: U. S. A., "Washington Territory", California, western Kansas, Arizona, Connecticut. Syntypes, KU.
- cinerescens** Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.), error.
- armata** (Hine), 1918: 4 (*Erax*). Type-locality: U. S. A., Texas, Davis Mountains, Fort Davis Quadrangle, Phantom Lake. Distr. – U. S. A. (Texas), Mexico (Coahuila, Nuevo León, Tamaulipas, Veracruz). Refs. – Hine, 1919: 112 (*Erax*); Baker, 1939: 169 (*Erax*); Carrera & d'Andretta, 1950: 172, figs. 12, 28 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 109 (*Efferia*). Type, OSU.
- bicolor** (Bellardi), 1861: 147 (47) (*Erax*). Type-locality: Mexico, surroundings of Mexico City. Distr. – U. S. A. (Arizona, New Mexico, Texas, Mississippi), Mexico (Distrito Federal, Michoacán). Refs. – Aldrich, 1905: 85 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Hine, 1919: 110 (*Erax*); Bromley, 1935: 4 (*Erax*); Carrera & d'Andretta, 1950: 172 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 111 (*Efferia*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- bimaculata** (Bellardi), 1861: 145 (45), pl. 2, fig. 11 (*Erax*). Type-locality: "Mexico". Distr. – Mexico (Jalisco, Morelos, Nayarit, Veracruz). Refs. – Schiner, 1868: 182 (*Erax*); Williston, 1891: 85 (cat.; *Erax*), 1901: 325 (*Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Hine, 1919: 115 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- cingulata** (Bellardi), 1861: 142 (42) (*Erax*). Type-locality: Mexico, Morelos, Cuautla. Distr. – Mexico (Michoacán, Morelos). Refs. – Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hine, 1919: 111, fig. 34 (*Erax*); Carrera & d'Andretta, 1950: 175 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- duncani** (Wilcox), 1966c: 112 (*Efferia*). Type-locality: U. S. A., Arizona, San Carlos Lake. Distr. – U. S. A. (Arizona), Mexico. Type, CAS.
- eximia** (Bellardi), 1861: 138 (38) (*Erax*). Type-locality: Mexico, surroundings of Mexico City. Distr. – Mexico (D. F., Morelos). Refs. – Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 236 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- eximeus** Hine, 1919: 146, fig. 46 (*Erax*), error.
- grandis** (Hine), 1919: 111 (*Erax*). Type-locality: U. S. A., Texas, Round Mountains. Distr. – U. S. A., Mexico (Coahuila, Nuevo León, Tamaulipas). Refs. – Baker, 1939: 169 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 113 (*Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.). Syntypes, OSU.
- ?**heteroptera** (Macquart), 1846: 211 (1846: 83), pl. 8, fig. 6 (*Erax*). Type-locality: "Colombia" (as "Nouvelle Grenade", cold region). Refs. – Walker, 1855: 634 (*Erax*); Schiner, 1868: 182 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 65 (cat.; *Efferia*, as *heteropterus*, error); Artigas & Papavero, 1997b: 68 (n. comb.). Type, OXF.
- incognita** (Forbes), 1987: 296, figs. 1, 3, 7 (*Efferia*). Type-locality: U. S. A., Texas, Davis Mountains, south side of Route 166, 6.3 miles southwest of junction with route 118, 1790 m. Distr. – U. S. A. (New Mexico, Texas), Mexico (Chihuahua). Type, CAS.
- marginata** (Bellardi), 1861: 146 (46) (*Erax*). Type-locality: Mexico, Morelos, Cuatla. Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 239 (cat.; *Erax*); Martin & Papavero, 1970: 66 (cat.; *Erax*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- peralta** (Wilcox), 1966c: 115 (*Efferia*). Type-locality: U. S. A., California, Orange County, Peralta (4 miles east of Olive). Distr. – U. S. A. (California), Mexico. Type, CAS.
- quadrimaculata** (Bellardi), 1861: 144 (44), pl. 2, fig. 13 (*Erax*). Type-locality: Mexico, Veracruz: Playa Vicente and Córdoba. Distr. – Mexico (Veracruz) Refs. – Williston, 1901: 325 (*Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Hine, 1919: 114 (*Erax*; ?specimens from Colombia); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 67 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican

- dipterol). Syntypes, TORO.
- sagax** (Williston), 1901: 324 (*Erax*). Type-locality: Mexico, Jalisco, Santiago Ixcuintla. Distr. – Mexico (Jalisco). Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 242 (cat.; *Erax*); Hine, 1919: 113 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 68 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.
- sonorensis** (Forbes), 1987: 298, figs. 2, 4, 8 (*Efferia*). Type-locality: Mexico, Sonora, 17 miles north of El Caballo (south of Hermosillo). Distr. – Mexico (Sinaloa, Sonora). Type, CAS.
- vertebrata** (Bromley), 1940: 14 (*Erax*). Type-locality: U. S. A., California, Warner Springs. Distr. – U. S. A. (California, Arizona), Mexico. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 396 (cat.; *Efferia*); Wilcox, 1966c: 108 (*Efferia*).

Genus *Carinefferia* Artigas & Papavero, 1997

- Carinefferia** Artigas & Papavero, 1997b: 69. Type-species, *Erax carinatus* Bellardi, 1861 (orig. des.).
- Erax, carinatus*-group of Hine, 1919: 131.
- Efferia, arida*-group of Wilcox, 1966c: 138.
- aper** (Walker), 1855: 621 (*Erax*). Type-locality: “Mexico”. Distr.- Mexico (Oaxaca), Guatemala. Refs. – Aldrich, 1905: 275 (cat.; *Erax*); Kertész, 1909: 233 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 69 (n. comb.); Papavero & Ibáñez Bernal, 2001: 95 (hist. Mexican dipterol.). Type, BMNH.
- comatus* Bellardi, 1861: 134 (34) (*Erax*). Type-locality: “Mexico”. Refs. – Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hull, 1962 (2): 478; Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.). Type, TORO.
- unicolor* Bellardi, 1861: 137 (37) (*Erax*). Type-locality: Mexico, Veracruz, Orizaba. Refs. – Wulp, 1882b: 114 (*Erax*); Aldrich, 1905: 278 (cat.; *Erax*); Kertész, 1909: 243 (cat.; *Erax*); Hine, 1919: 132, fig. 60 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.). Type, TORO.
- caliente** (Wilcox), 1966c: 153, figs. 109-110 (*Efferia*). Type-locality: U. S. A., California, San Diego County, Aguas Calientes. Distr. – U. S. A. (California). Type, CAS.
- carinata** (Bellardi), 1861: 136 (36), pl. 2, fig. 9 (*Erax*). Type-locality: “Mexico”. Distr. – Mexico (Guerrero). Refs. – Osten Sacken, 1887: 205 (*Erax*); Townsend, 1895: 598 (*Erax*); Williston, 1901: 323 (*Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Hine, 1919: 131, fig. 58 (*Erax*); Hull, 1962 (2): 478; Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 68 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.). Type, TORO.
- concinnata** (Williston), 1901: 323 (*Erax*). Type-locality: Mexico, Guerrero: Acaguizotla and Rincón. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hine, 1919: 131, fig. 57 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 70 (n. comb.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Syntypes, BMNH.
- cressoni** (Hine), 1919: 134, fig. 50 (*Erax*). Type-locality: U. S. A., New Mexico, Alamogordo. Distr. – U. S. A. (New Mexico, Texas), Mexico. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 157 (*Efferia*); Lavigne & Dennis, 1985: 146 (*Efferia*; ethology); Dennis, Lavigne & Bullington, 1986: 42 (*Efferia*). Type, ANSP.
- latruncula** (Williston), 1885: 67 (*Erax*). Type-locality: U. S. A., Montana and Arizona. Distr. – U. S. A. (Arizona, Utah), Mexico. Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Hine, 1919: 132, fig. 56 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 395 (cat.; *Efferia*); Wilcox, 1966c: 159 (*Efferia*). Type, KU.
- parvula** (Bellardi), 1861: 135 (35), pl. 2, fig. 8 (*Erax*). Type-locality: “Mexico”. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Hine, 1919: 132, fig. 55 (*Erax*); Martin & Papavero, 1970: 67 (cat.; *Efferia*); Artigas & Papavero, 1997b: 70 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.). Type, TORO.
- prolifica** (Osten Sacken), 1887: 202 (*Erax*). Type-locality: Mexico, northern Sonora. Distr. – U. S. A. (Arizona, New Mexico), Mexico (Sonora). Refs. – Williston, 1891: 86 (cat.; *Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Wilcox, 1966c: 160 (*Efferia*); Martin & Papavero, 1970: 67 (cat.; *Efferia*); Artigas & Papavero, 1997b: 70 (n. comb.); Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.
- willistoni** (Hine), 1919: 110 (*Erax*). Type-locality: U. S. A., Arizona, Williams. Distr. – U. S. A. (Arizona, Colorado, New Mexico), Mexico (Chihuahua). Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 396 (cat.; *Efferia*); Wilcox, 1966c: 162 (*Efferia*). Type, USNM.

Genus *Cratolestes* Hull, 1962

- Cratolestes** Hull, 1962 (2): 483. Type-species, *Asilus spectabilis* Philippi, 1865 = *Cratolestes chilensis* (Macquart, 1950) (orig. des.).

chiliensis (Macquart), 1850: 389 (1850: 85), pl. 8, fig. 5 (*Erax*). Type-locality: “Chile”. Distr. – Chile (Acogagua, Antofagasta, Biobío, Concepción, Cautín, Coquimbo, Maule, Ñuble, Santiago, Valparaíso). Refs. – Blanchard, 1852: 369 (*Erax*); Walker, 1855: addenda; Philippi, 1865: 692 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas, 1970a: 265, figs. 227, 231, 439 (n. comb.); Martin & Papavero, 1970: 71 (cat.); Artigas, 1971: 32, figs. 77–78; Artigas & Papavero, 1997b: 70. Type, MNHNP.
spectabilis Philippi, 1865: 695 (*Asilus*). Type-locality: Chile: Santiago, and Choapa, Illapel. Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 264 (cat.; *Asilus*); Stuardo Ortiz, 1946: 85 (cat.; *Asilus*); Hull, 1962 (2): 484, figs. 360, 749, 1438, 1447, 2178, 2204, 2455, 2469. Syntypes, MNHNS.
albescens Schiner, 1868: 180 (*Erax*). Type-locality: “Chile”. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 233 (cat.; *Erax*); Stuardo Ortiz, 1946: 84 (cat.; *Lochmorhynchus*); Hull, 1962 (2): 478 (*Nerax*). Type, NMW.
perniger Schiner, 1868: 180 (*Erax*). Type-locality: unknown. Refs. – Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Stuardo Ortiz, 1946: 84 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*). Type, NMW.
wirthi Artigas, 1970a: 268, figs. 232–235, 421. Type-locality: Chile, Tarapacá, Arica, río Azapa. Distr. – Chile (Antofagasta, Coquimbo, Tarapacá). Refs. – Artigas, 1971: 33, figs. 75–76; Artigas & Papavero, 1997b: 70. Type, DZUC.

Genus **Diplosynapsis** Enderlein, 1914

Diplosynapsis Enderlein, 1914a: 259. Type-species, *Diplosynapsis argentifascia* Enderlein, 1914 (orig. des.).

argentifascia Enderlein, 1914a: 259, fig. 5. Type-locality: Peru, Junín, Chanchamayo, and Brazil, Pará. Distr. – Peru (Cuzco, Junín), Brazil (Pará). Refs. – Hull, 1962 (2): 479; Lamas M., 1972: 313; Artigas & Papavero, 1997b: 70. Syntypes, IZPAN.
argentifascia Bromley, 1946: 114 (cat.), error.
?cellata (Schiner), 1868: 181 (*Erax*). Type-locality: “South America” (Venezuela on ST’s labels). Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 70 (n. comb.). Syntypes, NMW.
halterata Enderlein, 1914a: 260. Type-locality: Colombia, Caldas, Victoria (Hacienda Pehlke), 900 m [cf. Papavero, 1973d: 299]. Distr. – Colombia (Caldas). Refs. – Hull, 1962 (2): 479; Artigas & Papavero, 1997b: 70. Type, IZPAN.
remus Tomasovic, 2002b: 60, fig. 19. Type-locality: French Guiana, Pointe Combi. Distr. – French Guiana. Type, FUSAGembloix.

Genus **Efferia** Coquillett, 1893

Efferia Coquillett, 1893d: 175. Type-species, *Efferia candida* Coquillett, 1893 (sub. des., Coquillett, 1910: 536).

Erax Scopoli of authors (part), misident.

Erax, anomalus-group of Hine, 1919: 119.

Erax, anomala-group of Wilcox, 1966c: 120.

alia Scarbrough & Perez-Gelabert, 2009: 8, figs. 1–4. Type-locality: Dominican Republic, Pedernales, 25 km n. of Cabo Rojo. Distr. – Dominican Republic. Type, USNM.
angusta Scarbrough & Perez-Gelabert, 2009: 11, figs. 5–8. Type-locality: Dominican Republic, Azua, RD-156 La Furnia, Barreras. Distr. – Dominican Republic. Type, USNM.
anomala (Bellardi), 1861: 132 (32), pl. 2, fig. 7 (*Erax*). Type-locality: Mexico, Morelos, Cuautla. Distr. – U. S. A. (Arizona, Kansas to Texas), Mexico. Refs. – Williston, 1885: 69 (*Erax*); Osten Sacken, 1887: 199 (*Erax*); Coquillett, 1893d: 175; Williston, 1894: 137 (*Eichoichemus*), 1901: 323 (*Erax*); Aldrich, 1905: 275 (*Erax*); Kertész, 1909: 244 (cat.; *Eichoichemus*); Hine, 1918: 3 (*Erax*), 1919: 119, figs. 5, 10 (*Erax*); Bromley, 1935c: 4 (*Erax*); Hull, 1962 (2): 475; Martin & Wilcox, 1965: 394 (cat.; n. comb.); Wilcox, 1966c: 126 (*Efferia*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.). Type, TORO.
argentifrons (Hine), 1911b: 308, fig. 35 (*Erax*). Type-locality: U. S. A., Kansas, Clark County. Distr. – U. S. A. (Kansas), Mexico (Coahuila). Refs. – Bromley, 1935c: 4 (*Erax*); Baker, 1939: 169 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 393 (cat.; n. comb.). Type, OSU.
belfragei (Hine), 1919: 121, fig. 21 (*Erax*). Type-locality: U. S. A., Texas. Distr. – U. S. A. (Texas), Mexico (Coahuila). Refs. – Baker, 1939: 168 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; n. comb.); Wilcox, 1966c: 101.
bullata Scarbrough & Perez-Gelabert, 2009: 13, figs. 9–14. Type-locality: Dominican Republic, Pedernales, PN Jaragua, trail to Carlitos, ca. km s. of Highway 44, 106 m. Distr. – Dominican Republic (Barahona, Isla Beata, Pedernales). Type, USNM.
candida Coquillett, 1893d: 176. Type-locality: U. S. A., California, Los Angeles Co. Distr. – U. S. A. (California, Arizona, New Mexico, Kansas), Mexico (Coahuila). Refs. – Williston, 1894: 137 (*Eichoichemus*); Aldrich, 1905: 276 (*Erax*); Kertész,

- 1909: 244 (cat.; *Eichoichemus*); Hine, 1919: 119, fig. 11 (*Erax*); Bromley, 1935c: 4 (*Erax*); Baker, 1939: 168 (*Erax*); Hull, 1962 (2): 475; Martin & Wilcox, 1965: 394 (cat.; n. comb.); Martin & Wilcox, 1965: 394 (cat.); Wilcox, 1966c: 130; Artigas & Papavero, 1997b: 71 (n. comb.).
- caymanensis** Scarbrough, 1988a: 151, figs. 1-2. Type-locality: Grand Cayman Island. Distr. – Great Cayman Island. Type, USNM.
- clava** Scarbrough & Perez-Gelabert, 2009: 15, figs. 15-20. Type-locality: Dominican Republic, Azua, RD-065 Monte Rio, 4.5 km w. of Beach. Distr. – Dominican Republic (Azua, Barahona, Independencia, Pedernales, San Juan). Type, USNM.
- davisi** Wilcox, 1966c: 132. Type-locality: U. S. A., Arizona, 12 miles north of Sasabe. Distr. – U. S. A. (Arizona, New Mexico, Nevada), Mexico. Type, CAS.
- exacta** Scarbrough & Perez-Gelabert, 2009: 18, figs. 21-26. Type-locality: Dominican Republic, Santiago, RD-248 entrance to Sabana, Iglesia. Distr. – Dominican Republic (Santiago, Santo Domingo), Haiti (Ouerst department). Type, USNM.
- forbesi** (Curran), 1931a: 10, fig. 3 (*Erax*). Type-locality: Puerto Rico, Coamo Springs. Puerto Rico, British Virgin Islands (Guana island). Refs. – Hull, 1962 (2): 479 (*Nerax*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71; Scarbrough & Perez-Gelabert, 2009: 47, figs. 90, 97-102. Type, AMNH.
- fortis** (Walker), 1855: 623 (*Erax*). Type-locality: Dominican Republic, Santo Domingo. Distr. – Dominican Republic (Azua, Barahona, Independencia, Monte Cristi, Peravia, San Juan, Santiago), Haiti (Ouest department). Refs. – Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 236 (cat.; *Erax*); Hull, 1962 (2): 479 (n. comb.; as syn. of *stylatus* Fabricius, 1775); Martin & Papavero, 1970: 65 (cat.; n. comb.); Artigas & Papavero, 1997b: 71 (*Nerax*); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.; *Erax*); Scarbrough & Perez-Gelabert, 2006: 7, 2009: 20, figs. 27-32, 91. Type, BMNH.
- rufitibia** Macquart, 1848: 187 (1848: 27) (*Erax*), part (only his Haiti specimen). Ref. – Hine, 1919: 128, fig. 30 (*Erax*; Bahamas, Cuba); Curran, 1931a: 9 (*Erax*; as syn. of *stylata* (Fabricius)), 1953: 4 (*Erax*); Hull, 1962 (2): 479 (*Nerax*; n. comb.); Martin & Papavero, 1970: 56 (cat.; *Efferia*, n. comb.). Type, OXF.
- flavotibia** Bigot, 1878: 329 (*Phoneus*). Type-locality: “Haiti”. Refs. – Kertész, 1909: 171 (cat.; *Neophoneus*); Hull, 1962 (1): 354 (*Neophoneus*); Martin & Papavero, 1970: 45 (cat.; *Neophoneus*); Scarbrough & Perez-Gelabert, 2006: 20 (n. comb.), 2009: 20 (n. syn.). Type, OXF.
- flavitibialis** Bigot, 1879: xxi (*Phoneus*), emend.
- flavipes** Bigot, 1880: xl (*Phoneus*), error.
- flavitibius** Bigot, 1880: xl (*Phoneus*), emend.
- flavitibia** Aldrich, 1905: 273 (cat.), error.
- fulvibarbis** (Macquart), 1848: 188 (1848: 28), pl. 2, fig. 13 (*Erax*). Type-locality: “Haiti”. Refs. – Walker, 1855: 624 (*Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Hine, 1919: 129, fig. 23 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 65 (cat.; n. comb.); Artigas & Papavero, 1997b: 72 (*Nerax*); Scarbrough & Perez-Gelabert, 2006: 8, 2009: 23, figs. 33-38. Type, lost.
- pachychaeta** Bromley, 1928: 1 (*Erax*). Type-locality: Haiti, La Molinière. Distr. – Haiti. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 66 (cat.; n. comb.); Artigas & Papavero, 1997b: 72 (*Nerax*); Scarbrough & Perez-Gelabert, 2006: 8, 2009: 23 (n. syn.). Type, AMNH.
- haitensis** (Macquart), 1848: 188 (1848: 28) (*Erax*). Type-locality: “Haiti”. Distr. – Dominican Republic (Azua, Peravia), Haiti (Ouest department). Refs. – Walker, 1855: 623 (*Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Curran, 1928b: 23 (*Erax*), 1931a: 9 (*Erax*); Scarbrough & Perez-Gelabert, 2009: 26, figs. 39-44, 94 (revalidation of sp.). Type, OXF.
- imperialis** Forbes, 1988: 556, figs. 1a, 2c. Type-locality: U. S. A., California, Imperial County, Black Mountain, 4.5 miles southeast of Route 78, 390-454m. Distr. – U. S. A. (California, Arizona), Mexico (Sonora). Type, CAS.
- incisura** Scarbrough & Perez-Gelabert, 2009: 29, figs. 46-51. Type-locality: Dominican Republic, Isla Beata, RD-079, trail to Cueva del Tubo, PN. Distr. – Dominican Republic (Isla Beata). Type, USNM.
- montensis** Scarbrough & Perez, 2009: 49, figs. 95, 103-108. Type-locality: Puerto Rico, Utuado. Distr. – Puerto Rico. Type, AMNH.
- nigrimystacea** (Macquart), 1847: 57 (1847: 41) (*Erax*). Type-locality: “Guadeloupe”. Distr. – Guadeloupe, Barbuda, Dominica, St. Lucia, Montserrat, Redonda, St. Vincent. Refs. – Walker, 1855: 633 (*Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Hine, 1919: 130, fig. 31 (*Erax*); Hull, 1962 (2): 475 (*Asilus*); Martin & Papavero, 1970: 70 (cat.; *Eicherax*); Scarbrough & Knutson, 1989: 523, fig. 3 (*Efferia*, n. comb.); Artigas & Papavero, 1997b: 72 (*Nerax*), 2009: 52, figs. 96, 110-115. Type, OXF.
- picea** Scarbrough & Perez-Gelabert, 2009: 31, figs. 52-57. Type-locality: Dominican Republic, La Vega, P. N. Armando Bermúdez, trail La Ciénega de Manabao-Los Tablones. Distr. – Dominican Republic (La Vega). Type, USNM.
- pilosa** (Hine), 1919: 150 (*Erax*). Type-locality: U. S. A., Texas. Distr. – U. S. A. (Texas), Mexico (Coahuila). Refs. – Baker, 1939: 168 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 395 (cat.; n. comb.).
- portoricensis** (Hine), 1919: 128, fig. 24 (*Erax*). Type-locality: Puerto Rico, Ensenada. Distr. – Puerto Rico. Refs. – Bromley, 1929a: 290 (*Erax*); Curran, 1931a: 9 (*Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.; *Erax*); Artigas & Papavero, 1997b: 72; Scarbrough & Perez-Gelabert, 2009: 56, figs. 116-121. Type, AMNH.
- serrula** Scarbrough & Perez-Gelabert, 2009: 33, figs. 58-63. Type-locality: Dominican Republic, Santiago, RD-243, Babosico, on way to Jánico. Distr. – Dominican Republic (Independencia, La Altagracia, Monte cristi, San Juan, Santiago). Type, USNM.
- sinuosa** Scarbrough & Perez-Gelabert, 2009: 35, figs. 64-69, 96. Type-locality: Dominican Republic, Peravia, RD-171, El Maizal, near Presa de Valdesia. Distr. – Dominican Republic (Independencia, Peravia, San Juan, Santiago). Type, USNM.
- sp. C** Scarbrough & Perez-Gelabert, 2009: 63, figs. 130-131. Tobago (Canoe Bay rd. scrub forest nr. Hilton Resort), Trinidad

(Arima Valley).

spinulosa Scarbrough & Perez-Gelabert, 2009: 38, figs. 70-75. Type-locality: Dominican Republic, Independencia, terrenos Zona Franca. Distr. – Dominican Republic (Barahona, Independencia, Peravia, San Juan). Type, USNM.

stylata (Fabricius), 1775: 795 (*Asilus*). Type-locality: “America”. Distr. – U. S. A. (extreme southern Florida), Cuba, Haiti, Puerto Rico, Santo Domingo, Virgin Islands (St. Croix, St. Thomas Islands), Hispaniola. Refs. – Fabricius, 1781: 464 (*Asilus*), 1787: 360 (*Asilus*); Olivier, 1789: 268 (*Asilus*); Gmelin, 1792: 2899 (*Asilus*); Fabricius, 1794: 384 (*Asilus*), 1805: 171 (*Dasypogon*); Wiedemann, 1821: 198 (*Asilus*), 1828: 462, pl. 6, fig. 6 (*Asilus*); Walker, 1849: 399 (*Asilus*), 1855: 638 (*Erax*); Schiner, 1866: 686 (*Erax*); Wulp, 1882b: 112 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 243 (cat.; *Erax*); Hine, 1919: 127, fig. 32 (*Erax*); Bromley, 1929a: 291, pl. 1, figs. 1, 4 (*Erax*); Curran, 1931a: 9 (*Erax*); Carrera & d’Andretta, 1950: 174 (*Erax*, misident.); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 68 (cat.; n. comb.); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.; *Erax*); Artigas & Papavero, 1997b: 72 (*Nerax*); Scarbrough & Perez-Gelabert, 2006: 8, 2009: 58, figs. 122-127. Type, KIEL [cf. Zimsen, 1964: 471].

suspiciosa Scarbrough & Perez-Gelabert, 2009: 40, figs. 76-81. Type-locality: Dominican Republic, La Vega, P. N. Armando Bermúdez, km 1-3 along trail w. of La Ciénega, 900-1000 m. Distr. – Dominican Republic (La Vega). Type, USNM.

texana (Banks) in Hine, 1919: 151, fig. 43 (*Erax*). Type-locality: U. S. A., Texas. Distr. – U. S. A. (Texas), Mexico (Coahuila). Refs. – Bromley, 1935c: 4 (*Erax*); Baker, 1939: 169 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 396 (cat.; n. comb.); Morgan & Shelly, 1988: 419 (body temperature regulation). Type, OSU.

tricella (Bromley), 1951: 32 (*Erax*). Type-locality: U. S. A., Arizona, Buckeye. Distr. – U. S. A. (Arizona, New Mexico), Mexico (Sonora). Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 396 (cat.; n. comb.); Wilcox, 1966c: 137. Type, USNM.

tuberculata (Coquillett), 1904b: 34 (*Erax*). Type-locality: U. S. A., Texas. Distr. – U. S. A. (California, Texas), Mexico (Coahuila). Refs. – Hine, 1918: 4 (*Erax*); Baker, 1939: 169 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; n. comb.). Type, USNM.

woodleyi Scarbough & Perez-Gelabert, 2009: 42, figs. 82-87. Type-locality: Dominican Republic, Independencia, P. N. Sierra de Bahoruco, Caseta 1 w. of Puerto Escondido, 1215 m. Distr. – Dominican Republic (Azua, Independencia, La Altagracia, La Romana, La Vega, San Juan, Santiago, Santo Domingo). Type, USNM.

Genus *Nerax* Hull, 1962

Nerax Hull, 1962 (2): 476. Type-species, *Asilus aestuans* Linnaeus, 1758 (orig. des.).

Erax Scopoli of authors (part), misident.

Efferia Coquillett of authors, misident.

Erax, aestuans-group of Hine, 1919: 120.

Efferia, aestuans-group of Wilcox, 1966c: 94.

Some of the species listed below may be belong to several different genera, other than *Nerax*; only examination of the types will help to clarify this group.

abdominalis (Wiedemann), 1821: 432 (*Asilus*). Type-locality: “South America” (Brazil, São Paulo, Ipanema in label of ST).

Refs. – Wiedemann, 1828: 432 (*Asilus*); Walker, 1855: 701 (*Asilus*); Schiner, 1866: 683 (*Erax*); Williston, 1891: 84 (cat.; *Erax*); Kertész, 1909: 232 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 62 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71 (n. comb.). Syntypes, NMW.

affinis (Bellardi), 1861: 141 (41) (*Erax*). Type-locality: Mexico, Veracruz, Córdoba. Distr. – Mexico (Morelos, Veracruz).

Refs. – Aldrich, 1905: 275 (cat.; *Asilus*); Kertész, 1909: 233 (cat.; *Erax*); Hine, 1919: 121, fig. 18 (*Erax*; ?specimens from Colombia); Hull, 1962 (2): 478 (n. comb.); Martin, 1965a: 29 (*Efferia*); Martin & Papavero, 1970: 62 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71; Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.

amazonicus (Bromley), 1934b: 359 (*Erax*). Type-locality: Brazil, Minas Gerais, Curralinho. Distr. – ?Peru, Brazil (Minas Gerais). Refs. – Bromley, 1946: 114 (cat.; *Erax*); Carrera & d’Andretta, 1953: 70; Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 62 (cat.; *Efferia*); ?Lamas M., 1972: 313 (*Efferia*); Artigas & Papavero, 1997: 71. Type, COR.

apicalis (Wiedemann), 1821: 191 (*Asilus*). Type-locality: “North America”. Distr. – U. S. A., Mexico. Refs. – Wiedemann, 1828: 443 (*Asilus*); Walker, 1849: 453 (*Asilus*), 1855: 619 (*Erax*); Schiner, 1866: 684 (*Asilus*); Osten Sacken, 1878: 79, 294 (cat.; *Erax*); Comstock, 1879: 291 (*Erax*); Aldrich, 1905: 275 (cat.; *Erax*); Kertész, 1909: 233 (cat.; *Erax*); Hine, 1919: 125, fig. 18 (*Erax*); Fattig, 1945: 24; Bromley, 1950: 237 (*Erax*); Hull, 1962 (2): 478 (n. comb.); Wilcox, 1966c: 98 (*Efferia*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997: 71; Alayo Dalmau & Garcia Ávila, 1983: 48 (*Erax*).

auribarbis (Wiedemann), 1821: 186 (*Asilus*). Type-locality: Brazil, Bahia [Gomez; cf. Papavero, 1971e: 48]. Distr. – Brazil (Bahia). Refs. – Wiedemann, 1828: 437 (*Asilus*); Walker, 1855: 702 (*Erax*); Schiner, 1866: 683 (*Erax*); Wulp, 1882: 113 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Syntypes, BZM [no.

- 2050; cf. Schumann, 1973: 89], ?SENCK [cf. Geller-Grimm, 1999: 206].
- badiapex** (Bromley), 1928: 4 (*Erax*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, AMNH.
- bardyllis** (Walker), 1849: 401 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 637 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, BMNH.
- bilineatus** (Wulp), 1882b: 115 (*Erax*). Type-locality: “Argentina”. Refs. – Lynch Arribálzaga, 1883b: 15 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, ZMA.
- brunnescens** (Bromley), 1929a: 291, fig. 13, pl. 1, figs. 6, 9 (*Erax*). Type-locality: Cuba, Guantánamo. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Alayo Dalmau & Garcia Ávila, 1983: 48 (cat.; *Erax*); Artigas & Papavero, 1997b: 71. Type, MCZ.
- camposianus** (Curran), 1931b: 23 (*Erax*). Type-locality: Ecuador, Guayas, Guayaquil. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, AMNH.
- caudex** (Walker), 1849: 404 (*Asilus*). Type-locality: “Jamaica”. Refs. – Walker, 1855: 622 (*Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 234 (cat.; *Erax*); Hine, 1919: 127, fig. 32 (as syn. of *stylatus* Fabricius); Hull, 1962 (2): 478 (n. comb.; as syn. of *stylatus* Fabricius); Farr, 1965: 30, fig. 24 (*Efferia*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, BMNH.
- invariatus* Walker, 1851: 131 (*Asilus*). Type-locality: “Jamaica”. Refs. – Walker, 1855: 623 (*Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Farr, 1965: 30 (*Efferia*; syn.). Type, BMNH.
- cazieri** (Curran), 1953: 5 (*Erax*). Type-locality: Bahama Islands, South Bimini Island. Refs. – Martin, 1957b: 7; Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, AMNH.
- chapadensis** (Bromley), 1928: 3 (*Erax*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*). Type, AMNH.
- cockerellorum** (James), 1953: 50, fig. 3 (*Erax*). Type-locality: Honduras, Morelos, Uroca. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, WASH.
- commiles** (Walker), 1851: 132 (*Erax*). Type-locality: “South America”. Refs. – Walker, 1855: 633 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, BMNH.
- cubensis** (Bromley), 1929a: 292, fig. 14 (*Erax*). Type-locality: Cuba, Soledad. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71; Alayo Dalmau & Garcia Ávila, 1983: 48 (cat.; *Erax*). Type, MCZ.
- eurylabis** (Wiedemann), 1828: 469 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 636 (*Erax*); Schiner, 1866: 687 (*Erax*); Williston, 1891: 90 (cat.; ?*Erax*); Kertész, 1909: 236 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, BZM [no. 2035; cf. Schumann, 1973: 108].
- flavofasciatus** (Wiedemann), 1828: 470 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1849: 400 (*Asilus*; misident.; cf. Martin, 1965a: 29), 1855: 636 (*Erax*); Schiner, 1866: 687 (*Asilus*); Wulp, 1882b: 113 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 236 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin, 1965a: 29 (*Efferia*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71. Type, BZM [no. 2039; cf. Schumann, 1973: 111].
- fuscanipennis** (Macquart), 1850: 390 (1850: 86) (*Erax*). Type-locality: “Guiana”. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, OXF.
- fuscus** (Wiedemann), 1828: 465 (*Erax*). Type-locality: “Brazil” [Olfers; cf. Papavero, 1971e: 80]. Refs. – Macquart, 1838: 112 (*Erax*); Walker, 1849: 398 (*Asilus*), 1855: 637 (*Erax*); Schiner, 1866: 686 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Bromley, 1946: 117 (cat.; *Pachychoeta*); Carrera & d’Andretta, 1950: 175 (*Erax*, misident.); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, BZM [no. 2027; cf. Schumann, 1973: 113].
- gossei** (Farr), 1965: 31, fig. 25 (*Efferia*). Type-locality: Jamaica, St. Thomas, Morant Point. Refs. – Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Holotype, BMNH.
- haloesus** (Walker), 1849: 405 (*Asilus*). Type-locality: “Jamaica”. Refs. – Walker, 1855: 624 (*Erax*); Jonson, 1894: 274 (*Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Hine, 1919: 130, fig. 36 (*Erax*); Johnson, 1919: 436, fig. 36 (*Erax*); Hull, 1962 (2): 478 (n. comb.); Farr, 1965: 28, fig. 23 (*Efferia*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, BMNH.
- jalaeosus* Gowdey, 1926: 77 (*Erax*), error.
- hubbelli** (James), 1953: 51, fig. 4 (*Erax*). Type-locality: Honduras, Morazán, Mt. Calucatepe, 4200-4500 ft. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, MICH.
- imbuda** (Curran), 1934a: 16, fig. 1 (*Erax*). Type-locality: Ecuador, Guayas, Golfo de Guayaquil, Isla Puná, Puerto Grande. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, AMNH.

- labidophorus** (Wiedemann), 1828: 459 (*Asilus*). Type-locality: “Brazil” [Sello; cf. Papavero, 1971e: 56-60, 69-80, 88-102]. Refs. – Macquart, 1838: 110 (*Erax*); Walker, 1849: 398 (*Erax*), 1855: 637 (*Erax*); Schiner, 1866: 686 (*Erax*); Wulp, 1882b: 113 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Carrera, 1960a: 165 (*Erax*); Carrera & Vulcano, 1961: 74 (*Erax*; prey); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Syntypes, BZM [no. 2029; cf. Schumann, 1973: 122].
labidiphorus Tomasovic, 2002b: 55, fig. 8 (*Efferia*), error.
- lades** (Walker), 1849: 403 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 632 (*Erax*); Schiner, 1866: 394 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, lost.
- lascivus** (Wiedemann), 1828: 474 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1849: 400 (*Asilus*), 1855: 637 (*Erax*); Schiner, 1866: 687 (*Erax*), 1867: 394 (*Erax*); Osten Sacken, 1878: 80, 124 (cat.; *Erax*); Williston, 1891: 86 (cat.; *Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Syntypes, NMW, BZM [no. 2038; cf. Schumann, 1973: 124].
amarynceus Walker, 1849: 400 (*Asilus*). Type-locality: unknown. Ref. – Martin, 1965a: 28 (*Efferia*); Martin & Papavero, 1970: 62 (cat.; *Efferia*). Type, BMNH.
- loewii** (Bellardi), 1862: 218 (21), pl. 3, fig. 17 (*Erax*). Type-locality: Mexico, Veracruz: Tuxpano near Orizaba. Distr. – Mexico (Chiapas, Morelos, Veracruz), Guatemala. Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Hine, 1919: 125, fig. 20 (*Erax*; ?specimens from Colombia); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, TORO.
- dolichogaster** Williston, 1901: 326, pl. 6, fig. 6 (*Erax*). Type-locality: Mexico: Veracruz, Atoyac; Tabasco, Teapa. Refs. – Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 236 (cat.; *Erax*); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Syntypes, BMNH.
- lowei* Papavero & Ibáñez Bernal, 2001: 110 (hist. Mexican dipterol), error.
- medianus** (Wiedemann), 1828: 473 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 636 (*Erax*); Schiner, 1866: 687 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 239 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, BZM [no. 2037; cf. Schumann, 1973: 131].
- mexicanus** (Hine), 1919: 123, fig. 27 (*Erax*). Type-locality: Mexico, Chiapas, Tapachula. Distr. – Mexico (Chiapas). Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, OSU.
- nigrinus** (Wiedemann), 1821: 197 (*Asilus*). Type-locality “Brazil” [Sello; cf. Papavero, 1971e: 56-60, 59-80, 99-102]. Refs. – Wiedemann, 1828: 462 (*Asilus*); Walker, 1855: 635 (*Erax*); Schiner, 1866: 686 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Syntypes, NMW [no. 2030; cf. Schumann, 1973: 137].
- nigripes** (Macquart), 1850: 389 (1850: 85), pl. 8, fig. 6 (*Erax*). Type-locality: “Bolivia”. Refs. – Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, MNHNP.
- nigritarsis** (Hine), 1919: 129, fig. 29 (*Erax*). Type-locality: Cuba, Holguín. Refs. – Bromley, 1929a: 290 (*Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72; Alayo Dalmau & Garcia Ávila, 1983: 48 (cat.; *Erax*). Type, OSU.
- obscurus** (Macquart), 1838: 112 (1839: 228) (*Erax*). Type-locality: “Brazil”. Distr. – ?Peru, Brazil (Acre), Bolivia. Refs. – Walker, 1849: 401 (*Asilus*), 1855: 633 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Carrera, 1945a: 162 (*Erax*) (preys); Bromley, 1946: 115 (cat.; *Erax*); Carrera & d’Andretta, 1952: 259 (*Erax*; preys); Carrera, 1959: 6 (*Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); ?Lamas M., 1972: 313 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, lost.
- parvus** (Walker), 1855: 639 (*Asilus*). Type-locality: Brazil, Pará, Santarém. Refs. – Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Syntypes, BMNH.
- pictipennis** (Schiner), 1868: 182 (*Erax*). Type-locality: “South America” (Venezuela). Refs. – Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, NMW.
- pilosulus** (Bromley), 1929a: 292, fig. 11 (*Erax*). Type-locality: Cuba, Sierra Maestra. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Alayo Dalmau & Garcia Ávila, 1983: 48 (cat.; *Erax*); Artigas & Papavero, 1997b: 72. Type, USNM.
- poecilolamprus** (James), 1953: 48, fig. 2 (*Erax*). Type-locality: Honduras, Tela. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, MICH.
- pyrrhogonus** (Wiedemann), 1828: 472 (*Asilus*). Type-locality: “Brazil” [Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-102]. Refs. – Walker, 1855: 636 (*Erax*); Schiner, 1866: 687 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Syntypes, NMW, BZM [no. 2036; cf. Schumann, 1973: 149].
- rufipes** (Macquart), 1838: 112 (1839: 228) (*Erax*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Refs. – Walker, 1855:

- 634 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, MHNTP.
- rufithorax** (Macquart), 1846: 210 (1846: 82) (*Erax*). Type-locality: "Brazil". Refs. – Walker, 1855: 634 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, OXF (abdomen missing).
- rufitibia** (Macquart), 1848: 187 (1848: 27), pl. 2, fig. 11 (*Erax*), part. Type-locality: "Rio Negro" [his Haiti specimen is a synonym of *fortis* Walker, 1855; cf. Curran, 1933: 4]. Refs. – Walker, 1855: 623 (*Erax*); Roeder, 1885: 339 (*Erax*); Williston, 1891: 87 (cat.; *Erax*), 1896a: 305, pl. 10, fig. 79 (*Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 242 (cat.; *Erax*); Bromley, 1929a: 292, pl. 1, fig. 18 (*Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 65 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, MHNTP.
- stigmatus** (Carrera & d'Andretta), 1950: 172 (*Erax*). Type-locality: Mexico, Veracruz, Orizaba. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 66 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, CMNH.
- subappendiculatus** (Macquart), 1838: 114 (1839: 28) (*Erax*). Type-locality: "Brazil". Refs. – Walker, 1855: 633 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 243 (cat.; *Erax*); Martin & Papavero, 1970: 66 (cat.; *Efferia*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Artigas & Papavero, 1997b: 72; Tomasovic, 2002b: 57, fig. 14 (*Efferia*). Type, MHNTP.
- subchalybeus** (Bromley), 1928: 2 (*Erax*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 66 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, AMNH.
- tibialis** (Macquart), 1838: 234 (*Erax*). Type-locality: U. S. A., Pennsylvania, Philadelphia; French Guiana, Cayenne; and San Domingo. Ref. – Martin, 1965a: 29 (*Efferia*); Macquart's specimen from Philadelphia misident.). **N. comb.**
- titan** (Bromley), 1934b: 359 (*Erax*). Type-locality: Guyana, Bartica. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 66 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, AMNH.
- tortola** (Curran), 1928b: 23 (*Erax*). Type-locality: Virgin Islands, Tortola. Refs. – Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 66 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, AMNH.
- vauriei** (Curran), 1953: 4 (*Erax*). Type-locality: Bahama Islands, South Bimini Island. Refs. – Martin, 1957b: 7; Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 67 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, AMNH.
- velox** (Wiedemann), 1828: 468 (*Asilus*). Type-locality: Brazil, Pernambuco. Refs. – Walker, 1855: 638 (*Erax*); Schiner, 1866: 687 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 243 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (n. comb.); Martin & Papavero, 1970: 67 (cat.; *Efferia*); Artigas & Papavero, 1997b: 72. Type, ?

Genus **Pogonioefferia** Artigas & Papavero, 1997

- Pogonioefferia** Artigas & Papavero, 1997b: 73. Type-species, *Asilus pogonias* Wiedemann, 1821 (orig. des.).
Efferia, *pogonias*-group of Wilcox, 1966c: 163 (*rufibarbis*-group, *caudata*-group) and *staminea*-group of Wilcox, 1966c: 177.
- cana** (Hine), 1916: 22 (*Erax*). Type-locality: U. S. A., California, Claremont. Distr. – U. S. A. (California, Arizona, Nevada, Utah, Colorado), Mexico. Refs. – Hine, 1919: 145, fig. 38 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 196 (*Efferia*). Type, OSU.
canus Hine, 1919: 145 (*Erax*). Type-locality: U. S. A., California, Los Angeles County. Type, USNM.
- cannella** (Bromley), 1934a: 95 (*Erax*). Type-locality: U. S. A., Arizona, Sierritas, Black Dike Prospect. Distr. – U. S. A. (Arizona), Mexico. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 197 (*Efferia*). Type, AMNH.
- coquilletti** (Hine), 1919: 149, fig. 14 (*Erax*). Type-locality: U. S. A., California, San Diego County. Distr. – U. S. A. (California, Arizona, Nevada), Mexico. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 198 (*Efferia*). Type, USNM.
- deserti** (Wilcox), 1966c: 199, figs. 186–187 (*Efferia*). Type-locality: U. S. A., California, 10 miles east of Desert Center. Distr. – U. S. A. (California, Arizona, Nevada), Mexico. Type, JW.
- ehrenbergi** (Wilcox), 1966c: 203, figs. 179, 181 (*Efferia*). Type-locality: U. S. A., Arizona, 35 miles east of Ehrenberg. Distr. – U. S. A. (California, Arizona), Mexico. Type, JW.
- inflata** (Hine), 1911b: 310 (*Erax*). Type-locality: U. S. A., California, Los Angeles County. Distr. – U. S. A. (California), Mexico. Refs. – Hine, 1919: 146 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 204 (*Efferia*). Type, OSU.
- parkeri** (Wilcox), 1966c: 213 (*Efferia*). Type-locality: U. S. A., Arizona, 8 miles north of Tucson. Distr. – U. S. A. (Arizona, Texas), Mexico. Type, CAS.
- pogonias** (Wiedemann), 1821: 198 (*Asilus*; nom. nov. for *barbatus* Fabricius, 1805, when preocc. in *Asilus*). Type-locality: "North America". Distr. – Southern U. S. A. east of Rocky Mts., Mexico, West Indies. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin, 1962b: 248 (descr. of type); Martin & Wilcox, 1965: 395 (cat.; *Efferia*); Wilcox, 1966c: 175, figs. 118, 125 (*Efferia*). Type, COPN.

- barbatus** Fabricius, 1805: 169 (*Dasypogon*; preocc. Fabricius, 1787). Type-locality: "North America". Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin, 1962b: 248 (descr. of type). Type, KIEL [cf. Zimsen, 1964: 470].
- rufibarbis** Macquart, 1838: 116 (1839: 232) (*Erax*). Type-locality: "North America". Refs. – Hine, 1919: 137 (*Erax*); Fattig, 1945: 26; Bromley, 1950: 237 (*Erax*).
- dascyllus** Walker, 1849: 401 (*Asilus*). Type-locality: U. S. A., Massachusetts.
- virginianus** Wulp, 1882: 109 (*Proctacanthus*). Type-locality: U. S. A., Virginia. Type, ZMA.
- rava** Coquillett, 1893d: 176 (*Efferia*). Type-locality: U. S. A., Texas. Type, ?USNNM.
- aeastuans** authors, not Linnaeus.
- rapax** (Osten Sacken), 1887: 201 (*Erax*). Type-locality: Mexico, northern Sonora. Distr. – U. S. A. (Arizona, New Mexico, Utah, Texas, Colorado, Kansas), Mexico (Sonora). Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Hine, 1911b: 307 (*Erax*), 1919: 142; Hull, 1962 (2): 478 (*Nerax*); Martin & Wilcox, 1965: 394 (cat.; *Efferia*); Wilcox, 1966c: 216 (*Efferia*); Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipteron). Type, BMNH.
- splendens** (Williston), 1901: 325, pl. 6, fig. 5 (*Erax*). Type-locality: Mexico, Jalisco, San Blás. Distr. – Mexico (Jalisco). Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 242 (cat.; *Erax*); Hine, 1919: 243 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipteron). Type, BMNH.
- triton** (Osten Sacken), 1887: 200 (*Erax*). Type-locality: Mexico, Sinaloa, Presidio, near Mazatlán. Distr. – U. S. A. (Arizona, New Mexico, Kansas, Texas), Mexico (Nayarit, Sinaloa), Honduras. Refs. – Aldrich, 1905: 278 (cat.; *Erax*); Kertész, 1909: 243 (cat.; *Erax*); Hine, 1919: 151 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Lavigne & Dennis, 1985: 146 (ethology); Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipteron). Type, BMNH.
- yuma** (Wilcox), 1966c: 224 (*Efferia*). Type-locality: U. S. A., Arizona, 16 miles east of Yuma. Distr. – U. S. A. (Arizona, California), Mexico. Type, JW.

Genus *Porasilus* Curran, 1934

Porasilus Curran, 1934a: 15. Type-species, *Porasilus barbiellinii* Curran, 1934 (orig. des.).

Ref. – Lamas M., 1971b (rev.).

- barbiellinii** Curran, 1934a: 15. Type-locality: Brazil, São Paulo. Distr. – Brazil (Minas Gerais, São Paulo). Refs. – Carrera, 1945a: 164 (preys); Bromley, 1946: 114 (cat.); Carrera, 1947d: 271 (prey); Hull, 1962 (2): 472, figs. 314, 709, 1386, 1397, 2477, 2487; Lamas M., 1971b: 46, figs. 1-3; Artigas & Papavero, 1997b: 73; Bueno, 1986: 17 (coupling, oviposition, preys), 1987: 903 (biol.; predator of *Deois flavopicta* (Stål, 1854) (Hemiptera, Homoptera, Cercopidae)); Bueno & Berti Filho, 1987: 141 (biol.; predator of *Deois flavopicta* (Stål, 1854) (Hemiptera, Homoptera, Cercopidae))). Type, AMNH.
- cerdai** Tomasovic, 2002b: 61, fig. 20. Type-locality: French Guiana, Petit Saut, Pointe Combi. Type, FUSA Gembloux.
- garciae** Lamas M., 1971b: 51, figs. 10-12. Type-locality: Peru, Lima, Lima. Distr. – Peru (Lima, Junín). Refs. – Lamas M., 1972: 313 (cat.); Artigas & Papavero, 1997b: 73. Type, MZUSP.
- intermedius** Lamas M., 1971b: 53, figs. 13-16. Type-locality: Brazil, Amazonas, Manaus. Distr. – Brazil (Amazonas, Pará). Ref. – Artigas & Papavero, 1997b: 73. Type, MZUSP.
- lesbius** Lamas M., 1971b: 49, figs. 7-9. Type-locality: Brazil, São Paulo, São Bernardo do Campo. Distr. – Brazil (Minas Gerais, São Paulo). Ref. – Artigas & Papavero, 1997b: 73. Type, MZUSP.
- satyrus** Lamas M., 1971b: 46, figs. 4-6. Type-locality: Brazil, Acre, Forte Príncipe da Beira. Distr. – Brazil (Acre, Mato Grosso do Sul, Goiás). Ref. – Artigas & Papavero, 1997: 73. Type, MZUSP.
- senilis** (Wiedemann), 1828: 471 (*Asilus*). Type-locality: Uruguay, Montevideo. Refs. – Walker, 1849: 400 (*Asilus*), 1855: 639 (*Erax*); Schiner, 1866: 687 (*Erax*); Lynch Arribálzaga, 1880: 113 (*Erax*); Wulp, 1882: 115 (*Erax*); Lynch Arribálzaga, 1883a: 149 (cat.; *Erax*), 1883b: 14 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 242 (cat.; *Erax*); Llano, 1959: 90 (*Erax*; biol.); Carrera & Vulcano, 1961: 75 (*Erax*; prey); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 73 (n. comb.). Syntypes, NMW.

Genus *Triorla* Parks, 1968

Trriorla Parks, 1968: 175. Type-species, *Asilus interruptus* Macquart, 1834 (orig. des.).

Ref. – Pamplona & Aires, 1999 (rev.).

- ambigua** (Macquart), 1846: 212 (1846: 84) (*Erax*). Type-locality: U. S. A., Texas, Galveston; Mexico, Yucatán, Mérida. Lectotype MNHNP. Refs. – Walker, 1855: 620 (*Erax*); Osten Sacken, 1878: 234 (cat.; *Erax*); Hull, 1962 (2): 478

- (*Nerax*); Tomasovic, 2002b: 53, fig. 2 (Lectotype des., but does indicate which of the two specimens at LILLE!).
- argyrogaster** (Macquart), 1846: 212 (1846: 84) (*Erax*). Type-locality: Mexico, Yucatán, Mérida. Distr. – Mexico (Oaxaca, Yucatán), Honduras. Refs. – Walker, 1855: 620 (*Erax*); Osten Sacken, 1878: 224 (cat.; *Erax*); Aldrich, 1905: 276 (cat.; *Erax*); Hine, 1919: 152 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Lavigne, 1979: 544 (*Efferia*; ethology); Papavero & Ibáñez Bernal, 2001: 85 (hist. Mexican dipterol.; *Efferia*); Tomasovic, 2002: 55 (n. comb.). Lectotype LILLE [des. by Tomasovic, 2002b: 54].
- argirogaster* Tomasovic, 2002b: 54, fig. 3, error.
- demifasciata** (Macquart), 1850: 390 (1850: 86) (*Erax*). Type-locality: “Guyana”. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas & Papavero, 1997b: 71 (*Nerax*); Tomasovic, 2002: 55, fig. 6 (n. comb.). Lectotype LILLE.
- interrupta** (Macquart), 1834: 310 (*Asilus*). Type-locality: U. S. A., Georgia. Distr. – U. S. A., Mexico, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, Colombia. Type, lost. Refs. – Malloch, 1917 (pupal case of a male); Hine, 1918: 5 (*Erax*); Hine, 1919: 153 (*Erax*); Osterberger, 1930 (pupa); Fattig, 1945: 25 (*Erax*); Bromley, 1950: 237 (*Erax*); Carrera & d’Andretta, 1950: 171 (*Erax*); Martin, 1961b: 1 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin, 1965a: 28 (*Erax*), 1965b: 126, 128 (*Erax*); Martin & Wilcox, 1965: 394 (cat.; *Erax*); Wilcox, 1966c: 205, 206 (*Erax*); Parks, 1968: 176, figs. 4, 7, 12, 13 (n. comb.); Dennis & Knutson, 1988: 854, figs. 2A-C (pupa in ventral, lateral and dorsal views); Artigas & Papavero, 1997b: 73.
- maculatus* Macquart, 1838: 111 (1839: 227), pl. 9, figs. 6a-b (*Erax*), part. Type-locality: Guiana, Colombia and Guadeloupe. Lectotype-locality: “Colombia”. Refs. – Rondani, 1848: 95 (*Erax*); Walker, 1855: 635 (*Erax*); Schiner, 1867: 393 (*Erax*); Osten Sacken, 1878: 234 (cat.; *Erax*), 1887: 200 (*Erax*); Williston, 1891: 86 (cat.; *Erax*), 1901: 323 (*Erax*); Brauer & Bischof, 1903: 244; Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 239 (cat.; *Erax*); Davis, 1919 (*Erax*; pupa); Brown, 1929: 310, pl. 30, figs. 38-39 (*Erax*); Martin, 1965b: 128 (*Efferia*); Tomasovic, 2002b: 55, fig. 9 (Lectotype des.). Lectotype, LILLE.
- lateralis* Macquart, 1838: 116 (1839: 232) (*Erax*). Type-locality: U. S. A., Pennsylvania, Philadelphia. Refs. – Walker, 1855: 617 (*Erax*); Williston, 1885: 70 (*Erax*); Titus & Pratt, 1905: 15, figs. 6a-c (*Erax*).
- concolor* Walker, 1851: 130 (*Erax*). Type-locality: unknown. Refs. – Walker, 1855: 646 (*Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hull, 1962 (2): 479 (*Nerax*). Type, BMNH.
- perrumpens* Walker, 1860: 283 (*Asilus*). Type-locality: “Mexico”. Refs. – Aldrich, 1905: 283 (cat.; *Asilus*); Kertész, 1909: 262 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*). Type, BMNH.
- villosus* Bellardi, 1861: 149 (49) (*Erax*). Type-locality: “Mexico”. Refs. – Aldrich, 1905: 278 (cat.; *Neoeristicus*); Hull, 1962 (2): 475 (*Eicherax*); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.
- perrumpeus* Parks, 1968: 176; Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.), error.
- nervosa** (Macquart), 1838: 118 (*Erax*). Type-locality: unknown. Refs. – Walker, 1855: 646 (*Erax*); Kertész, 1909: 240 (cat.; *Erax*); Tomasovic, 2002: 56, fig. 10. Type, MHNTP.
- parastriola** Pamplona & Aires, 1999: 1128, figs. 1-2, 7-8, 11, 14. Type-locality: Brazil, Roraima, Surumu. Distr. – Brazil (Roraima). Type, MZUSP.
- rubidiventris** (Macquart), 1850: 388 (1850: 84) (*Erax*). Type-locality: “Brazil. M. Ghisbrecht” (in error; Mexico; cf. Papavero, 1971e: 176-178). Refs. – Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 90; Tomasovic, 2002b: 56, fig. 12. Lectotype MHNTP.
- spinosa** Tomasovic, 2002b: 58, fig. 17. Type-locality: “Colombia”. Type, LILLE.
- striola** (Fabricius), 1805: 172 (*Dasyopogon*). Type-locality: “South America”. Distr. – Panama, Colombia, Venezuela, Guyana and Surinam south to Paraguay. Refs. – Wiedemann, 1821: 199 (*Asilus*), 1828: 464 (*Asilus*); Walker, 1849: 432 (*Asilus*), 1855: 635 (*Erax*); Burmeister, 1861: 171 (*Asilus (Eristicus)*); Schiner, 1866: 686 (*Erax*), 1867: 392 (*Erax*); Lynch Arribálzaga, 1880: 117 (*Erax*), 1883a: 113 (cat.; *Erax*), 1883b: 16 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 243 (cat.; *Erax*); Engel, 1930: 461 (*Erax*); Carrera, 1945a: 164 (*Erax*) (preys); Bromley, 1946: 115 (cat.; *Erax*); Carrera, 1947a: 47 (*Erax*); Carrera & d’Andretta, 1952: 258 (*Erax*) (prey); Carrera, 1959: 6 (*Erax*), 1960a: 165 (*Erax*); Carrera & Vulcano, 1961: 75 (*Erax*; preys); Martin, 1961b: 1 (*Efferia*); Hull, 1962 (2): 479 (*Nerax*); Carrera & Machado-Allison, 1963: 254 (*Eicherax*); Parks, 1968: 177, figs. 2, 8, 9 (n. comb.); Dennis & Knutson, 1988: 854, figs. 2A-C (pupa in ventral, lateral and dorsal views); Artigas & Papavero, 1997b: 90; Geller-Grimm, 1999: 215 (material in SENCK). Syntypes, COPN [cf. Zimsen, 1964: 471].
- maculatus* Macquart, 1838: 116 (1839: 232), pl. 9, figs. 6a-b (*Erax*), part. Type-locality: Guiana, Colombia and Guadeloupe. Syntypes, (males only) MHNTP.
- sicyon* Walker, 1849: 399 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 640 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 242 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 90 (syn.). Type, BMNH.
- stimicon* Walker, 1851: 129 (*Erax*). Type-locality: “Colombia”. Refs. – Walker, 1855: 640 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 242 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 90 (syn.). Type, BMNH.
- dilectus* Walker, 1855: 632 (*Erax*). Type-locality: Brazil, Pará, Santarém. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 236 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 90 (syn.). Syntypes, BMNH.
- zetterstedti* Jaennicke, 1867: 362 (*Erax*). Type-locality: “Venezuela”. Refs. – Williston, 1891: 87 (cat.; *Erax*); Kertész,

- 1909: 244 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 90 (syn.). Type, SENCK [cf. Geller-Grimm, 1999: 215; eaten by dermestids, antennae and abdomen missing, eyes eaten away].
curranii Bromley, 1951: 27, fig. 4 (*Erax*). Type-locality: Panama, Canal Zone, Corozal. Refs. – Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 90 (syn.). Type, AMNH.
?*trichina* Tomasovic, 2002b: 59, fig. 18. Type-locality: French Guiana, Kourou. Type, FUSAGemblo.

Eicherax-group

Genus **Eicherax** Bigot, 1857

Eristicus Loew, 1848: 396 (preocc. Wesmael, 1844). Type-species, *Erax nigripes* Bellardi, 1861 (sub. des., Hull, 1962: 476) = *Eicherax bellardii* (Schiner, 1868).

Eicherax Bigot, 1857b: 548 (in key). Type-species, *Erax simplex* Macquart, 1848 (orig. des.).

Neoeristicus Osten Sacken, 1878: 81 (cat.; nom. nov. for *Eristicus* Loew, 1848). Type-species, *Erax nigripes* Bellardi, 1861 (aut.) = *Eicherax bellardii* (Schiner, 1868).

bellardii (Schiner), 1868: 182 (*Eristicus*; nom. nov. for *nigripes* Bellardi, 1861). Type-locality: “Mexico”. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 232 (cat.; *Neoeristicus*); Martin & Papavero, 1970: 70 (cat.; as syn. of *nigripes* Bellardi, 1861); Artigas & Papavero, 1995e: 37.
nigripes Bellardi, 1861: 148 (*Erax*; preocc. Macquart, 1850). Type-locality: “Mexico”. Refs. – Aldrich, 1905: 278 (cat.; *Neocriaticus*); Kertész, 1909: 232 (cat.; *Neoeristicus*); Hine, 1917: 298; Bromley, 1946: 115 (cat.); James, 1953: 47; Hull, 1962 (2): 475, figs. 322, 742, 1419, 1428, 2252; Martin, 1965a: 30; Martin & Papavero, 1970: 70 (cat.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, ?OXF (Coll. Bigot).

disjunctus Williston, 1901: 326 (*Erax*). Type-locality: Mexico, Veracruz, Atoyac. Refs. – Aldrich, 1905: 276 (cat.; *Erax*); Kertész, 1909: 236 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.

costatus (Schiner), 1868: 181 (*Erax*). Type-locality: “Venezuela”. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 70 (cat.); Artigas & Papavero, 1995e: 37. Syntypes, NMW.

culiciformis (Walker), 1855: 632 (*Asilus*). Type-locality: Brazil, Pará, rio Tapajós. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 70 (cat.); Artigas & Papavero, 1995e: 37. Type, BMNH.

fulvithorax (Macquart), 1838: 113 (1839: 229) (*Erax*). Type-locality: Brazil, Pará, Belém, and French Guiana. Distr. – Colombia, French Guiana, Peru (Cuzco, Huánuco, Junín), Brazil (Pará). Refs. – Rondani, 1848: 96 (*Erax*); Walker, 1855: 631 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Carrera & Machado-Allison, 1963: 255 (n. comb.); Martin & Papavero, 1970: 70 (cat.); Lamas M., 1972: 313; Artigas & Papavero, 1995e: 37. Syntypes, MNHNP.

parphorus Walker, 1851: 127 (*Erax*). Type-locality: “Colombia”. Refs. – Walker, 1855: 632 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Carrera & d'Andretta, 1953: 70 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 73. Type, BMNH (apex of abdomen eaten away; first submarginal cell very narrowly open).

macularis (Wiedemann), 1821: 193 (*Asilus*). Type-locality: “Brazil (Bescke)” [probably Rio de Janeiro, Nova Friburgo; cf. Papavero, 1971e: 88]. Distr. – Peru, Brazil (Bahia, Mato Grosso and Rio de Janeiro), Paraguay. Refs. – Wiedemann, 1828: 447 (*Asilus*); Macquart, 1838: 147, pl. 12, fig. 5 (*Asilus*); Walker, 1855: 705 (*Asilus*); Schiner, 1866: 684 (*Erax*), 1868: 180 (*Erax*); Wulp, 1882b: 112 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 239 (cat.; *Erax*); Carrera, 1945a: 164 (*Erax*) (preys); Bromley, 1946: 115 (cat.); Carrera, 1947a: 46, 1947d: 271 (prey); Carrera & d'Andretta, 1952: 258 (preys); Carrera, 1958a: 152, 1960a: 165; Carrera & Vulcano, 1961: 75 (preys); Hull, 1962 (2): 475, fig. 275, 697, 2384, 2401; Fisher, 1985: 34; Artigas & Papavero, 1995e: 37; Geller-Grimm, 1999: 211 (material in SENCK). Syntypes, NMW (some other species mixed up with the syntypic series).

eupator Walker, 1851: 147 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 703 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 256 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 475; Martin & Papavero, 1970: 70 (cat.); Artigas & Papavero, 1995e: 37 (syn.). Type, BMNH.

minor (Macquart), 1847: 57 (1847: 41), pl. 1, fig. 8 (*Erax*). Type-locality: “Brazil”. Refs. – Walker, 1855: 635 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995e: 37 (n. comb.). Type, MNHNP.

ricnotes (Engel), 1930: 461, fig. 4 (*Neoeristicus*). Type-locality: Argentina, Formosa, Estancia Tapikiolé; and Bolivia, La Paz, Mapiri, Sarampiuní, 700 m. Distr. – ?Peru (Huánuco), Bolivia, Argentina (Formosa). Refs. – Bromley, 1946: 115 (cat.); Carrera & d'Andretta, 1953: 74; Hull, 1962 (2): 475; Martin & Papavero, 1970: 70 (cat.); Rinaldi, Pailhé & Popolizio, 1971: 451; ?Lamas M., 1972: 313 (cat.); Artigas & Papavero, 1995e: 37, figs. 11–20. Syntypes, ZSBS.

simplex (Macquart), 1848: 187 (1848: 27), pl. 2, fig. 14 (*Erax*). Type-locality: “Rio Negro”. Distr. – Venezuela (Zulia). Refs. – Walker, 1855: 638 (*Erax*); Bigot, 1857b: 545; Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 244 (cat.); Bromley, 1946: 115 (cat.); Hull, 1962 (2): 475, figs. 73, 1627, 1628; Carrera & Machado-Allison, 1963: 154; Martin & Papavero, 1970: 70 (cat.); Artigas & Papavero, 1995e: 37. Type, ?

Genus **Eraxasilus** Carrera, 1959

Eraxasilus Carrera, 1959: 6. Type-species, *Eraxasilus pruinosus* Carrera, 1959 (orig. des.).

acuminatus Carrera, 1959: 9, fig. 5. Type-locality: Paraguay. Nueva Germania, río Aguaray-Guazú. Distr. – Paraguay. Ref. – Artigas & Papavero, 1995e: 37. Type, ZSBS.

amphissa (Walker), 1849: 406 (*Erax*). Type-locality: “Brazil”. Refs. – Walker, 1855: 633 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 233 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995e: 37 (n. comb.). Type, BMNH.

fuscipennis (Macquart), 1847: 56 (1856: 40) (*Erax*). Type-locality: “Brazil”. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995e: 37. Type, OXF (tip of abdomen missing).

gerion (Walker), 1849: 433 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 701 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 256 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995e: 37 (n. comb.). Type, BMNH.

hebes (Walker), 1855: 704 (*Asilus*). Type-locality: Brazil, Pará, Santarém. Refs. – Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 257 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995e: 37 (n. comb.). Syntypes, BMNH.

longiusculus (Walker), 1855: 706 (*Asilus*). Type-locality: Brazil, Pará, Santarém. Refs. – Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 259 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995e: 37. Type, BMNH (tip of abdomen missing).

luctuosus (Macquart), 1838: 146 (1848: 262) (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 701 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 259 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Carrera & d’Andretta, 1952: 258 (*Asilus*; preys), 1961: 77 (*Asilus*; preys); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995: 37e, figs. 1-10. Syntypes, MNHNP.

?**tibialis** Rondani, 1850: 187 (*Asilus*; preocc. Pallas in Wiedemann, 1818). Type-locality: Brazil, São Paulo, ilha de São Sebastião. Ref. – Hull, 1962 (2): 546 (*Asilus*). Type, TORO.

peticus (Walker), 1849: 434 (*Asilus*). Type-locality “Brazil”. Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 262 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995e: 37 (n. comb.). Type, BMNH.

potamon (Walker), 1851: 128 (*Erax*). Type-locality: “South America”. Refs. – Walker, 1855: 633 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Carrera, 1959: 10, fig. 6 (n. comb.); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995e: 37. Type, BMNH.

pruinosus Carrera, 1959: 7, figs. 2-4. Type-locality: Brazil, São Paulo, Onda Verde (Fazenda São João). Distr. – Brazil (Goiás, São Paulo). Refs. – Carrera & Vulcano, 1961: 77 (preys); Hull, 1962 (2): 508; Artigas & Papavero, 1995e: 37. Type, MZUSP.

Eichoichemus-group

Genus **Eichoichemus** Bigot, 1857

Eichoichemus Bigot, 1857b: 543 (in key). Type-species, *Erax flavianalis* Macquart, 1848 (orig. des.).

Proctophorus Schiner, 1866: 66 (key), 674 (diagnosis), 712 (in list of spp.). Type-locality: *Asilus pygophorus* Wiedemann, 1828 (orig. des.).

connexus (Wiedemann), 1828: 497 (*Asilus*). Type-locality: Uruguay. Montevideo. Refs. – Walker, 1855: 592 (*Trupanea*); Schiner, 1866: 689 (*Proctophorus*); Lynch Arribálzaga, 1880: 120 (*Proctophorus*), 1883a: 150 (cat.; *Proctophorus*); Williston, 1891: 84 (cat.; *Proctophorus*); Kertész, 1909: 230 (cat.; *Proctophorus*); Bromley, 1946: 114 (cat.); Hull, 1962 (2): 460; Martin & Papavero, 1970: 60 (cat.); Artigas & Papavero, 1995a: 98. Syntypes, NMW.

eraxoides (Curran), 1935: 10 (in key), 11 (*Promachina*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Bromley, 1946: 114 (cat.); Hull, 1962 (2): 467 (*Promachina*); Martin & Papavero, 1970: 64 (cat.; *Efferia*); Artigas &

- Papavero, 1995a: 98 (n. comb.). Type, AMNH.
- flavianalis** (Macquart), 1848: 186 (1848: 26), pl. 2, figs. 12, 12a (not 13 as in text, p. 186; see “Explanation des figures”, p. 230) (*Erax*). Type-locality: Brazil, Minas Gerais. Refs. – Walker, 1855: 636 (*Erax*); Bigot, 1857b: 543; Williston, 1891: 85 (cat.; *Erax*), 1894: 137; Kertész, 1909: 245 (cat.); Bromley, 1946: 114 (cat.); Hull, 1962 (2): 460; Martin & Papavero, 1970: 60 (cat.); Artigas & Papavero, 1995a: 99. Type, OXF.
- gavius** (Walker), 1851: 151 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 703 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 256 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 89 (cat.; *Asilus*); Artigas & Papavero, 1995a: 99. Type, BMNH (?syn. of *pyrrhomystax* Wiedemann, 1828; the TP is a female and therefore this species cannot be correctly identified).
- kalettae** Ayala, 1978: 34, pl. 1, figs. 4-7. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Ref. – Artigas & Papavero, 1995a: 99. Holotype, FAUCV.
- lizbethae** Ayala, 1978: 32, pl. 1, figs. 1-3. Type-locality: Venezuela, Distrito Federal, Macuto, río Cojo. Ref. – Artigas & Papavero, 1995a: 99. Type, FAUCV.
- lycorius** (Walker), 1851: 143 (*Asilus*). Type-locality: “South America”. Refs. – Walker, 1855: 703 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 259 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*); Artigas & Papavero, 1995a: 99. Type, BMNH. (As occur with *gavius* Walker, 1851, this species may be a syn. of *pyrrhomystax* Wiedemann, 1828, but as the TP is a female, we cannot be sure of this synonymy).
- melaleucus** (Wiedemann), 1828: 498 (*Asilus*). Type-locality: “Brazil”. Refs. – Schiner, 1866: 690 (*Proctophorus*); Williston, 1891: 84 (cat.; *Proctophorus*); Bromley, 1946: 114 (cat.; [*Eichoichemus* (*Proctophorus*)]); Carrera, 1947d: 270 (prey); Hull, 1962 (2): 460; Martin & Papavero, 1970: 60 (cat.); Artigas & Papavero, 1995a: 99. Syntypes, NMW, BZM [no. 731; cf. Schumann, 1973: 131].
- neowillistoni** (Bromley), 1933: 15 (*Erax*; nom. nov. for *willistoni* Bromley, 1928). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Curran, 1935: 10 (in key); Bromley, 1946: 114 (cat.); Hull, 1962 (2): 475 (*Efferia*); Martin & Papavero, 1970: 66 (cat.; *Efferia*); Artigas & Papavero, 1995a: 99, figs. 1-8.
- willistoni* Bromley, 1928: 2 (*Erax*; preocc. Hine, 1919). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Type, AMNH.
- propinquus** (Bromley), 1928: 5 (*Erax*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Bromley, 1946: 114 (cat.); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 67 (cat.; *Efferia*); Artigas & Papavero, 1995a: 99 (n. comb.). Type, AMNH.
- pumilus** (Macquart), 1850: 389 (1850: 85), pl. 8, fig. 9 (*Erax*). Type-locality: “South America”. Refs. – Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Carrera & d’Andretta, 1953: 74 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 67 (cat.; *Efferia*); Artigas & Papavero, 1995a: 99. Type, MNHNP (tip of abdomen missing; the poor drawing of the male terminalia makes it difficult to identify this species with certainty).
- pyrrhomystax** (Wiedemann), 1828: 497 (*Asilus*). Type-locality: “Brazil” [Freyreiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1855: 590 (?*Trupanea*); Schiner, 1866: 690 (*Proctophorus*), 1867: 391 (*Proctophorus*); Williston, 1891: 84 (cat.; *Proctophorus*); Kertész, 1909: 231 (cat.; *Proctophorus*); Carrera, 1945: 163 (prey); Bromley, 1946: 114 (cat.; [*Eichoichemus* (*Proctophorus*)]); Carrera & d’Andretta, 1952: 258 (prey), 1961: 76 (preys); Hull, 1962 (2): 460, figs. 197, 719, 1440, 1449, 2399, 2402; Martin & Papavero, 1970: 60 (cat.); Artigas & Papavero, 1995a: 99. Syntypes, NMW, BZM [no. 732; cf. Schumann, 1973: 149].

Genus *Proctophoroides* Artigas & Papavero, 1995

Proctophoroides Artigas & Papavero, 1995a: 99. Type-species, *Erax hyalipennis* Macquart, 1838 (orig. des.).

hyalipennis (Macquart), 1838: 114 (1839: 230) (*Erax*). Type-locality: Brazil, western Minas Gerais. Refs. – Walker, 1855: 639 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995a: 99 (n. comb.). Type, MNHNP.

latiforceps (Bromley), 1928: 3 (*Erax*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Bromley, 1946: 114 (cat.; *Eichoichemus*); Hull, 1962 (2): 475 (*Efferia*); Artigas & Papavero, 1995a: 99 (n. comb.). Type, AMNH.

Genus *Wygodasilus* Artigas & Papavero, 1995

Wygodasilus Artigas & Papavero, 1995a: 99. Type-species, *Erax pulchripes* Bromley, 1928 (orig. des.).

pulchripes (Bromley), 1928: 4 (*Erax*). Type-locality: “Piedra, Brazil” [in error; actually Bolivia, Santa Cruz, Piedra Blanca (4 miles west of Corumbá, Mato Grosso do Sul, Brazil); cf. Papavero, 1973d: 380]. Distr. – Bolivia (Santa Cruz). Refs. –

Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995a: 99 (n. comb.). Holotype, AMNH.

***Glaphyropyga*-group**

Genus *Glaphyropyga* Schiner, 1866

Glaphyropyga Schiner, 1866: 668 (in key), 647 (1868: 187, second erection of genus). Type-species, *Asilus himantocerus* Wiedemann, 1828 (orig. des.).

Tapinostylus Enderlein, 1914a: 256. Type-species, *Tapinostylus setosifemur* Enderlein, 1914 (orig. des.).

Opopotes Hull, 1957: 884. Type-species, *Opopotes attenuatus* Hull, 1957 (orig. des.). Ref. – Artigas & Papavero, 1995b: 15 (syn.).

aristata Carrera, 1950a: 29, fig. Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. – Costa Rica, Panama, ?Peru (Huánuco). Refs. – Carrera & d'Andretta, 1953: 70; Hull, 1962 (2): 502; Lamas M., 1972: 313 (cat.); Artigas & Papavero, 1995b: 15, fig. 4. Type, USNM.

attenuata (Hull), 1957: 884 (*Opopotes*). Type-locality: “Costa Rica”. Refs. – Hull, 1962 (2): 503, figs. 363, 747, 1493, 1502, 2357, 2400 (*Opopotes*); Artigas & Papavero, 1995b: 15 (n. comb.). Type, BMNH.

bolivari Ayala, 1983: 8, figs. 2, 10, 17. Type-locality: Venezuela, Aragua, Rancho Grande, via la Toma, 1100 m. Distr. – Venezuela (Aragua, Bolívar, Carabobo, Guárico, Miranda). Type, IZAUCV.

carrerai Ayala, 1983: 6, figs. 1, 8, 23. Type-locality: Venezuela, Miranda, Caracas (Quebrada Pasaquire), 1000 m. Distr. – Venezuela (Aragua, Distrito Federal, Mérida, Miranda, Monagas). Type, IZAUCV.

dryas Fisher & Hespenheide, 1982: 712, figs. 1-6 Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. – Costa Rica (Cartago, Heredia), Panama, Venezuela (Aragua, Guárico, Lara, Miranda, Zulia). Refs. – Ayala, 1983: 12, figs. 4, 12, 20; Artigas & Papavero, 1995b: 15. Type, USNM.

himantocera (Wiedemann), 1828: 447 (*Asilus*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Peru (Huánuco), Brazil (Goiás, Rio de Janeiro, São Paulo, Paraná). Refs. – Walker, 1855: 702 (*Asilus*); Schiner, 1866: 685; Williston, 1891: 89 (cat.); Kertész, 1909: 284 (cat.); Carrera, 1945c: 177, figs. 1-11, 36-38; Bromley, 1946: 117 (cat.); Carrera & d'Andretta, 1953: 69; Carrera, 1960a: 166; Hull, 1962 (2): 502, figs. 324, 731, 1401, 1410, 2186, 2462, 2481; Carrera & Machado-Allison, 1963: 255, fig. 3; Lamas M., 1972: 313 (cat.); Artigas & Papavero, 1995b: 15, fig. 5. Type, SENCK [cf. Geller-Grimm, 1999: 209].

pollinifera Carrera, 1945c: 181, figs. 12-21. Type-locality: Brazil, São Paulo, São Paulo. Distr. – Peru, Brazil (Rio de Janeiro, São Paulo), Bolivia. Refs. – Carrera, 1959: 6, 1960a: 166; Hull, 1962 (2): 502, fig. 2187; Artigas & Papavero, 1995b: 15, figs. 1-2, 7, 8-14; Fisher, 1985: 34. Type, MZUSP.

renatoi Ayala, 1983: 12, figs. 6, 14, 19. Type-locality: Venezuela, Táchira, San Joaquín de Navay, 300 m. Distr. – Venezuela (Aragua). Type, IZAUCV.

setosifemur (Enderlein), 1914a: 256, figs. 3-4 (*Tapinostylus*). Type-locality: Ecuador, Napo, Archidona and Tungurahua: Hacienda Santa Inés, 1244 m (10°25'S, 78°12'W) [cf. Brown, 1941: 846]. Refs. – Bromley, 1946: 117 (cat.; *Tapinostylus*); Hull, 1962 (2): 502; Artigas & Papavero, 1995b: 15. Syntypes, IZPAN.

tachirensis Ayala, 1983: 14, figs. 7, 15, 22. Type-locality: Venezuela, Táchira, Río Frío, 450 m. Distr. – Venezuela (Táchira). Type, IZAUCV.

tiarensis Ayala, 1983: 10, figs. 3, 11, 18. Type-locality: Venezuela, Aragua, Campamento Rangel, vía La Tiara, 1200 m. Distr. – Venezuela (Aragua). Type, IZAUCV.

venezuelensis Carrera & Machado-Allison, 1963: 256. Type-locality: Venezuela, Aragua, Rancho Grande, 1100 m. Distr. – Venezuela (Aragua, Carabobo). Refs. – Artigas & Papavero, 1995b: 15, figs. 3, 6; Ayala, 1983: 12, figs. 5, 13, 21. Type, FAUCV.

Genus *Leptoharpacticus* Lynch Arribálzaga, 1880

Leptoharpacticus Lynch Arribálzaga, 1880: 178. Type-species, *Asilus mucius* Walker, 1849 (orig. des.).

mucius (Walker), 1849: 463 (*Asilus*). Type-locality: Uruguay, Montevideo. Refs. – Walker, 1855: 707 (*Asilus*); Lynch Arribálzaga, 1880: 180, 1883: 152 (cat.); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 265 (cat.; *Asilus*); Hull, 1962 (2): 511, figs. 373, 687, 1494, 1503, 2346, 2359; Artigas & Papavero, 1995b: 15. Type, BMNH (lacking one antenna, both wings, pair of middle legs and hind left leg).

sp. Artigas & Papavero, 1995b: 32, figs. 87-99.

Genus **Megalometopon** Artigas & Papavero, 1995

Megametopon Artigas, 1970a: 312 (preocc. Giglio-Tos, 1891). Type-species, *Asilus occidentalis* Philippi, 1865 (orig. des.).
Megalometopon Artigas & Papavero, 1995b: 15 (*nom. nov.* for *Megametopon* Artigas, 1970). Type-species, *Asilus occidentalis* Philippi, 1865 (aut.).

immisericorde (Artigas), 1970a: 314, figs. 301-304, 422 (*Megametopon*). Type-locality: Chile, Valparaíso, Perales. Distr. – Chile (Santiago, Valparaíso). Refs. – Artigas, 1971: 41, figs. 99-100; Artigas & Papavero, 1995b: 15, figs. 36-48 (n. comb.). Type, DZUC.

occidentale (Philippi), 1865: 696 (*Asilus*). Type-locality: Chile, Valdivia. Neotype-locality: Chile, Talca, Arnerillo. Distr. – Chile (Aconcagua, Arauco, Cautín, Concepción, Curicó, Linares, Malleco, Ñuble, O'Higgins, Santiago, Talca, Valdivia, Valparaíso). Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 261 (cat.; *Asilus*); Bromley, 1932: 281 (*Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas, 1970a: 315, figs. 305-311, 430 (*Megametopon*), 1971: 41, figs. 101-102; Artigas & Papavero, 1995b: 15 (n. comb.). Type, lost. NT DZUC [des. by Artigas, 1970a: 315].

Genus **Neotes** Artigas & Papavero, 1995

Nesiotes Artigas, 1970a: 333 (preocc. Martens, 1860). Type-species, *Nesiotes chiloensis* Artigas, 1970 (orig. des.).
Neotes Artigas & Papavero, 1995b: 15 (*nom. nov.* for *Nesiotes* Artigas, 1970). Type-species, *Nesiotes chiloensis* Artigas, 1970 (aut.).

chiloensis (Artigas), 1970: 334, figs. 345-352, 423 (*Nesiotes*). Type-locality: Chile, Chiloé, Isla de Chiloé, Dalcahue. Distr. – Chile (Cautín, Chiloé, Osorno). Refs. – Artigas, 1971: 45, fig. 113 (*Nesiotes*); Artigas & Papavero, 1995b: 15, figs. 15-28 (n. comb.). Type, DZUC.

Genus **Nomomyia** Artigas, 1970

Nomomyia Artigas, 1970a: 336. Type-species, *Erax murinus* Philippi, 1865 (orig. des.).

ivetteae Artigas, 1970a: 337, figs. 353-357, 475. Type-locality: Chile, Coquimbo, El Pangue, south of Vicuña. Distr. – Chile (Coquimbo, Santiago). Refs. – Artigas, 1971: 46, figs. 114-115; Artigas & Papavero, 1995b: 16, figs. 61-68. Type, DZUC.
murina (Philippi), 1865: 694 (*Erax*). Type-locality: Chile, Santiago. Distr. – Chile. Refs. – Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Bromley, 1932: 266; Stuardo Ortiz, 1946: 84 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas, 1970a: 339, figs. 358-362, 456 (n. comb.), 1971: 47, figs. 116-117; Artigas & Papavero, 1995b: 16, figs. 69-72. Type, MNHNS.

Genus **Regasilus** Curran, 1931

Regasilus Curran, 1931b: 24. Type-species, *Regasilus strigarius* Curran, 1931 (orig. des.).

strigarius Curran, 1931b: 24 (as *strigaria*, error). Type-locality: Ecuador, Guayas, Guayaquil. Distr. – Ecuador, Peru (Cuzco, Piura). Type, AMNH. Refs. – Hull, 1962 (2): 506, figs. 207, 685, 1406, 1415, 2200, 2209, 2421, 2422; Lamas M., 1972: 314 (cat.); Artigas & Papavero, 1995b: 16.

Genus **Tsacasia** Artigas & Papavero, 1995

Tsacasia Artigas & Papavero, 1995b: 16. Type-species, *Tacasia wagneri* Artigas & Papavero, 1995 (orig. des.).

wagneri Artigas & Papavero, 1995b: 17, figs. 29-35. Type-locality: Argentina, Santiago del Estero, 56 km south of Icaño. Distr. – Argentina (Santiago Del Estero). Type, MNHNP.

Genus **Zoticus** Artigas, 1970

Zoticus Artigas, 1970a: 368. Type-species, *Zoticus toconaoensis* Artigas, 1970 (orig. des.)/

fitzroyi Artigas, 1974c: 119, figs. 1-7. Type-locality: Argentina, Santa Cruz, 24 km south of FitzRoy. Distr. – Argentina (Santa Cruz). Ref. – Artigas & Papavero, 1995b: 17, figs. 73-86 TP DZUC.

toconaoensis Artigas, 1970a: 369, figs. 400-405, 489. Type-locality: Chile, Antofagasta, Toconao. Distr. – Chile (Antofagasta). Ref. – Artigas, 1971: 52, figs. 134-135; Artigas & Papavero, 1995b: 17. Type, DZUC.

Lecania-group

Genus **Cerozodus** Bigot, 1857

Cerozodus Bigot, 1857b: 543 (in key). Type-species, *Asilus nodicornis* Wiedemann, 1828 (orig. des.).

nodicornis (Wiedemann), 1828: 469 (*Asilus*). Type-locality: “Brazil”. Distr. – Brazil (São Paulo). Refs. – Macquart, 1838: 108 (*Erax*); Walker, 1855: 638 (?*Erax*); Bigot, 1857b: 543; Schiner, 1866: 667 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 244 (cat.); Bromley, 1946: 115 (cat.); Carrera, 1946d: 250, figs. 1-14; Hull, 1962 (2): 485, figs. 372, 712, 1399, 1408, 2442, 2443; Artigas & Papavero, 1995b: 36, figs. 1-11. Type, BZM [no. 2040; cf. Schumann, 1973: 138].

Genus **Ctenodontina** Enderlein, 1914

Ctenodontina Enderlein, 1914a: 260. Type-species, *Ctenodontina pectinatipes* Enderlein, 1914 (orig. des.).

Catostola Hull, 1958c: 320. Type-species, *Catostola carrerai* Hull, 1958 (orig. des.) = *Ctenodontina maya* Carrera & d’Andretta, 1953. Ref. – Martin & Papavero, 1970: 70 (syn.).

martini Fisher in Fisher & Hespenheide, 1992: 630 (note 7) (*nom. nov.* for *maya* Martin, 1975). Type-locality: Mexico, Chiapas, Tacaná, San Jerónimo. Distr. – Mexico (Chiapas, Yucatán), Guatemala.

maya Martin, 1975c: 87, fig. 78 (*Pachychaeta*; preocc. Carrera & d’Andretta, 1953). Type-locality: Mexico, Chiapas, Tacaná, San Jerónimo. Ref. – Artigas & Papavero, 1995: 38. Type, CAS.

maya Carrera & d’Andretta, 1953: 75, figs. 9-11. Type-locality: Peru, Huánuco, Tingo María. Distr. – Peru (Cuzco, Huánuco, Junín, Madre de Dios). Refs. – Hull, 1962 (2): 482; Lamas M., 1972: 313 (cat.), 1973b: 276, figs. 1-3; Martin & Papavero, 1970: 70 (cat.); Fisher, 1985: 34; Artigas & Papavero, 1995b: 36, fig. 12. Type, MZUSP.

carrerai Hull, 1958c: 323 (*Catostola*). Type-locality: Peru, Junín, Valle de Chanchamayo. Distr. – Peru (Junín). Refs. – Hull, 1962 (2): 482, figs. 349, 394, 1545, 1554, 2201, 2205; Lamas M., 1972: 313 (cat.), 1973b: 276 (syn.); Artigas & Papavero, 1995b: 36, figs. 17-27, 29-34. Type, MZUSP.

mochica Lamas M., 1973b: 277, figs. 4-7. Type-locality: Peru, La Libertad, Cartavio. Distr. – Peru. Ref. – Artigas & Papavero, 1995b: 36. Type, Museo de Entomología, Universidad Nacional Agraria, Lima, Peru.

pectinatipes Enderlein, 1914a: 261, figs. 6-7. Type-locality: Colombia, Caldas, Victoria (Hacienda Pehlke), 900 m [cf. Papavero, 1973d: 299]. Distr. – Colombia. Refs. – Carrera & d’Andretta, 1953: 77; Hull, 1962 (2): 480, figs. 2508, 2524; Martin & Papavero, 1970: 71 (cat.); Lamas M., 1973b: 276; Artigas & Papavero, 1995b: 36. Type, IZPAN.

sp. Artigas & Papavero, 1995b: fig. 28. Argentina, Jujuy, La Legua.

Genus **Lecania** Macquart, 1838

Lecania Macquart, 1838: 131 (1839: 247). Type-species, *Lecania rufipes* Macquart, 1838 (sub. des., Hull, 1962 (2): 480).

Pachychaeta Bigot, 1857b: 545 (as *Pachychoeta* in key). Type-species, *Erax annulipes* Macquart, 1946 (orig. des.).

Nyssoprosopa Hull, 1962 (2): 519. Type-species, *Nyssoprosopa pollinosa* Hull, 1962 (orig. des.).

Aristofolia Ayala, 1978: 38. Type-species, *Aristofolia lapila* Ayala, 1978 (orig. des.). **N. syn.**

annulipes (Macquart), 1846: 211 (1846: 83), pl. 8, fig. 8 (*Erax*). Type-locality: “Brazil”. Refs. – Walker, 1855: 642 (*Erax*); Bigot, 1857b: 545; Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 245 (cat.); Bromley, 1946: 116 (cat.; *Pachychoeta*); Hull, 1962

- (2): 481 [*Lecania (Pachychaeta)*]; Martin, 1975c: 85 (*Pachychoeta*); Artigas & Papavero, 1995b: 37. Type, ?lost.
- apicalis** (Bromley), 1934b: 355 (*Pachychaeta*). Type-locality: Guyana, Bartica. Refs. – Hull, 1962 (2): 481; Martin, 1975c: 85 (*Pachychoeta*); Artigas & Papavero, 1995b: 37. Type, AMNH.
- baleta** (Walker), 1849: 427 (*Asilus*). Type-locality: “Venezuela”. Refs. – Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995b: 37. Type, BMNH.
- ?*caracasae* Martin, 1975c: 85 (*Pachychaeta*). Type-locality: Venezuela, Quebrada Caxia, south of Caracas. Type, CAS.
- boraceae** Carrera, 1958a: 150, figs. 3-5. Type-locality: Brazil, São Paulo, Salesópolis (Estação Biológica de Boracéia). Ref. – Artigas & Papavero, 1995b: 37, figs. 65-69. Type, MZUSP.
- clavata** (Macquart), 1838: 147 (1839: 263), pl. 12, fig. 3 (*Asilus*). Type-locality: Brazil, western Minas Gerais. Refs. – Walker, 1855: 706 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 253 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995b: 37. Type, MNHNP.
- ?*calatina* Walker, 1849: 436 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 708 (*asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 252 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*). Type, BMNH.
- complicata** (James), 1953: 52, fig. 6 (*Pachychaeta*). Type-locality: Honduras, Guaimas, Tela. Refs. – Hull, 1962 (2): 481; Martin, 1975c: 86 (*Pachychoeta*); Artigas & Papavero, 1995b: 37. Type, MICH.
- copulata** (Wiedemann), 1819: 48 (*Asilus*). Type-locality: “Brazil”. Refs. – Wiedemann, 1821: 192 (*Asilus*), 1828: 457 (*Asilus*); Walker, 1855: 637 (*Erax*); Schiner, 1866: 686 (*Erax*), 1867: 394 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hull, 1962 (2): 481; Artigas & Papavero, 1995b: 37; Geller-Grimm, 1999: 208 [specimen in SENCK]. Syntypes, NMW.
- ctesicles** (Walker), 1851: 149 (*Asilus*). Type-locality: “South America”. Refs. – Walker, 1855: 705 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 255 (cat.; *Asilus*); Bromley, 1946: 117 (cat.; *Pachychoeta*); Hull, 1962 (2): 481; Artigas & Papavero, 1995b: 38. Type, BMNH.
- femorata** Macquart, 1838: 132 (1839: 248). Type-locality: unknown. Refs. – Walker, 1855: 762; Williston, 1891: 89 (cat.; *Senoprosopis*); Kertész, 1909: 285 (cat.; *Stenoprosopis*); Hull, 1962 (2): 481; Artigas & Papavero, 1995b: 38. Syntypes, MNHNP, OXF.
- flavida** (Wiedemann), 1828: 473 (*Asilus*). Type-locality: Uruguay, Montevideo. Refs. – Walker, 1855: 639 (*Erax*); Schiner, 1866: 687 (*Asilus*); Lynch Arribálzaga, 1880: 116 (*Erax*), 1883a: 148 (cat.; *Erax*); Osten Sacken, 1891: 420 (*Erax*); Kertész, 1909: 236 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995b: 38. Type, BZM [no. 2041; cf. Schumann, 1973: 111].
- flavidus** Macquart, 1838: 114 (1839: 230) (*Erax*; preocc. Wiedemann, 1828). Type-locality: “Brésil, à l’embouchure de l’Uruguay jusqu’aux Missions” [= Uruguay, western part, along Uruguay river; cf. Papavero, 1973d: 121]. Refs. – Walker, 1855: 637 (*Erax*); Lynch Arribálzaga, 1880: 37 (*Erax*), 1883a: 149 (cat.; *Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 236 (cat.; *Erax*); Bromley, 1946: 114 (cat.; *Erax*); Martin & Papavero, 1970: 64 (cat.; syn.). Type, MNHNP.
- genitalis** (Bromley), 1934: 256, fig. 20 (*Pachychaeta*). Type-locality: Guyana, Tropical Research Station. Refs. – Hull, 1962 (2): 481, figs. 699 (as *Lecania (Pachychaeta)* in caption of figure), 1403, 1412, 1604, 2434; Martin, 1975c: 86 (*Pachychoeta*); Artigas & Papavero, 1995b: 38. Type, AMNH.
- hilarii** (Macquart), 1838: 147 (1839: 263) (*Asilus*). Type-locality: Brazil, Rio de Janeiro. Refs. – Walker, 1855: 707 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 257 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 481; Artigas & Papavero, 1995b: 38. Type, MNHNP.
- inca** (Martin), 1975: 86 (*Pachychaeta*). Type-locality: Venezuela, San Esteban. Ref. – Artigas & Papavero, 1995b: 38. Type, CAS.
- lapila** (Ayala), 1978: 40, pl. 2, figs. 1-8 (*Aristofolia*). Type-locality: Venezuela, Guárico, San Francisco de Tiznados, Quebrada Coronado. Distr. – Venezuela (Guárico). Type, IZAUCV. **N. comb.**
- leucopyga** (Wiedemann), 1828: 586 (*Asilus*). Type-locality: “Brazil”. Refs. – Schiner, 1866: 690 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 481; Artigas & Papavero, 1995b: 38, figs. 81-83. Type, ?
- mellina** (Wiedemann), 1828: 465 (*Asilus*). Type-locality: Uruguay, Montevideo. Refs. – Walker, 1855: 638 (*Erax*); Schiner, 1866: 686 (*Erax*), 1867: 393 (*Erax*); Lynch Arribálzaga, 1880: 119 (*Erax*), 1883a: 149 (cat.; *Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995b: 38. Syntypes, NMW.
- mygdon** (Walker), 1851: 127 (*Erax*). Type-locality: “South America”. Refs. – Walker, 1855: 634 (*Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Artigas & Papavero, 1995b: 38. Type, BMNH.
- migdon** Hull, 1962 (2): 478 (*Nerax*), error.
- pollinosa** (Hull), 1962 (2): 521, figs. 770, 1418, 1427, 1451, 2251, 2279, 2381, 2416 (*Nyssoprosopa*). Type-locality: “Paraguay”. Ref. – Artigas & Papavero, 1995b: 38. Type, ? (not in ZSBS).
- rufina** (Wiedemann), 1819: 47 (*Asilus*). Type-locality: “Brazil”. Refs. – Wiedemann, 1828: 441 (*Asilus*); Walker, 1855: 705 (*Asilus*); Schiner, 1866: 684 (*Erax*), 1867: 393 (*Erax*), Wulp, 1882a: 78 (*Erax*), 1882b: 113 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Bromley, 1946: 115 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995b: 38, figs. 35-38; Geller-Grimm, 1999: 214 (material in SENCK). Syntypes, NMW, BZM [no. 10290; cf. Schumann, 1973: 153].
- rufipes** Macquart, 1838: 131 (1839: 247), pl. 11, figs. 2, 2a-b. Type-locality: Brazil, Pará, surroundings oof Belém. Refs. – Walker, 1855: 761; Williston, 1891: 89 (cat.; *Senoprosopis*); Kertész, 1909: 285 (cat.; *Stenoprosopis*); Bromley, 1946:

117 (cat.; *Pachychoeta*); Carrera & Vulcano, 1961: 76 (preys); Hull, 1962 (2): 478 (*Nerax*), 481 (*Lecania*), figs. 2185, 2406, 2409; Artigas & Papavero, 1995b: 38, figs. 16, 61-64. Type, MNHNP.
spp. Artigas & Papavero, 1995: figs. 13-15, 39-55, 70-80
virilis (Wiedemann), 1828: 468 (*Asilus*). Type-locality: "Brazil". Refs. – Macquart, 1838: 113 (*Erax*); Walker, 1849: 401 (*Asilus*), 1855: 636 (*Erax*); Schiner, 1866: 687 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 244 (cat.; *Erax*); Bromley, 1946: 117 (cat.; *Pachychoeta*); Hull, 1962 (2): 481; Martin, 1975c: 87 (*Pachychoeta*); Artigas & Papavero, 1995b: 38. Type, NMW.
?sexmaculatus Walker, 1855: 707 (*Asilus*). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 263 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*). Type, BMNH.

***Lochmorhynchus*-group**

Genus ***Apotinocerus*** Hull, 1956

Apotinocerus Hull, 1956: 69. Type-species, *Proctacanthus brevistylatus* Wulp, 1882 (orig. des.).

brevistylatus (Wulp), 1882b: 111 (*Proctacanthus*). Type-locality: "Argentina". Ref. – Lynch Arribálzaga, 1883b: 13 (*Prctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 246 (cat.; *Proctacanthus*); Hull, 1956: 70 (n. comb.), 1962 (2): 493, figs. 395, 738, 1367, 1376, 2181, 2457-2460; Artigas & Papavero, 1995c: 133, figs. 1-9. Type, ZMA (Left antenna missing, flagellum of right antenna missing). [There are several specimens of this species in ZSBS; it was actually Hermann who recognized this genus and who gave it the manuscript name *Apotinocerus*, profited by Hull].

Genus ***Lochmorhynchus*** Engel, 1930

Lochmorhynchus Engel, 1930: 459. Type-species, *Proctacanthus senectus* Wulp, 1882 (orig. des.).

Philonerax Bromley, 1932: 270. Type-species, *Asilus mucidus* Walker, 1837 (orig. des.).

Lonchodogonus Hull, 1962 (2): 496. Type-species, *Lonchodogonus cibratus* Hull, 1962 (orig. des.) = *mucidus* (Walker, 1837). *Eremomyia* Artigas, 1970a: 184 (preocc. Stein, 1898). Type-species, *Eremomyia tuimbreensis* Artigas, 1970 (orig. des.). Ref. – Artigas & Papavero, 1995: 133 (syn.).

Eretomyia Artigas, 1971: 36 (nom. nov. for *Eremomyia* Artigas, 1970). Type-species, *Eremomyia tumbrensis* Artigas, 1970 (aut.). Ref. – Artigas & Papavero, 1995: 133 (syn.).

albicans (Carrera & d'Andretta), 1953: 7, fig. 2-8 (*Erax*). Type-locality: Peru, Lima, Pachacámac, near Lima. Distr. – Peru (Ica, Lima, Piura). Refs. – Hull, 1962 (2): 478 (*Nerax*); Lamas M., 1972: 314 (cat.); Artigas & Papavero, 1995c: 133, figs. 25-32. Type, MZUSP.

albinigrus Artigas, 1981: 187, figs. 1-7. Type-locality: Argentina, Córdoba, Salina Mascasín. Distr. – Argentina (Córdoba). Ref. – Artigas & Papavero, 1995c: 133. Type, DZUC.

albispinosus (Macquart), 1850: 388 (1850: 84), pl. 8, fig. 7 (only body; wing belongs to *longiterebratus* Macquart, 1850) (*Erax*). Type-locality: "South America". Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 233 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1995c: 133. Type, MNHNP.

amblayoensis (Hengst & Artigas), 2003: 290, figs. 1, 2B, E, G, 3A-H, 5. Type-locality: Argentina, Salta, Salta, Alturas de Amblayo, 3500 m. Distr. – Argentina (La Rioja, Salta, San Juan). Type, DZUC. N. comb.

tibialis Hengst & Artigas, 2003: 287, *nomen nudum*.

ablayoensis Hengst & Artigas, 2003: 291 (caption of fig. 3D), error.

borroi Artigas, 1970a: 288, figs. 261, 264, 268, 270, 496. Type-locality: Chile, Ñuble, Chillán. Distr. – Chile (Biobío, Concepción, Curicó, Linares, Malleco, Ñuble, Santiago, Valparaíso). Refs. – Artigas, 1971: 36, figs. 89-90; Artigas & Papavero, 1995c: 133, figs. 33-38. Type, DZUC.

borroi Schumann, 1973: 93, error.

cienagaensis (Hengst & Artigas), 2003: 293, figs. 1, 3D-E, G, 6A-F, 7. Type-locality: Argentina, San Juan, La Ciénaga. Distr. – Bolivia (Cochabamba), Argentina (Catamarca, Mendoza, San Juan, Tucumán). Type, DZUC. N. comb.

sagittata Hengst & Artigas, 2003: 287, *nomen nudum*.

griseus (Guérin-Méneville), 1838: 292 (*Asilus*). Type-locality: "Chile". Distr. – Chile (Arauco, Biobío, Concepción, Coquimbo, Curicó, Malleco, Maule, Ñuble, O'Higgins, Santiago, Valparaíso). Refs. – Macquart, 1838: 115 (*Erax*); Blanchard,

- 1852: 369, pl. 3, fig. 2 (*Erax*); Walker, 1855: 634 (*Erax*); Philippi, 1865: 692 (*Erax*); Schiner, 1868: 180 (*Erax*); Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 237 (cat.; *Erax*); Engel, 1930: 459, fig. 3 (n. comb.); Hull, 1962 (2): 496, figs. 206A, 716, 1437, 1446, 2203, 2217, 2494; Artigas, 1971: 37, figs. 87-88; Artigas & Reyes, 1975: 199, figs. 4-6, 12; Artigas & Papavero, 1995c: 134. Type, MNHNP.
- cinereus* Philippi, 1865: 693 (*Erax*). Type-locality: Chile, Santiago, Colchagua. Refs. – Williston, 1891: 85 (cat.; *Erax*); Kertész, 1909: 235 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*). Type, MNHNS.
- albescens* Schiner, 1868: 180 (*Erax*). Type-locality: “Chile”. Type, NMW. Ref. – Hull, 1962 (2): 478 (*Nerax*).
- leoninus* Artigas, 1970b: 67, figs. 1-4. Type-locality: Chile, Tarapacá, Mollepampa, Lluta. Distr. – Southern Peru, northern Chile (Tarapacá). Ref. – Artigas & Papavero, 1995c: 134. Type, DZUC.
- longiterebratus* (Macquart), 1850: 387 (1850: 83), pl. 8, fig. 7 (only wing; body belongs to *albispinosus* Macquart, 1850) (*Erax*). Type-locality: Argentina, Patagonia. Refs. – Lynch Arribálzaga, 1880: 120 (*Erax*), 1883a: 149 (cat.; *Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 238 (cat.; *Erax*); Engel, 1930: 460 (n. comb.); Bromley, 1932: 269 (*Erax*); Hull, 1962 (2): 496; Artigas & Papavero, 1995c: 134. Type, lost.
- mucidus* (Walker), 1837: 340 (*Asilus*). Type-locality: Argentina, Puerto Santa Elena. Distr. – Argentina (Chubut, Mendoza, Neuquén, Río Negro, Santa Cruz), Chile (Aysén). Refs. – Lynch Arribálzaga, 1883a: 151 (cat.; *Asilus*), 1883b: 86 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 260 (cat.; *Asilus*); Bromley, 1932: 279, figs. 23, 24a-c (*Philonerax*); Hull, 1962 (2): 495, figs. 212, 689, 705, 750, 1439, 1448, 2225 (*Philonerax*); Artigas, 1970a: 342, figs. 363-365, 411 (*Philonerax*); Artigas & Papavero, 1995c: 134, 18-24 (*Lochmorhynchus*); Hengst & Artigas, 2003: 289, figs. 1, 2A-E, 3C, F (*Philonerax*). Type, BMNH. N. comb.
- cribratus* Hull, 1962 (2): 498, figs. 209, 751, 1435, 1444, 2256, 2281, 2373, 2382 (*Lonchodogonus*). Type-locality: “Believed to be from the Cordilleran region of southern South America”. Refs. – Artigas & Papavero, 1995c: 134; Hengst & Artigas, 2003: 289 (*Philonerax*; syn.). Type, ZSBS.
- cribatus* Hengst & Artigas, 2003: 289, error.
- chilechicoensis* Artigas, 1970a: 344, figs. 366-370, 500 (*Philonerax*). Type-locality: Chile, Aysén, Chile Chico, laguna Buenos Aires. Distr. – Chile (Aysén). Refs. – Artigas, 1971: 47, figs. 120-121 (*Philonerax*); Artigas & Papavero, 1995c: 133; Hengst & Artigas, 2003: 289 (*Philonerax*; syn.). Type, DZUC.
- patagoniensis* (Macquart), 1850: 387 (1850: 83) (*Erax*). Type-locality: Argentina, Patagonia. Distr. – Argentina (Mendoza, Buenos Aires, Patagonia), Chile. Refs. – Schiner, 1866: 713 (*Erax*); Lynch Arribálzaga, 1880: 113 (*Erax*), 1883a: 149 (cat.; *Erax*); Williston, 1891: 86 (cat.; *Erax*); Kertész, 1909: 240 (cat.; *Erax*); Bromley, 1832: 269 (*Erax*); Hull, 1962 (2): 478 (*Nerax*); Artigas & Papavero, 1997b: 72 (*Nerax*); Tomasovic, 2002: 56 (n. comb.). Type, MNHNP.
- patagonensis* Artigas, 1970a: 282, figs. 249, 251, 413 (*Efferia*), error.
- puntarenensis* Artigas, 1970a: 294, figs. 262, 263b, 271, 272, 431. Type-locality: Chile, Magallanes, Punta Arenas. Distr. – Chile (Magallanes). Refs. – Artigas, 1971: 37, figs. 91-92; Artigas & Papavero, 1995c: 134. Type, DZUC.
- senectus* (Wulp), 1882b: 110 (*Proctacanthus*). Type-locality: “Argentina”. Refs. – Lynch Arribálzaga, 1883b: 13 (*Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 248 (cat.; *Proctacanthus*); Engel, 1930: 459, fig. 2 (n. comb.); Hull, 1962 (2): 496, figs. 732, 1420, 2363; Artigas & Papavero, 1995c: 134. Syntypes, ZMA (in bad shape, eaten by dermestids, the two ST without terminalia and only parts of the abdomen left).
- sp. Artigas & Papavero, 1995c: 143-144, figs. 41-48. Argentina.
- tumbrensis* (Artigas), 1970a: 285, figs. 253-260, 479 (*Eremomyia*). Type-locality: Chile, Antofagasta, Tumbre. Distr. – Chile (Antofagasta). Refs. – Artigas, 1971: 36, figs. 85-86 (*Eretomyia*); Artigas & Papavero, 1995c: 134, figs. 10-17 (n. comb.). Type, DZUC.

Mallophora-group

Promachina Lehr, 1969: 532, 549, *nomen nudum*. Ref. – Dikow, 2003: 461.

Genus **Amblyonychus** Hermann, 1921

- Promachus*, subg. **Amblyonychus** Hermann, 1921: 118. Type-species, *Promachus wiedemannii* Schiner, 1867 (orig. des.).
Promachina Bromley, 1934a (March): 96. Type-species, *Promachus trapezoidalis* Bellardi, 1861 (orig. des; based on specimens from U. S. A., Texas, Brownsville).
Promachina Bromley, 1934b (July): 352 (preocc. Bromley, 1934). Type-species, *Promachus trapezoidalis* Bellardi, 1861 (orig. des., misident.; based on specimens from Guyana). Ref. – Martin, 1967a.

annularis (Fabricius), 1805: 171 (*Dasyopogon*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Venezuela, Guyana, Peru (Huánuco, Lima), Bolivia. Refs.

- Wiedemann, 1821: 211 (*Asilus*), 1828: 492 (*Asilus*); Macquart, 1834; 311 (*Asilus*); Walker, 1849: 394 (*Asilus*), 1855: 589 (*Trupanea*); Schiner, 1866: 689 (?gen.); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 217 (cat.; *Promachus*); Curran, 1935: 10 (in key; *Promachina*); Carrera & d'Andretta, 1953: 69, fig. 1 (*Promachina*); Hull, 1962 (2): 462 (*Promachus*), 467 (*Promachina*); Carrera & Machado-Allison, 1963: 249, fig. 2a (*Promachina*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Lamas M., 1972: 314 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 99 (n. comb.). Syntypes, COPN, KIEL (in bad condition) [cf. Zimšen, 1964: 471].
- flavifasciatus** (Macquart), 1838: 102 (1839: 218), pl. 9, figs. 1, 1a, 1b (*Trupanea*). Type-locality: Brazil, Pará, surroundings of Belém. Distr. – Surinam, Brazil (Pará). Refs. – Walker, 1849: 394 (*Asilus*), 1855: 591 (*Trupanea*); Williston, 1891: 84 (cat.); Kertész, 1909: 220 (cat.); Bromley, 1946: 113 (cat.; *Promachus*); Carrera, 1960a: 164 (*Promachina*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 99 (n. comb.). Syntypes, MNHNP.
- gracilis** (Macquart), 1838: 102 (1839: 218) (*Trupanea*). Type-locality: “Brazil”. Refs. – Walker, 1955: 589 (*Trupanea*); Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 221 (cat.; *Promachus*); Bromley, 1946: 113 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 80 (cat.; *Promachus*); Artigas & Papavero, 1997c: 99 (n. comb.). Type, lost.
- horni** (Bromley), 1935b: 109, fig. 2 (*Promachus*). Type-locality: Peru, Huallaga, Jumbatis, 350 m. Distr. – Peru, Brazil (Acre), Bolivia. Refs. – Carrera & d'Andretta, 1950: 171 (*Promachus*, misident.), 1952: 294 (*Promachus*); Carrera, 1959: 4 (*Promachus*), 1960a: 164 (*Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin, 1965a: 33; Martin & Papavero, 1970: 81 (cat.; *Promachus (Promachus)*); Lamas M., 1972: 314 (cat.; *Promachus (Promachus)*); Artigas & Papavero, 1997c: 99 (n. comb.). Type, DEI.
- incisuralis** (Macquart), 1838: 107 (1839: 223) (*Trupanea*). Type-locality: unknown. Distr. – Brazil. Refs. – Macquart, 1846: 80 (*Trupanea*); Walker, 1855: 610 (*Trupanea*); Williston, 1891: 84 (cat.); Kertész, 1909: 221 (cat.; *Promachus*); Bromley, 1946: 114 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Promachus)*); Artigas & Papavero, 1997c: 100 (n. comb.; = *nigripes* Fabricius). Type, MNHNP.
- lateralis** (Walker), 1860: 283 (*Trupanea*). Type-locality: “Mexico”. Refs. – Kertész, 1909: 221 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 100 (n. comb.); Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.). Type, BMNH.
- nigripes** (Fabricius), 1787: 360 (*Asilus*). Type-locality: “French Guiana, Cayenne (Rohr)” [cf. Papavero, 1971e: 20]. Refs. – Olivier, 1789: 268 (*Asilus*); Gmelin, 1792: 2899 (*Asilus*); Fabricius, 1794: 385 (*Asilus*), 1805: 171 (*Dasyopogon*); Wiedemann, 1821: 209 (*Asilus*), 1828: 480 (*Asilus*); Rondani, 1848: 95 (*Mallophora*); Walker, 1855: 592 (*Trupanea*); Schiner, 1866: 688 (*Promachus*), 1867: 388 (*Promachus*); Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 223 (cat.; *Promachus*); Carrera, 1945a: 162 (*Promachina*; preys); Bromley, 1946: 114 (cat.; *Promachus*); Carrera, 1947d: 270 (*Promachina*; preys); Hull, 1962 (2): 467 (*Promachina*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 100 (n. comb.). Type, KIEL (only thorax and wings) [cf. Zimšen, 1964: 471].
- ?*heydenii* Wiedemann, 1828: 490 (*Asilus*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1855: 592 (*Trupanea*); Schiner, 1866: 589 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 257 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*). Type, SENCK [cf. Geller-Grimm, 1999: 209].
- ?*perfectus* Walker, 1851: 120 (*Trupanea*). Type-locality: “South America”. Refs. – Walker, 1855: 592 (*Trupanea*); Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 224 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 80 (cat.; *Promachus*). Type, BMNH (without abdomen; the colour of the hairs indicates a male).
- ?*breviusculus* Walker, 1855: 591 (*Trupanea*). Type-locality: Brazil, Pará, rio Tapajós. Refs. – Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 218 (cat.; *Promachus*); Bromley, 1946: 113 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 79 (cat.; *Promachus*). Syntypes, BMNH.
- ?*nigripes* Macquart, 1838: 101 (1839: 217) (*Trupanea*; preocc. Fabricius, 1787). Type-locality: Brazil, Pará, Belém. Refs. – Walker, 1855: 589 (*Trupanea*); Williston, 1891: 84 (cat.). Type, MNHNP.
- ?*macquartii* Rondani, 1848: 95 (*Mallophora*; nom. nov. for *nigripes* Macquart, 1837). Type-locality: Brazil, Pará, Belém. Refs. – Kertész, 1909: 222 (cat.); Bromley, 1929a: 287, pl. 1, fig. 10 (*Mallophora*); Curran, 1934a: 5 (in key); Bromley, 1946: 114 (cat.; *Promachus*; as syn. of *nigripes* Macquart, 1838); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 80 (cat.; *Promachus*). Type, MZUN.
- ovatus** (Martin), 1967a: 158 (*Promachus (Amblyonychus)*). Type-locality: Mexico, Colima, 6 miles south of Colima. Distr. – Mexico (Colima). Refs. – Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 100 (n. comb.). Type, KANS.
- pulchellus** (Bellardi), 1861: 129 (29), pl. 2, fig. 5 (*Promachus*). Type-locality: “Mexico”. Refs. – Aldrich, 1905: 280 (cat.; *Promachus*); Kertész, 1909: 224 (cat.; *Promachus*); Curran, 1935: 10 (in key; *Promachina*); Hull, 1962 (2): 467 (*Promachina*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 100 (n. comb.); Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.). Type, TORO.
- quatuorlineatus** (Macquart), 1838: 102 (1839: 218) (*Trupanea*). Type-locality: Brazil, Rio de Janeiro. Refs. – Walker, 1855: 592 (*Trupanea*); Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 224 (cat.; *Promachus*); Bromley, 1946: 114 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 80 (cat.; *Promachus*); Artigas & Papavero, 1997c: 100 (n. comb.). Type, lost.
- substitulus** (Walker), 1851: 119 (*Trupanea*). Type-locality: “South America”. Distr. – Brazil (São Paulo), Paraguay. Refs. –

- Walker, 1855: 591 (*Trupanea*); Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 226 (cat.; *Promachus*); Carrera & Vulcano, 1961: 74 (*Promachina*; prey); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 80 (cat.; *Promachus*); Artigas & Papavero, 1997c: 100 (n. comb.). Type, BMNH.
- mitescens** Walker, 1851: 118 (*Trupanea*). Type-locality: "South America". Refs. – Walker, 1855: 591 (*Trupanea*); Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 223 (cat.; *Promachus*); Carrera, 1958a: 150 (*Promachina*); Hull, 1962 (2): 462 (*Promachus*). Type, BMNH (this is the female of the species; the type of *substitutulus* Walker is the male).
- barbiellinii** Curran, 1935: 10 (in key), 11 (*Promachina*). Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 113 (cat.; *Promachina*); Carrera, 1947a: 45 (*Promachina*); Carrera & d'Andretta, 1952: 258 (*Promachina*) (prey); Hull, 1962 (2): 467 (*Promachina*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997: 100 (syn.). Type, AMNH.
- titan** (Carrera), 1959: 4 (*Promachina*). Type-locality: Bolivia, Chapare, rio Chipirí, 400 m. Distr. – Peru (Cuzco), Bolivia. Refs. – Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Lamas M., 1972: 314 (cat.; *Promachus (Amblyonychus)*); Fisher, 1985: 33 (n. comb.); Artigas & Papavero, 1997c: 100. Type, ZSBS.
- trapezoidalis** (Bellardi), 1861: 128 (28), pl. 2, fig. 4 (*Promachus*). Type-locality: "Mexico". Refs. – Williston, 1901: 320; Aldrich, 1905: 279 (*Mallophora*); Kertész, 1909: 215 (cat.; *Mallophora*); Bromley, 1934a: 96; Curran, 1935: 10 (in key); Bromley, 1946: 113 (cat.; *Promachina*); Carrera & d'Andretta, 1950: 169 (*Promachina*), 1953: 69 (*Promachina*; misident.); Hull, 1962 (2): 467, figs. 346, 744, 1383, 1392, 2190, 2217 (*Promachina*); Cole & Pritchard, 1964: 64 (*Promachus*); Martin, 1965a: 32; Martin & Wilcox, 1965: 399 (cat.; *Promachina*); Artigas & Papavero, 1997c: 100 (n. comb.); Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, ?
- trichonotus** (Wiedemann), 1828: 490, pl. 6, fig. 4 (*Asilus*). Type-locality: Brazil, Bahia, Salvador. Refs. – Loew, 1848: 405 (*Promachus*); Walker, 1855: 591 (*Trupanea*); Schiner, 1866: 689 (*Promachus*), 1867: 69 (*Promachus*); Wulp, 1882b: 108 (*Promachus*); Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 226 (cat.; *Promachus*); Engel, 1930: 458 (*Promachus (Trypanoides)*); Curran, 1935: 10 (in key; *Promachina*); Carrera, 1945a: 163 (*Promachina*; preys); Bromley, 1946: 113 (cat.; *Promachina*); Carrera, 1960a: 164 (*Promachina*); Hull, 1962 (2): 467 (*Promachina*); Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 100 (n. comb.). Type, NMW.
- wiedemanni** (Schiner), 1867: 388 (*Promachus*; nom. nov. for *nigripes* Fabricius of Wiedemann, 1828). Type-locality: Brazil, Bahia [i. e., Bahia, Salvador]. Refs. – Williston, 1891: 84 (cat.; *Promachus*); Kertész, 1909: 227 (cat.; *Promachus*); Bromley, 1946: 114 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*), 463 (*Promachus (Amblyonychus)*); Martin, 1965a: 32; Martin & Papavero, 1970: 81 (cat.; *Promachus*). Syntypes, NMW.
- nigripes* Fabricius of Wiedemann, 1828: 480 (*Asilus*), misident.
- yepezi** (Carrera & Machado-Allison), 1963: 250, fig. 2b (*Promachina*). Type-locality: Venezuela, Bolívar, Guyaraca, Auyantepuy, 1100 m. Distr. – Venezuela (Bolívar). Refs. – Martin & Papavero, 1970: 81 (cat.; *Promachus (Amblyonychus)*); Artigas & Papavero, 1997c: 100 (n. comb.). Type, FAUCV.

Genus **Carreraomyia** Cole, 1969

- Carreraia** Cole & Pritchard, 1964: 60 (preocc. Correa, 1947). Type-species, *Carreraia acapulquensis* Cole & Pritchard, 1964 (orig. des.).
- Carreraomyia** Cole, 1969: 543 (nom. nov. for *Carreraia* Cole & Pritchard, 1964). Type-species, *Carreraia acapulquensis* Cole & Pritchard, 1964 (aut.).

- acapulquensis** (Cole & Pritchard), 1964: 62, fig. 7 (*Carreraia*). Type-locality: Mexico, Guerrero, Acapulco. Distr. – Mexico (Guerrero, Oaxaca). Refs. – Martin & Papavero, 1970: 77 (cat.); Artigas & Papavero, 1997c: 100. Type, CAS.
- alpuyeca** (Cole & Pritchard), 1964: 63 (*Carreraia*). Type-locality: Mexico, Morelos, Alpuyeca. Distr. – Mexico (Morelos). Refs. – Martin & Papavero, 1970: 77 (cat.); Artigas & Papavero, 1997c: 100. Type, CAS.

Genus **Mallophora** Macquart, 1838

- Mallophora** Macquart, 1834: 300. Type-species, *Asilus bombooides* Wiedemann, 1821 (sub. des., Coquillett, 1910: 565).

Ref. – Artigas & Angulo, 1980 (rev.).

- aeaca** Williston, 1901: 319. Type-locality: Mexico, Veracruz, Atoyac. Distr. – Mexico (Guerrero, Veracruz, Yucatán), El Salvador, French Guiana, Brazil (Acre, Pará), Bolivia (Cochabamba, Santa Cruz), Paraguay, Uruguay, Argentina (Catamarca, Córdoba, Entre Ríos, Jujuy, Mendoza, Misiones, Salta, Tucumán). Refs. – Aldrich, 1905: 278 (cat.); Kertész, 1909: 209 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 27, fig. 125;

- Artigas & Papavero, 1997c: 100; Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.
- duellojuradoi* Gemignani, 1930: 136. Type-locality: Argentina, Jujuy. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 27 (syn.). Type, MACN.
- annuliventris* Artigas & Angulo, 1980: 29, fig. 126. Type-locality: Guatemala, Sacatepéquez, Capetillo. Distr. – Guatemala (Sacatepéquez). Ref. – Artigas & Papavero, 1997c: 100. Type, AMNH.
- antiqua* Walker, 1855: 581. Type-locality: Mexico, Veracruz. Refs. – Kertész, 1909: 209 (cat.); Martin & Papavero, 1970: 72 (cat.). Hull, 1962 (2): 468; Artigas & Papavero, 1997: 105 (unrecogn.); Papavero & Ibáñez Bernal, 2001: 95 (hist. Mexican dipterol.). [Species ignored by Artigas & Angulo, 1980].
- ?*ardens* Macquart, 1834: 362. Type-locality: “North America”. Distr. – unknown. Refs. – Macquart, 1838: 89, pl. 8, fig. 2; Walker, 1855: 576; Aldrich, 1905: 278 (cat.); Kertész, 1909: 209 (cat.); Hull, 1962 (2): 468; Artigas & Papavero, 1997c: 100. *ardens* Macquart of Artigas & Angulo, 1980: 29, figs. 122, 127 is most probably misident., they designated as NT a specimen from Brazil, Rio de Janeiro, Rio de Janeiro (Copacabana), in the MZUSP.
- argentipes* Macquart, 1838: 38 (1839: 204). Type-locality: “Guyana”. Distr. – Panama, Colombia, Venezuela, Guyana, Trinidad-Tobago, Surinam, French Guiana, Brazil (Roraima, Rio Grande do Norte, Mato Grosso, Goiás, Rio de Janeiro, São Paulo, Santa Catarina), Paraguay, Argentina (Buenos Aires, Formosa, Salta, Santa Fe, Santiago del Estero). Refs. – Williston, 1891: 82 (cat.); Kertész, 1909: 208 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 31, figs. 20, 25, 75-76, 122, 128; Artigas & Papavero, 1997c: 101. Type, MNHNP.
- armata* Artigas & Angulo, 1980: 33, figs. 17, 122, 129. Type-locality: Paraguay, Caaguazú. Distr. – Paraguay. Ref. – Artigas & Papavero, 1997c: 101. Type, DZUC.
- atra* Macquart, 1834: 302. Type-locality: “Brazil”. Neotype-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – U. S. A. (Minnesota, Florida), Mexico (Guerrero, Morelos), Honduras, Colombia, Venezuela, Guyana, Surinam, French Guiana, Trinidad-Tobago, Ecuador, Peru, Brazil (Roraima, Rondônia, Amazonas, Pará, Amapá, Paraíba, Pernambuco, Bahia, Mato Grosso, Mato Grosso do Sul, Tocantins, Goiás, Distrito Federal, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul), Bolivia (Cochabamba, Santa Cruz), Paraguay, Argentina (Buenos Aires, Chaco, Córdoba, Formosa, Jujuy, Misiones, Salta, Santa Fe, Santiago del Estero, Tucumán), Chile (Arica). Refs. – Walker, 1849: 389 (*Asilus*), 1855: 583; Schiner, 1866: 711; Williston, 1891: 82 (cat.); Carrera, 1945a: 161 (preys); Bromley, 1946: 112 (cat.); Carrera & Vulcano, 1961: 74 (preys); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 34, figs. 92-93, 122, 130; Artigas & Papavero, 1997c: 101; Dennis, Barnes & Knutson, 2008: 234 (pupal case). Type, lost. NT DZUC [des. by Artigas & Angulo, 1980: 34].
- freycineti* Macquart, 1838: 85 (1839: 201). Type-locality: “Brazil” (probably Rio de Janeiro, the only place where Freycinet landed in Brazil; cf. Papavero, 1971e: 124). Refs. – Macquart, 1846: 77; Walker, 1855: 579; Schiner, 1866: 711; Williston, 1891: 82 (cat.), 1901: 319; Aldrich, 1905: 278 (cat.); Bristowe, 1925: 483 (as *infernalis* var. *freycineti*; “bears a very striking resemblance to a big black Humble Bee, *Bremus kohli* (*carbobarius* Handl.), occurring in the same locality”); Carrera, 1945a: 162 (preys), 1947d: 270 (preys); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469 (as syn. of *infernalis* Wiedemann, 1821); Martin, 1965a: 34; Martin & Papavero, 1970: 74 (cat.); Artigas, 1970a: 310; Artigas & Angulo, 1980: 34 (syn.). Type, MNHNP.
- ventralis* Macquart, 1838: 91 (1839: 207). Type-locality: unknown. Refs. – Walker, 1855: 580; Schiner, 1866: 711; Williston, 1891: 84 (cat.); Kertész, 1909: 216 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, MNHNP.
- albifrons* Walker, 1850a: xcvi. Type-locality: “South America”. Refs. – Walker, 1855: 579; Schiner, 1866: 710; Williston, 1891: 82 (cat.); Kertész, 1909: 209 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, lost.
- testaceipes* Macquart, 1850: 380 (1850: 76). Type-locality: “South America”. Refs. – Schiner, 1866: 710; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, OXF.
- xylocopides* Walker, 1851: 111. Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, lost.
- xyocopides* Walker, 1855: 579; Schiner, 1866: 711; Williston, 1891: 83 (cat.), error.
- testaceitarsis* Macquart, 1854: 74 (1854: 54). Trype-locality: “South America”. Refs. – Schiner, 1866: 710; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, lost.
- coeruleiventris* Thomson, 1869: 467. Type-locality: Peru, Callao. Refs. – Williston, 1891: 82 (cat.); Kertész, 1909: 211 (cat.); Engel, 1930: 457; Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, ?lost.
- nigra* Williston, 1885: 58. Type-locality: U. S. A., Minnesota. Refs. – Coquillett, 1893c: 118; Curran, 1930a: 12 (in key); Bromley, 1950: 235; Cole & Pritchard, 1960: 53; Hull, 1962 (2): 468; Martin & Wilcox, 1965: 396 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, lost.
- lynchi* Gemignani, 1930: 141. Type-locality: Argentina, Jujuy. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, MACN.
- flavicollis* Curran 1941: 272 (in key), *nomen nudum*.
- nitidula* Curran, 1941: 272 (in key), *nomen nudum*.

- vorax* Curran 1941: 272 (in key), 278. Type-locality: Peru, Lima. Refs. – Carrera & d'Andretta, 1953: 67 (allotype des.); Carrera & Machado-Allison, 1963: 247; Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, AMNH.
- lucida* Curran, 1941: 272 (in key), 278. Type-locality: Argentina, Posadas. Refs. – Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, AMNH.
- curiosa* Curran, 1941: 272 (in key), 279. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 34 (syn.). Type, AMNH.
- jemima* Curran, 1941: 272 (in key), 279. Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Artigas & Angulo, 1980: 34 (syn.). Type, AMNH.
- jemina* Martin & Papavero, 1970: 74 (cat.), error.
- barbipes** (Wiedemann), 1819: 48 (*Laphria*). Type-locality: "Brazil". Neotype-locality: Bolivia, Beni, Rurrenabaque. Distr. – French Guiana, Ecuador (Napo, Oriente), Peru (Cuzco, Huánuco, Loreto), Brazil (Pará, Mato Grosso, Goiás, Bahia), Bolivia (Beni, Sara), Paraguay (Guairá), Argentina (Formosa, Salta, Tucumán). Refs. – Wiedemann, 1821: 239 (*Laphria*), 1824: 60 (*Asilus*), 1828: 481 (*Asilus*); Walker, 1855: 590 (*Trupanea*); Schiner, 1866: 688; Williston, 1891: 82 (cat.); Kertész, 1909: 210 (cat.); Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 40, figs. 114-115, 121-122, 130; Fisher, 1984: 33; Artigas & Papavero, 1997c: 101. Type, lost. NT DZUC [des. by Artigas & Angulo, 1980: 41].
- semiviolacea* Rondani, 1848: 92. Refs. – Schiner, 1866: 711; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 40 (syn.). Type, MZUN.
- cilicrura* Rondani, 1850: 867. Type-locality: "Equatorial America". Refs. – Osten Sacken, 1891: 420; Kertész, 1909: 210 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 40 (syn.). Type, MZUN.
- subviolacea* Rondani, 1850: 369, error.
- purpurea* Walker, 1850a: xcvi (Trupanea). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Walker, 1855: 584; Williston, 1891: 83, 84 (*Promachus*); Kertész, 1909: 214 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 40 (syn.).
- auromystacea* Macquart, 1855: 75 (1855: 55). Type-locality: "South America". Refs. – Williston, 1891: 82 (cat.); Kertész, 1909: 209 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 40 (syn.). Type, lost.
- campestris* Curran, 1941: 270 (in key), 274. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 40 (syn.). Type, AMNH.
- dana* Curran, 1941: 270 (in key), 274. Type-locality: Peru, río Marañón. Refs. – Carrera & d'Andretta, 1953: 68; Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 40 (syn.). Type, AMNH.
- cora* Curran, 1941: 272 (in key), 280. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 40 (syn.). Type, AMNH.
- bassleri** Curran, 1941: 270 (in key), 273. Type-locality: Brazil, Amazonas, Tefé. Distr. – Peru (Huánuco), Brazil (Amazonas, Santa Catarina). Refs. – Bromley, 1946: 112 (cat.); Carrera & d'Andretta, 1953: 68; Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Lamas M., 1972: 313 (cat.); Artigas & Angulo, 1980: 43, figs. 122, 131; Artigas & Papavero, 1997c: 101. Type, AMNH.
- bellengeri** Farr, 1965: 26, figs. 11-12. Type-locality: Jamaica, St. Andrew, Long Mountain. Distr. – Jamaica. Refs. – Artigas & Angulo, 1980: 44, figs. 122, 132; Artigas & Papavero, 1997c: 101. Type, USNM.
- bellingeri* Martin & Papavero, 1970: 73 (cat.), error.
- bigotii** Lynch Arribálzaga, 1883a: 146. Type-locality: Uruguay, Montevideo. Neotype-locality: Argentina, Buenos Aires, Tigre. Distr. – Brazil (Santa Catarina), Uruguay, Argentina (Buenos Aires, Junín). Refs. – Lynch Arribálzaga, 1883b: 6; Williston, 1891: 82 (cat.); Osten Sacken, 1891: 419; Kertész, 1909: 210 (cat.); Brèthes, 1908: 287 (cat.); Kertész, 1909: 210 (cat.); Gemignani, 1930: 240; Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 46, figs. 81-82, 122, 133; Artigas & Papavero, 1997c: 101. Type, ? NT DZUC [des. by Artigas & Angulo, 1980: 46].
- soccata* Thomson of Lynch Arribálzaga, 1880: 256, misident. Ref. – Kertész, 1909: 210 (cat.).
- bomboides** (Wiedemann), 1821: 203 (*Asilus*). Type-locality: U. S. A., Georgia. Distr. – U. S. A. (Florida, Georgia, North Carolina, Pennsylvania), Mexico (Veracruz). Refs. – Wiedemann, 1828: 476 (*Asilus*); Macquart, 1834: 302, 1838: 89; Walker, 1849: 387 (*Asilus*), 1855: 576; Schiner, 1866: 687; Williston, 1885: 57; Coquillett, 1893c: 118; Aldrich, 1905: 278 (cat.); Kertész, 1905: 278 (cat.); Fattig, 1945: 26; Bromley, 1950: 228, 234; Brower, Brower & Westcott, 1960: 94 (mimic of *Bombus pensylvanicus* (De Geer, 1773) [as *Bombus americanorum* Fabricius, 1804], Hymenoptera, Apidae); Hull, 1962 (2): 468; Cole & Pritchard, 1964: 51; Martin & Wilcox, 1965: 396 (cat.); Artigas & Papavero, 1997c: 101. Type, BZM [no. 421; cf. Schumann, 1973: 93].
- trichostica* Williston, 1901: 320. Type-locality: Mexico, Veracruz, Atoyac. Refs. – Aldrich, 1905: 279 (cat.); Kertész, 1909: 215 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 48 (syn.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.
- rex* Bromley, 1925: 192. Lectotype-locality: U. S. A., South Carolina, Southern Pines. Refs. – Cresson Jr., 1926: 91 (Lectotype des.); Curran, 1930a: 12 (in key); Bromley, 1950: 235; Hull, 1962 (2): 468, fig. 2206, 2231; Cole & Pritchard, 1964: 51 (syn.); Martin & Wilcox, 1965: 397 (cat.); Artigas & Angulo, 1980: 48. Lectotype ANSP [des. by

- Cresson Jr., 1926: 91].
- chrysomela* Bromley, 1925: 193. Type-locality: U. S. A., Georgia, Atlanta. Refs. – Cresson Jr., 1926: 91 (Lectotype des.); Curran, 1930a: 12 (in key); Fattig, 1945: 26; Hull, 1962 (2): 468; Cole & Pritchard, 1964: 51 (syn.); Martin & Wilcox, 1965: 396 (cat.); Artigas & Angulo, 1980: 48. Lectotype ANSP [des. by Cresson Jr., 1926: 91].
- cheysomela* Cresson Jr., 1926: 91, error.
- calida** (Fabricius), 1787: 358 (*Asilus*). Type-locality: “French Guiana, Cayenne (Rohr)” [cf. Papavero, 1971e: 20]. Neotype-locality: Bolivia, Cordillera, San Antonio. Distr. – Mexico (Veracruz), Colombia (Santander), Venezuela, Guyana, French Guiana, Peru (San Marytín), Brazil (Amazonas, Pará, Mato Grosso, Mato Grosso do Sul, Goiás, Ceará, Rio Grande do Norte, Paraíba, Bahia, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul), Bolivia (Beni, Cordillera, Sara), Paraguay (Guairá), Uruguay, Argentina (Chaco, Jujuy, Salta, Santa Fe, Santiago del Estero). Refs. – Olivier, 1789: 265 (*Asilus*); Gmelin, 1792: 2897 (*Asilus*); Fabricius, 1794: 380 (*Asilus*), 1805: 159 (*Laphria*); Wiedemann, 1821: 210 (*Asilus*), 1828: 482 (*Asilus*); Walker, 1855: 590 (*Trupanea*); Schiner, 1866: 688, 1867: 383; Williston, 1891: 82 (cat.); Kertész, 1909: 210 (cat.); Bromley, 1934: 350; Carrera, 1960a: 162; Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 48, figs. 28, 38, 58-59, 122, 235; Artigas & Papavero, 1997c: 102. Type, KIEL (only indeterminable remains) [cf. Zimsen, 1964: 468]. NT DZUC [des. by Artigas & Angulo, 1980: 49].
- nigritarsis* Fabricius of Macquart, 1838: 86, misident.
- albicincta* Bromley, 1929b: 47. Type-locality: Mexico, Veracruz, Coatepec. Refs. – Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 49 (syn.). Type, USNM.
- callida* Curran, 1934a: 5 (in key), 1941: 271 (in key), error.
- abana* Curran, 1934a: 5 (in key), 7. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Curran, 1941: 271 (in key); Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 49 (syn.). Type, AMNH.
- clavitarsis* Curran, 1941: 271 (in key), 282. Type-locality: Peru, San Martín, Achinamiza. Refs. – Bromley, 1946: 112 (cat.); Carrera & d'Andretta, 1952: 257 (prey); Carrera, 1960a: 162; Carrera & Vulcano, 1961: 74 (prey); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 49 (syn.). Type, AMNH.
- candens** (Walker), 1851: 117 (*Trupanea*). Type-locality: “South America”. Refs. – Walker, 1855: 583; Williston, 1891: 82 (cat.); Kertész, 1909: 210 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Papavero, 1997c: 105 (unrecogn.).
- cingulata* Artigas & Angulo, 1980: 53, figs. 2, 122, 136, 176. Type-locality: Peru, Pucallpa, río Ucayali. Distr. – Panama, Colombia, Peru. Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- circumflava* Artigas & Angulo, 1980: 54, figs. 86-89, 122, 137. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. – Brazil ((Rio de Janeiro, São Paulo, Santa Catarina)). Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- clavipes* Curran, 1941: 270, 271 (in key), 281. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Distr. – Panama, Venezuela (Bolívar), Guyana, French Guiana, Ecuador, Peru (Huánuco), Brazil (Amazonas, Pará, Amapá, Mato Grosso, Rio de Janeiro). Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 55, figs. 18, 30, 112-113, 116-117, 122, 139; Fisher, 1984: 33; Artigas & Papavero, 1997c: 102. Type, AMNH.
- cortesi* Artigas & Angulo, 1980: 57, figs. 122, 138, 175. Type-locality: Mexico, Sonora, Álamos, Estrella. Distr. – Mexico (Sonora). Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- cubana* Artigas & Papavero, 1997c: 102 (*nom. nov.* for *macquarti* Rondani of Artigas & Angulo, 1980). Type-locality: Cuba, Soledad. Distr. – Cuba. Type, DZUC.
- macquarti* Rondani of Artigas & Angulo, 1980: 77, figs. 22, 31-32, 54-55, 122, 154. misident.
- crocuscopha* Artigas & Angulo, 1980: 58, figs. 23-24, 52-53, 122, 140. Type-locality: Paraguay, Guairá, Villarica. Distr. – Bolivia (Santa Cruz, Sara), Paraguay (Caaguazú, Guairá, Paraguarí), Brazil (Mato Grosso, Goiás, Minas Gerais, Rio de Janeiro, Paraná, Rio Grande do Sul), Argentina (Chaco, Formosa, Tucumán). Type, DZUC.
- umbrina* Artigas & Angulo, 1980: 60, *nomen nudum*.
- dureti* Artigas & Angulo, 1980: 60, figs. 37, 83-85, 122, 141. Type-locality: Argentina, Salta, Orán. Distr. – Brazil (Rio Grande do Norte), Bolivia (Santa Cruz, Sara), Paraguay, Argentina (Catamarca, Córdoba, Formosa, La Rioja, Río Negro, Salta, Santa Fe, Santiago del Estero, Tucumán). Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- emiliae* Carrera, 1960b: 48. Type-locality: Brazil, Paraná, Curitiba. Distr. – Brazil (Mato Grosso, Paraná, Santa Catarina, Rio Grande do Sul), Uruguay. Refs. – Carrera & Vulcano, 1961: 74 (preys); Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 63, figs. 60-61, 122, 142; Artigas & Papavero, 1997c: 102. Type, MZUSP.
- fautrix* Osten Sacken, 1887: 191, pl. 3, fig. 14. Type-locality: Mexico, Sinaloa, Presidio, near Mazatlán. Distr. – U. S. A. (Arizona, California, New Mexico, Texas, Utah), Mexico (Chihuahua, Morelos, Nayarit, Puebla, Sinaloa, Veracruz), French Guiana. Refs. – Coquillett, 1893c: 118; Cockerell, 1894: 173; Snow, 1896: 186; Williston, 1901: 320; Aldrich, 1905: 278 (cat.); Kertész, 1909: 211 (cat.); Hine, 1918: 5; Curran, 1930a: 12 (in key); Hull, 1962 (2): 469; Cole & Pritchard, 1964: 57; Martin & Papavero, 1970: 74 (cat.); Alcock, 1974: 68 (behaviour); Artigas & Angulo, 1980: 64, figs. 62-63, 122, 143; Artigas & Papavero, 1997c: 102; Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.
- fautricoides* Curran, 1930a: 14. Type-locality: U. S. A., California, San Diego. Refs. – Hull, 1962 (2): 468; Cole & Pritchard, 1964: 58 (as *fautrix* ssp.); Martin & Wilcox, 1965: 396 (cat.); Hogue, 1974: 100, fig.; Artigas & Angulo,

- 1980: 64 (syn.). Type, AMNH.
- bromleyi** Curran, 1930a: 13. Type-locality: U. S. A., Texas, Marathon. Refs. – Bromley, 1934: 104, 1935: 3; Hull, 1962 (2): 468, fig. 729, 1369; Cole & Pritchard, 1964: 59 (as *fautrix* ssp.); Martin & Wilcox, 1965: 396 (cat.); Artigas & Angulo, 1980: 64 (syn.). Type, AMNH.
- fritzii** Artigas & Angulo, 1980: 66, figs. 122, 144. Type-locality: Uruguay, Montevideo [Sello; cf. Papavero, 1971e: 71]. Distr. – Brazil (Rio Grande do Sul), Uruguay. Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- fulviventris** Macquart, 1850: 381 (1850: 77). Type-locality: “Mexico”. Distr. – U. S. A. (Utah), Mexico. Refs. – Schiner, 1866: 711; Loew in Osten Sacken, 1878: 77 (cat.); Osten Sacken, 1887: 191; Aldrich, 1905: 278 (cat.); Kertész, 1909: 211 (cat.); Brown, 1929: 313, pl. 30, figs. 42–45; Bromley, 1934a: 104, Bromley, 1935c: 3; Hull, 1962 (2): 468; Martin & Wilcox, 1965: 396 (cat.); Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 100; Artigas & Papavero, 1997: 105 (unrecogn.); Artigas & Papavero, 1997c: 105 (as unrecognized). Papavero & Ibáñez Bernal, 2001: 86 (hist. Mexican dipterol.). ?TP MNHNP (a male; Macquart says female!).
- ?**fulvianalis** Macquart, 1850: 382 (1850: 78). Type-locality: “Mexico”. Refs. – Schiner, 1866: 711; Loew in Osten Sacken, 1878: 78 (cat.); Osten Sacken, 1887: 191; Aldrich, 1905: 278 (cat.); Kertész, 1909: 211 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 100; Papavero & Ibáñez Bernal, 2001: 86 (hist. Mexican dipterol.). Type, MNHNP.
- gracipes** Artigas & Angulo, 1980: 66, figs. 33, 122, 145, 176. Type-locality: Brazil, Santa Catarina, Caúna. Distr. – Brazil (Santa Catarina, Rio Grande do Sul). Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- hemivitrea** Artigas & Angulo, 1980: 68, figs. 122, 146. Type-locality: Brazil, Espírito Santo, Viana. Distr. – Brazil (Espírito Santo). Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- inca** Curran, 1941: 270 (in key), 273. Type-locality: Peru, río Santiago. Distr. – Colombia, Peru, Brazil (Mato Grosso, Paraná), Paraguay. Refs. – Carrera & d’Andretta, 1953: 88; Hull, 1962 (2): 469; Lamas M., 1972: 314 (cat.); Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 68, figs. 122, 147; Artigas & Papavero, 1997c: 102. Type, AMNH.
- incanipes** Artigas & Angulo, 1980: 69, figs. 122, 148. Type-locality: Brazil, Amazonas, Tabatinga. Distr. – Peru, Brazil (Amazonas). Refs. – Fisher, 1984: 33; Artigas & Papavero, 1997c: 102. Type, DZUC.
- infernalis** (Wiedemann), 1821: 202, 244 (*Asilus*). Type-locality: “Brazil”. Neotype-locality: Brazil, Rio de Janeiro, Nova Friburgo. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo), Argentina (Misiones). Refs. – Wiedemann, 1828: 475, pl. 6, fig. 7 (*Asilus*); Perty, 1833: 181, pl. 36, fig. 5 (*Asilus*); Macquart, 1834: 301, pl. 7, fig. 7; Walker, 1849: 387 (*Asilus*), 1855: 579; Bellardi, 1861: 21; Schiner, 1866: 687; Wulp, 1882b: 106; Osten Sacken, 1887: 189; Williston, 1891: 82 (cat.); Kertész, 1909: 211 (cat.); Curran, 1934a: 5 (in key), 1941: 270 (in key); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 468, fig. 194; Martin & Papavero, 1970: 74 (cat.); Artigas & Papavero, 1997c: 102. Type, lost. NT DZUC [des. by Artigas & Angulo, 1980: 70].
- leschenaulti** Macquart, 1838: 86 (1839: 202). Type-locality: French Guiana, La Mana river (as Lamana). Distr. – Mexico (Chihuahua, Colima, Guanajuato, Guerrero, Jalisco, Morelos, Nuevo León, Oaxaca, Tamaulipas, Veracruz), Venezuela, Trinidad-Tobago, Guyana, Surinam, French Guiana, Ecuador, Peru, Brazil (Pará, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Espírito Santo, Rio de Janeiro, Rio Grande do Sul), Paraguay, Argentina (Córdoba, Jujuy, Misiones). Refs. – Walker, 1855: 580; Schiner, 1866: 711; Williston, 1891: 83 (cat.); Kertész, 1909: 212 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 72, figs. 36, 50–51, 122, 153; Artigas & Papavero, 1997c: 102. Type, MNHNP.
- breviventris** Macquart, 1850: 381 (1850: 70), pl. 7, figs. 7, 7a. Type-locality: “Brazil”. Distr. – Brazil (Rio de Janeiro). Refs. – Schiner, 1866: 711; Williston, 1891: 82 (cat.), 1901: 321; Aldrich, 1905: 210 (cat.); Kertész, 1909: 210 cat.); Curran, 1934a: 5 (in key), 1941: 270 (in key); Bromley, 1946: 112 (cat.); Carrera, 1960a: 163; Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 72 (syn.). Type, MNHNP.
- belzebul** Schiner, 1866: 711 (*nomen nudum*), 1867: 385. Type-locality: “Brazil”. Refs. – Williston, 1891: 82 (cat.); Kertész, 1909: 210 (cat.); Bromley, 1934a: 104, 1946a: 112 (cat.); Hull, 1962 (2): 468; Cole & Pritchard, 1964: 60; Martin, 1965a: 34; Martin & Wilcox, 1965: 396 (cat.); Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 72 (syn.). Type, NMW.
- zottai** Gemignani, 1930: 141. Type-locality: Argentina, Jujuy. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 72 (syn.). Type, MACN.
- speciosa** Curran, 1934: 1. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Refs. – Bromley, 1934b: 349; Curran, 1934a: 5 (in key), 1941: 270 (in key); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 72 (syn.). Type, AMNH.
- leucopyga** Artigas & Angulo, 1980: 74, figs. 122, 150. Type-locality: Mexico, Guerrero, Acapulco. Distr. – Mexico (Guerrero). Ref. – Artigas & Papavero, 1997c: 102. Type, DZUC.
- leucothrica** Carrera & d’Andretta, 1953: 66. Type-locality: Peru, Lima. Distr. – Panama, Peru. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 74, figs. 19, 26, 40, 122, 151; Artigas & Papavero, 1997c: 102. Type, MZUSP.
- lugubris** Lynch Arribálzaga, 1880: 252. Type-locality: Argentina, Formosa. Distr. – Brazil (Mato Grosso, Mato Grosso do Sul, São Paulo), Bolivia (Santa Cruz, Yungas), Paraguay (Guairá), Argentina (Formosa, Jujuy). Refs. – Lynch Arribálzaga, 1883a: 146 (cat.); Williston, 1891: 83 (cat.); Brèthes, 1908: 187 (cat.); Kertész, 1909: 212 (cat.); Gemignani, 1930: 142; Curran, 1934a: 4 (in key), 272 (in key); Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 75, figs. 71–74, 122, 152; Artigas & Papavero, 1997c: 103. Type, MACN.

- macquarti** Rondani, 1850b: 367. Type-locality: "Equatorial America". Distr. – Cuba, Brazil. Refs. – Loew in Osten Sacken, 1878: 78, 233 (cat.; note), 1891: 420; Kertész, 1909: 212; Bromley, 1929: 187; Curran, 1930a: 12 (in key); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.; misident. ?); Artigas & Papavero, 1997c: 105 (as unrecognized). Type, MZUN.
- scopifera** Wiedemann of Macquart, 1838: 89 (1839: 205) (Cuba), Walker, 1855: 577; Rondani, 1850: 367, Bigot, 1857: 790; Jaennicke, 1867: 362; misident. Refs. – Martin & Papavero, 1970: 75 (cat.), Artigas & Angulo, 1980: 77.
- media** Clements & Bennett, 1969: 455 (also descr. of pupa). Type-locality: Trinidad, Curepe. Distr. – Trinidad-Tobago. Refs. – Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 79, figs. 122, 155; Artigas & Papavero, 1997c: 103. Type, BMNH.
- minos** (Wiedemann), 1824: 478 (*Asilus*). Type-locality: Uruguay, Montevideo. Distr. – Mexico (Distrito Federal, Jalisco, Nayarit, Oaxaca, Tabasco, Veracruz), Belize, Guatemala, El Salvador, Panama, Colombia, Venezuela, Trinidad-Tobago, Ecuador, Peru, Brazil (São Paulo), Uruguay. Refs. – Wiedemann, 1828: 478 (*Asilus*); Walker, 1855: 579; Schiner, 1866: 688; Wulp, 1882b: 106; Williston, 1891: 83 (cat.); Brèthes, 1908: 287 (cat.); Kertész, 1909: 212; Gemignani, 1930: 144; Curran, 1934a: 4 (in key), 1941: 272 (in key); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75; Artigas & Angulo, 1980: 80, figs. 29, 90-91, 122, 156; Fisher, 1984: 33; Artigas & Papavero, 1997c: 103.
- craverii** Bellardi, 1861: 122 (22). Type-locality: Mexico, surroundings of Mexico City, and Oaxaca. Refs. – Schiner, 1866: 711; Osten Sacken, 1887: 191; Williston, 1901: 320; Aldrich, 1905: 278 (cat.); Kertész, 1909: 211 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 80 (syn.). Type, TORO.
- pulchella** Curran, 1925a: 260. Type-locality: Colombia, Cauca. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 80 (syn.). Type, ANSP.
- schwarzi** Curran, 1934a: 4 (in key), 5. Type-locality: Panama, Canal Zone, Barro Colorado Island. Refs. – Curran, 1941: 271, 272 (in key); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 80 (syn.); Shelly & Weinberger, 1981: 380 (courtship and diet). Type, AMNH.
- tolteca** Curran, 1941: 272 (in key), 277. Type-locality: "Belize" (as British Honduras). Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 80 (syn.). Type, AMNH.
- pipiens** Curran, 1941: 272 (in key), 277. Type-locality: "Surinam". Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 80 (syn.). Type, AMNH.
- nigrifemorata** Macquart, 1838: 90 (1839: 206). Type-locality: "?Brazil". Distr. – Panama, Colombia, Guyana, Surinam, Ecuador, Brasil (Amazonas, Rio de Janeiro, São Paulo, Santa Catarina), Argentina (Buenos Aires). Refs. – Walker, 1855: 582; Williston, 1891: 83 cat.); Kertész, 1909: 213 (cat.); Hull, 1962 (2): 469 (as ?syn. of *nigritarsis* Fabricius, 1805); Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 82, figs. 64-66, 122, 157; Artigas & Papavero, 1997c: 103. Type, MNHNP.
- tibialis** Macquart, 1846: 77, pl. 8, fig. 3 (preocc. Macquart, 1838). Type-locality: "Guyana". Refs. – Walker, 1855: 580; Carrera & d'Andretta, 1953: 68.
- cruralis** Rondani, 1863: 46 (nom. nov. for *tibialis* Macquart, 1846). Type-locality: "Guyana". Refs. – Williston, 1891: 82 (cat.); Kertész, 1909: 211 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 82 (syn.). Type, MZUN.
- ada** Curran, 1941: 171 (in key), 184. Type-locality: Brazil, Rio de Janeiro. Refs. – Bromley, 1946: 112 (cat.); Carrera, 1960a: 162; Carrera & Vulcano, 1961: 74 (preys); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 82 (syn.). Type, AMNH.
- geijskesi** Curran, 1942: 58. Type-locality: Surinam, Brownsberg. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 82 syn.). Type, AMNH.
- fairchildi** Curran, 1942: 58. Type-locality: Panama, Coclé, El Valle. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 82 (syn.). Type, AMNH.
- nigritarsis** (Fabricius), 1805: 168 (*Dasytopogon*). Type-locality: "South America (Smidt)" [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Cuba, Panama, Colombia, Venezuela (Bolívar, Carabobo), Guyana, Surinam, French Guiana, Trinidad-Tobago, Peru (Loreto), Ecuador (Los Ríos, Oriente), Brazil (Amazonas, Pará, Mato Grosso, Pernambuco, Minas Gerais, Santa Catarina), Bolivia (Beni, Sara), Paraguay, Argentina (Salta, Tucumán). Refs. – Wiedemann, 1821: 208 (*Asilus*), 1828: 479 (*Asilus*); Macquart, 1838: 86; Walker, 1849: 389, 1855: 582; Schiner, 1866: 688, 1867: 386; Wulp, 1882a: 77, 1882b: 107; Williston, 1891: 83 (cat.); Kertész, 1909: 213 (cat.); Bromley, 1934b: 350; Curran, 1934a: 5 (in key), 1941: 271 (in key); Carrera, 1945a: 160 (preys); Bromley, 1946: 113 (cat.); Carrera, 1947d: 270 (preys); Carrera & d'Andretta, 1952: 237 (preys); Carrera, 1960a: 162; Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 85, figs. 69-70, 122, 158; Artigas & Papavero, 1997c: 103. Type, COPN [cf. Zimsen, 1964: 470. [NT from Bolivia, Chapare, 400 m DZUC, des. by Artigas & Angulo, 1980: 85, invalid].
- cinerascens** Walker, 1855: 581. Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Schiner, 1866: 711; Williston, 1891: 82 (cat.); Kertész, 1909: 210 (cat.); Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 85 (syn.).
- socculata** Rondani, 1848: 93. Type-locality: "Brazil" [i. e., Belém, Pará; cf. Papavero, 1973d: 342]. Refs. – Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.). Type, MZUN.

- bruneri*** Bromley, 1929a: 287. Type-locality: Cuba, Sierra Rangel, Pinar del Río. Refs. – Curran, 1930a: 12 (in key); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 85 (syn.); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.). Type, USNM.
- fusca*** Bromley, 1934b: 350. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Refs. – Curran, 1934a: 5 (in key), 1941: 271 (in key); Carrera, 1960a: 162; Hull, 1962 (2): 469; Carrera & Machado-Allison, 1963: 247; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 85 (syn.). Type, AMNH.
- orcina*** (Wiedemann), 1828: 477 (*Asilus*). Type-locality: “U. S. A., Savannah”. Distr. – U. S. A. (Alabama, Arizona, Florida, Georgia, Indiana, North Carolina, Oklahoma, South Carolina, Texas), Mexico (Hidalgo, Puebla, Sonora). Refs. – Walker, 1855: 576; Schiner, 1866: 687, 1868: 711; Loew in Osten Sacken, 1878: 78 (cat.); Williston, 1885: 58; Aldrich, 1905: 279 (cat.); Kertész, 1909: 213 (cat.); Dozier, 1920: 371; Curran, 1930a: 12 (in key); Bromley, 1931c: 12; Fattig, 1933: 119 (feeding); Bromley, 1934: 104; Fattig, 1945: 27; Bromley, 1947: 68, 1950a: 231, 1950b: 234; Goslin, 1950: 305; Cole & Pritchard, 1960: 54; Hull, 1962 (2): 468, fig. 2429, 2431; Martin & Wilcox, 1965: 396 (cat.); Artigas & Angulo, 1980: 88, figs. 104-105, 122, 159; Alayo Dalmau & Garcia Ávila, 1983: 50 (cat.); Bambara, 1983: 143; Artigas & Papavero, 1997c: 103.
- ciliata*** Walker, 1851: 114. Type-locality: unknown. Refs. – Walker, 1855: 581; Schiner, 1866: 710; Williston, 1891: 82 (cat.); Kertész, 1909: 210 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 88 (syn.).
- fulva*** Banks, 1911: 130. Type-locality: U. S. A., Arizona. Refs. – Curran, 1930a: 12 (in key); Hull, 1962 (2): 468; Cole & Pritchard, 1964: 55; Martin & Wilcox, 1965: 396 (cat.); Artigas & Angulo, 1980: 88 (syn.). Type, ?
- papaveroi*** Artigas & Angulo, 1980: 90, figs. 3, 96-99, 122, 160. Type-locality: Brazil, Amazonas, rio Urupadi. Distr. – Brazil (Acre, Amazonas, Pará, Mato Grosso, Minas Gerais, Tocantins). Ref. – Artigas & Papavero, 1997c: 103. Type, DZUC.
- parasylveirii*** Artigas & Angulo, 1980: 91, figs. 77-78, 122, 161. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (Rio de Janeiro, São Paulo, Santa Catarina). Ref. – Artigas & Papavero, 1997c: 103. Type, DZUC.
- pica*** Macquart, 1850: 382 (1850: 78). Type-locality: “Mexico”. Distr. – Bolivia (Sara), Uruguay, Argentina (Buenos Aires, Córdoba, San Luís). Refs. – Schiner, 1866: 711; Osten Sacken, 1887: 191; Snow, 1896: 186; Aldrich, 1905: 279 (cat.); Kertész, 1909: 213 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 92, figs. 106-107, 122, 162 [No specimens seen from Mexico; probably misident.]; Ref. – Artigas & Papavero, 1997c: 103; Papavero & Ibáñez Bernal, 2001: 86 (hist. Mexican dipterol.). Type, MNHNP.
- ?***bergii*** Lynch Arribálzaga, 1880: 257. Type-locality: “Uruguay”. Refs. – Lynch Arribálzaga, 1883a: 146 (cat.), 1883b: 10; Williston, 1891: 82 (cat.); Osten Sacken, 1891: 419; Brèthes, 1908: 287; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 93 (syn.).
- pluto*** (Wiedemann), 1828: 477 (*Asilus*). Type-locality: “Brazil”. Neotype-locality: Argentina, Sierra Córdoba. Distr. – Colombia, Brazil (Goiás, Paraná, Santa Catarina, Rio Grande do Sul), Uruguay, Argentina (Chaco, Córdoba, La Rioja, Misiones, Salta, Santa Fe, Tucumán). Refs. – Walker, 1849: 387 (*Asilus*), 1855: 579; Schiner, 1866: 688, 1868: 176; Wulp, 1882b: 106; Lynch Arribálzaga, 1882b: 142, 1883b: 5; Osten Sacken, 1887: 190; Williston, 1891: 83 (cat.); Aldrich, 1905: 279 (cat.); Kertész, 1909: 214 (cat.); Curran, 1934a: 4 (in key), 271 (footnote); Carrera, 1945a: 162 (prey); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 94, figs. 27, 39, 45-46, 49, 122, 163; Ref. – Artigas & Papavero, 1997c: 103. Type, ? NT DZUC [des. by Artigas & Angulo, 1980: 94].
- vegeta*** Lynch Arribálzaga, 1882b: 140. Type-locality: Argentina, Santa Fe. Refs. – Osten Sacken, 1891: 420; Kertész, 1909: 216 (cat.); Engel, 1930: 457; Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 94 (syn.). Type, MACN.
- copelloi*** Gemignani, 1930: 138. Type-locality: Argentina, La Rioja. Refs. – Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 94 (syn.). Type, MACN.
- pusilla*** Macquart, 1838: 88 (1839: 204). Type-locality: “Brazil, Rio Grande do Sul, eastern part (as : Capitainerie de Rio Grande)”. Refs. – Walker, 1855: 582; Williston, 1891: 83 (cat.); Kertész, 1909: 214 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Papavero, 1997c: 105 (as unrecognized). [Species ignored by Artigas & Angulo, 1980].
- robusta*** (Wiedemann), 1828: 478 (*Asilus*). Type-locality: unknown. Neotype-locality: Brazil, São Paulo, Boa Esperança do Sul (Fazenda Itaquerê). Distr. – Mexico (Veracruz), Guadeloupe, Venezuela, Guyana, Surinam, French Guiana, Peru, Ecuador, Brazil (Amazonas, Pará, Mato Grosso, Goiás, Rio de Janeiro, São Paulo), Bolivia (Beni, Santa Cruz, Sara, Yungas), Paraguay, Argentina (Chaco, Jujuy). Refs. – Macquart, 1846: 78; Walker, 1855: 580; Schiner, 1866: 688; Loew in Osten Sacken, 1878: 78 (cat.); Wulp, 1882b: 106; Lynch Arribálzaga, 1882b: 142; Wulp, 1882b: 106; Osten Sacken, 1887: 191; Williston, 1891: 83 (cat.); Aldrich, 1905: 279 (cat.); Kertész, 1909: 214 (cat.); Bromley, 1934b: 351; Curran, 1934a: 4 (in key), 1941: 272 (in key); Carrera, 1945a: 161 (preys); Bromley, 1946: 113 (cat.); Carrera, 1960a: 163; Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 96, figs. 47-48, 122, 164, 175; Fisher, 1984: 33; Artigas & Papavero, 1997c: 103. Type, ? NT DZUC [des. by Artigas & Angulo, 1980: 97].
- tisiphones*** Rondani, 1848: 94. Type-locality: “Brazil” [i. e., Pará, Belém; cf. Papavero, 1973d: 342]. Refs. – Rondani, 1863: 47; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.). Type, MZUN.
- alecto*** Rondani, 1848: 94. Type-locality: “Brazil” [i. e., Pará, Belém; cf. Papavero, 1973d: 342]. Refs. – Schiner, 1866: 711; Williston, 1891: 82 (cat.); Kertész, 1909: 209 (cat.); Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 209 (cat.). Artigas & Angulo, 1980: 97 (syn.). Artigas & Angulo, 1980: 97 (syn.). Type, MZUN.

- lampon** Walker, 1849: 388 (*Asilus*). Type-locality: unknown. Refs. – Walker, 1855: 582; Schiner, 1866: 711; Williston, 1891: 82 (cat.); Kertész, 1909: 212 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 97 (syn.).
- hirtiventris** Macquart, 1850: 383 (1850: 79), pl. 8, fig. 1 (*Trupanea*). Type-locality: “Bolivia (d’Orbigny)” [cf. Papavero, 1973d: 139-141]. Refs. – Williston, 1891: 84 (cat.); Kertész, 1909: 221 (cat.); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 80 (*Promachus*); Artigas & Angulo, 1980: 97 (syn.). Type, MNHNP.
- rufipes** Macquart, 1850: 380 (1850: 76). Type-locality: “Brazil”. Refs. – Schiner, 1866: 711; Williston, 1891: 83 (cat.); Curran, 1941: 272 (in key); Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 97 (syn.). Type, OXF.
- tissiphones** Schiner, 1866: 711, error.
- dallasi** Gemignani, 1931: 265. Type-locality: Argentina, Chaco, Formosa. Refs. – Hull, 1962 (2): 468; Carrera & Machado-Allison, 1963: 247 (?misident.); Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 97 (syn.). Type, ?MACN.
- aria** Curran, 1941: 272 (in key), 276. Type-locality: Peru, río Santiago. Refs. – Carrera & d’Andretta, 1953: 67; Carrera, 1960a: 163; Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Lamas M., 1972: 314; Artigas & Angulo, 1980: 97 (syn.). Type, AMNH.
- aletō** Martin & Papavero, 1970: 72 (cat.), error.
- hertiventris** Artigas & Angulo, 1980: 97, error.
- ruficauda** (Wiedemann), 1828: 476 (*Asilus*). Type-locality: Uruguay, Montevideo. Neotype-locality: Argentina, Buenos Aires, Devoto. Distr. – Mexico (Guerrero, Yucatán), Brazil (Amapá, Paraná, Santa Catarina, Rio Grande do Sul), Paraguay, Uruguay, Argentina (Buenos Aires, Córdoba, Entre Ríos, Jujuy, Santa Fe, Tucumán). Refs. – Walker, 1855: 580; Schiner, 1866: 687, 1867: 386; Lynch Arribálzaga, 1880: 230; Wulp, 1882b: 106; Lynch Arribálzaga, 1882b: 140, 1883a: 146 (cat.); Williston, 1891: 83 (cat.); Brèthes, 1908: 288 (cat.); Kertész, 1909: 214 (cat.); Copello, 1922: 1 (biol.), 1927: 699 (biol.); Bromley, 1929a: 287; Gemignani, 1930: 136; Curran, 1934a: 5 (in key), 1941: 270 (in key); Copello, 1942; Carrera, 1960b: 48; Hull, 1962 (2): 469; Remedi de Gavotto, 1964: 151 (larvae preying upon *Cyclocephala signaticollis* Burmeister, 1847 (Coleoptera, Scarabaeidae)); Crouzel, 1965 (biol.); Martin & Papavero, 1970: 75 (cat.); Naveiro, 1975 (biol.); Tricca & Trujillo Cenóz, 1980: 627 (sexual behavior); Rabinovich & Corley, 1997 (biol.); Rabinovich, Quiroga & Castelo, 1997 (biol.); Artigas & Angulo, 1980: 100; Trica & Trujillo-Cenóz, 1980 (biol.); Dennis & Knutson, 1988: 858, 861, figs. 4A-C, 8 (pupa in ventral, lateral and dorsal views; host: *Cyclocephala signaticollis* Burmeister, 1857 (Coleoptera, Scarabaeidae)); De Santis, 1989: 16 (biol.); De Santis & Cornejo, 1990 (biol.); Artigas & Papavero, 1997c: 104; Marcangeli, 1998 (biol.); Castelo & Capurro, 2000 (biol.); Castelo, 2001a, 2001b (biol.); Castelo & Capurro, 2000 (biol.); Castelo, 2002a, 2002b, 2004a (biol.); Castelo & Corley, 2004a, 2004b (biol.); Castelo & Lazzari, 2004 (biol.); Castelo, Ney-Nifle, Corley & Bernsatein, 2006 (biol.). Type, ? NT DZUC [des. by Artigas & Angulo, 1980: 100].
- pyrura** Rondani, 1868: 47. Type-locality: unknown. Refs. – Osten Sacken, 1891: 420; Kertész, 1909: 214 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 100 (syn.). Type, MZUN.
- soccata** Thomson, 1869: 467. Type-locality: Argentina, Buenos Aires. Refs. – Lynch Arribálzaga, 1883a: 146 (cat.); Hull, 1962 (2): 469; Artigas & Angulo, 1980: 100 (syn.).
- rufiventris** Macquart, 1838: 91 (1839: 207). Type-locality: unknown. Distr. – Colombia, French Guiana, Brazil (Pernambuco, Mato Grosso, Goiás, São Paulo, Paraná, Santa Catarina), Bolivia (Sara), Paraguay, Argentina (Formosa, Misiones, Salta, Tucumán). Refs. – Walker, 1855: 580; Schiner, 1866: 711; Lynch Arribálzaga, 1882b: 143; Williston, 1891: 83 (cat.); Kertész, 1909: 214 (cat.); Curran, 1934a: 5 (in key), 1941: 270 (in key); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 184; Artigas & Papavero, 1997c: 104. Type, MNHNP.
- ?**nigriventris** Jaennicke, 1867: 361. Type-locality: “Paraguay”. Refs. – Lynch Arribálzaga, 1880: 254, 1882b: 143, 1883a: 119 (cat.); Williston, 1891: 83 (cat.); Brèthes, 1908: 287 (cat.); Kertész, 1909: 213 (cat.); Gemignani, 1930: 143; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 104 (type given as “lost”, error). Type, SENCK [cf. Geller-Grimm, 1999: 212; several differences found in relation to *rufiventris* Macquart, 1838].
- scopifera** (Wiedemann), 1828: 487 (*Asilus*). Type-locality: “Brazil”. Distr. – Brazil (Minas Gerais, Paraná, Santa Catarina) Refs. – Walker, 1855: 577; Schiner, 1866: 688; Jaennicke, 1867: 362; Schiner, 1868: 176; Osten Sacken, 1878: 78, 233 (cat.); Lynch Arribálzaga, 1883b: 9; Williston, 1891: 83 (cat.); Brèthes, 1908: 214 (cat.); Kertész, 1909: 215 (cat.); Engel, 1930: 458; Curran, 1941: 270 (in key); Bromley, 1946: 113 (cat.); Carrera, 1947a: 45, 1960a: 161; Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 106, figs. 34, 38, 108-109, 122, 167; Artigas & Papavero, 1997c: 104. Syntypes, BZM [no. 478; cf. Schumann, 1973: 156].
- scopipeda** Rondani, 1863: 46. Type-locality: “Brazil”. Refs. – Lynch Arribálzaga, 1883b: 9; Williston, 1891: 83 (cat.); Aldrich, 1905: 279 (cat.); Artigas & Angulo, 1980: 72; Artigas & Papavero, 1997c: 105 (as unrecognized). Type, MZUN.
- scopitarsis** Rondani, 1863: 46. Type-locality: Brazil, Pará, Belém. Refs. – Lynch Arribálzaga, 1883: 9; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 72 (syn.); Artigas & Papavero, 1997c: 105 (as unrecognized). Type, MZUN.
- singularis** Macquart, 1838: 87 (1839: 203), pl. 8, fig. 4. Type-locality: Brazil, Paraná, Guaratuba. Refs. – Walker, 1855: 582; Schiner, 1867: 387; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Papavero, 1997c: 105 (as unrecognized). [Species ignored by Artigas & Angulo, 1980]. Type, lost.
- sylveirii** Macquart, 1838: 86 (1839: 202). Type-locality: “Brazil” [probably Rio de Janeiro, Rio de Janeiro; cf. Papavero,

- 1973d: 144-147]. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Santa Catarina, Rio Grande do Sul), Bolivia (Sara, Yungas), Paraguay, Uruguay, Argentina (Chaco, Formosa, Misiones). Refs. – Walker, 1855: 581; Schiner, 1866: 711; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 107, figs. 67-68, 122, 168; Dennis & Knutson, 1988: 856, figs. 3A-C (pupa in dorsal, lateral and ventral views; *Dyscinetus rugifrons* (Burmeister, 1847) (Coleoptera, Scarabaeidae, Dynastinae)); Artigas & Papavero, 1997c: 104. Type, MNHNP (headless).
- geniculata* Macquart, 1838: 87 (1839: 203). Type-locality: Brazil, Rio Grande do Sul, São Leopoldo. Refs. – Walker, 1855: 582; Schiner, 1866: 711, 1867: 384; Williston, 1891: 82 (cat.); Kertész, 1909: 211 (cat.); Bromley, 1946: 113 (cat.); Hull, 1962 (2): 469; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 107 (syn.). Type, MNHNP.
- opposita* Walker, 1851: 112. Type-locality: “Brazil”. Refs. – Walker, 1855: 581; Schiner, 1867: 385; Williston, 1891: 83 (cat.); Kertész, 1909: 2134 (cat.); Carrera, 1945a: 160, fig. 1 (preys); Bromley, 1946: 113, pl. 5, fig. 1 (cat.); Carrera & d’Andretta, 1952: 257 (preys); Carrera, 1958a: 149, 1960a: 163; Carrera & Vulcano, 1961: 72 (preys); Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 107 (syn.).
- contraria* Walker, 1851: 113. Type-locality: “South America”. Refs. – Walker, 1855: 579; Schiner, 1866: 711; Williston, 1891: 82 (cat.); Kertész, 1909: 211 (cat.); Curran, 1941: 272 (in key); Carrera, 1945a: 161 (preys), 1947d: 270 (preys); Carrera & d’Andretta, 1952: 257 (preys); Carrera, 1960a: 163; Carrera & Vulcano, 1961: 73 (preys); Hull, 1962 (2): 468; Martin & Papavero, 1970: 73 (cat.); Artigas & Angulo, 1980: 107 (syn.).
- affinis* Schiner, 1866: 711 (*nomen nudum*), 1868: 176. Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Refs. – Williston, 1891: 82 (cat.); Kertész, 1909: 209 (cat.); Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 107 (syn.). Type, NMW.
- scutellaris* Lynch Arribálzaga, 1880: 254. Type-locality: Argentina, Misiones. Refs. – Lynch Arribálzaga, 1882b: 143, 1883a: 121 (cat.); Williston, 1891: 83 (cat.); Brèthes, 1908: 288 (cat.); Kertész, 1909: 215 (cat.); Gemignani, 1930: 138; Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 107 (syn.). Type, ?MACN.
- antica* Curran, 1934a: 5 (in key), 6. Type-locality: Brazil, São Paulo. Refs. – Curran, 1941: 270 (in key); Bromley, 1946: 112 (cat.); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 107 (syn.). Type, AMNH.
- sexualis* Curran, 1934c: 331. Type-locality: Brazil, São Paulo, Campos do Jordão. Refs. – Curran, 1941: 270 (in key); Bromley, 1946: 113 (cat.); Carrera & Vulcano, 1961: 72 (preys); Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 107 (syn.). Type, AMNH.
- ajax* Curran, 1941: 270 (in key), 275. Type-locality: “Brazil”. Refs. – Bromley, 1946: 112 (cat.); Carrera, 1947d: 270 (preys); Carrera & Vulcano, 1961: 73 (preys); Hull, 1962 (2): 468; Martin & Papavero, 1970: 72 (cat.); Artigas & Angulo, 1980: 107 (syn.). Type, AMNH.
- tertiavitrea* Artigas & Angulo, 1980: 117, figs. 122, 169. Type-locality: Brazil, Mato Grosso, Cuiabá. Distr. – Brazil (Mato Grosso, Minas Gerais). Ref. – Artigas & Papavero, 1997c: 104. Type, DZUC.
- thompsoni* Artigas & Angulo, 1980: 118, figs. 1, 122, 170. Type-locality: Paraguay, P. G. Lageado, Voarido [?]. Distr. – Paraguay. Ref. – Artigas & Papavero, 1997c: 104. Type, DZUC.
- tibialis* Macquart, 1838: 85 (1939: 201). Type-locality: Brazil, Pará, surroundings of Belém; French Guiana, “dans les grands bois aux bords de l’Oyapock”, and Cayenne. Distr. – Colombia, Venezuela, Guyana, Surinam, French Guiana, Peru (Loreto), Brazil (Amazonas, Pará), Bolivia (Sara), Argentina (La Rioja). Refs. – Rondani, 1848: 93; Walker, 1849: 387 (*Asilus*); Schiner, 1866: 711; Wulp, 1882b: 107; Williston, 1891: 83 (cat.); Kertész, 1909: 215 (cat.); Curran, 1934a: 4 (in key), 1941: 271 (in key); Bromley, 1946: 113 (cat.); Carrera & d’Andretta, 1953: 68; Carrera, 1960a: 164; Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 118, figs. 21, 35, 110-111, 118-120, 122, 171, 175; Artigas & Papavero, 1997c: 104. Syntypes, MNHNP.
- fascipennis* Macquart, 1850: 379 (1850: 75). Type-locality: “Brazil”. Refs. – Williston, 1891: 82 (cat.); Carrera, 1960a: 164; Martin & Papavero, 1970: 74 (cat.); Artigas & Angulo, 1980: 118 (syn.). Type, lost.
- fasciata* Walker, 1851: 112. Type-locality: Brazil, Pará [i. e., Pará, Belém]. Ref. – Walker, 1855: 583; Schiner, 1866: 711; Williston, 1891: 82 (cat.); Martin & Papavero, 1970: 76 (cat.); Artigas & Angulo, 1980: 118 (syn.).
- tsacasi* Artigas & Angulo, 1980: 121, figs. 122, 172. Type-locality: “Mexico, Púrpú[s sic]” [Karl August Purpus was the collector, not a locality; cf. Sousa Sánchez, 1969]. Distr. – Mexico, ?Bolivia. Ref. – Artigas & Papavero, 1997c: 104. Type, DZUC.
- wilhelmi* Artigas & Angulo, 1980: 121, figs. 122, 173. Type-locality: México, Veracruz, Cotaxtla, Estación Experimental de Cotaxtla. Distr. – Mexico (Veracruz). Ref. – Artigas & Papavero, 1997c: 104. Type, DZUC.
- zita* Curran, 1941: 270, 271 (in key), 282. Type-locality: Peru, Achinamiza. Distr. – México (Nuevo León), Panama, Venezuela, Peru, Surinam, French Guiana, Ecuador, Brazil (Roraima, Amazonas, Pará, Minas Gerais, São Paulo), Bolivia (Beni), Paraguay, Uruguay, Argentina (Chaco). Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 76 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 122, figs. 79-80, 122, 174; Artigas & Papavero, 1997c: 104. Type, AMNH.
- neta* Curran, 1941: 271 (in key), 283. Type-locality: Peru, río Santiago. Refs. – Carrera, 1960a: 162; Hull, 1962 (2): 469; Carrera & Machado-Allison, 1963: 246; Martin & Papavero, 1970: 75 (cat.); Lamas M., 1972: 314 (cat.); Artigas & Angulo, 1980: 122 (syn.). Type, AMNH.
- mexicana* Carrera & d’Andretta, 1950: 167. Type-locality: Mexico, Nuevo León, Villa Santiago. Refs. – Hull, 1962 (2): 469; Martin & Papavero, 1970: 75 (cat.); Artigas & Angulo, 1980: 122 (syn.). Type, FMNH.

Genus **Megaphorus** Bigot, 1857

Megaphorus Bigot, 1857b: 542. Type-species, *Mallophora heteroptera* Macquart, 1838 (orig. des.) = *laphroides* (Wiedemann, 1828).

Mallophorina Curran, 1934c: 183. Type-species, *Mallophora guildiana* Williston, 1885 (orig. des.).

acrus (Curran), 1931b: 21 (*Mallophora*). Type-locality: U. S. A., Oklahoma, Woods County. Distr. – U. S. A. (Oklahoma, Texas, Kansas, New Mexico, Arizona), Mexico (Coahuila). Refs. – Baker, 1939: 168 (*Mallophora (Mallophorina)*); Hull, 1962 (2): 468 [*Mallophora*], 469 (*Mallophora (Megaphorus)*); Cole & Pritchard, 1964: 79 (*Mallophorina*); Martin & Wilcox, 1965: 397 (cat.; *Mallophorina*).

durangoensis (Cole) in Cole & Pritchard, 1964: 82 (*Mallophorina*). Type-locality: Mexico, Durango, 23 miles northeast of Durango, 6300 ft. Distr. – Mexico (Durango). Refs. – Martin & Papavero, 1970: 77 (cat.; *Mallophorina*); Artigas & Papavero, 1997c: 105. Type, CAS.

frustus flavidus (Cole) in Cole & Pritchard, 1964: 87 (*Mallophorina*). Type-locality: U. S. A., California, San Diego Co., Scissors Spring. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Martin & Papavero, 1970: 77 (cat.; *Mallophorina*); Artigas & Papavero, 1997c: 105. Type, CAS.

guildianus (Williston), 1885: 60 (*Mallophora*). Type-locality: U. S. A., California, Los Angeles Co. Distr. – U. S. A. (California, Utah), Mexico (Baja California). Refs. – Coquillett, 1893: 118 (*Mallophora*); Aldrich, 1905: 278 (cat.; *Mallophora*); Kertész, 1909: 211 (cat.; *Mallophora*); Hine, 1918: 6; Brown, 1929: 313, pl. 30, figs. 46-47 (*Mallophora*); Curran, 1930b: 21 (in key; *Mallophora*); Pritchard, 1935: 10 (in key; *Mallophorina*); Hull, 1962 (2): 469, figs. 727, 1368, 1377 (*Mallophora (Megaphorus)*); Cole & Pritchard, 1964: 73; Martin & Wilcox, 1965: 397 (cat.; *Mallophorina*); Martin & Papavero, 1970: 77 (cat.; *Mallophorina*); Dennis & Lavigne, 1976 (pupa); Artigas & Papavero, 1997c: 105. Type, USNM.

megachile (Coquillett), 1893a: 118 (*Mallophora*). Type-locality: U. S. A., California, Los Angeles County. Distr. – U. S. (California), Mexico (Baja California). Refs. – Hull, 1962 (2): 468 (*Mallophora*); Cole & Pritchard, 1964: 85 (*Mallophorina*); Martin & Wilcox, 1965: 397 (cat.; *Mallophorina*); Martin & Papavero, 1970: 77 (cat.; *Mallophorina*). **N. comb.**

prudens (Pritchard), 1935: 10 (*Mallophorina*). Type-locality: U. S. A., Arizona, Oracle. Distr. – Mexico (Coahuila). Refs. – Baker, 1939: 168 (*Mallophora (Mallophorina)*); Hull, 1962 (2): 468 (*Mallophora*), 469 (*Mallophora (Megaphorus)*); Cole & Pritchard, 1964: 80 (*Mallophorina*); Martin & Wilcox, 1965: 397 (cat.; *Mallophorina*). Syntypes, AMNH. **N. comb.**

willistoni (Cole) in Cole & Pritchard, 1964: 74, (*Mallophorina*). Type-locality: U. S. A., California, Lassen County, Hallelujah Junction. Distr. – U. S. A. (California), Mexico. Type, CAS. **N. comb.**

Genus **Promachella** Cole & Pritchard, 1964

Promachella Cole & Pritchard, 1964: 65. Type-species, *Promachina pilosa* Wilcox, 1937 (orig. des.).

pilosa (Wilcox), 1937: 43 (*Promachina*). Type-locality: U. S. A., Arizona, Sabino Canyon, Catalina Mts. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Hull, 1962 (2): 467 (*Promachina*); Cole & Pritchard, 1964: 66 (n. comb.); Martin & Wilcox, 1965: 399 (cat.; *Promachina*); Martin & Papavero, 1970: 77 (cat.); Artigas & Papavero, 1997c: 115. Type, CAS.

Genus **Promachus** Loew, 1848

Trupanea Macquart, 1838: 91 (1839: 209) (preocc. Guettard, 1762). Type-species, *Asilus maculatus* Fabricius, 1775 (orig. des.).

Promachus Loew, 1848: 390. Type-species, *Asilus maculatus* Fabricius, 1775 (sub. des., Coquillett, 1910: 595).

albifacies Williston, 1885: 63. Type-locality: U. S. A., Arizona. Distr. – U. S. A. (Arizona, California, Colorado, New Mexico), Mexico (northern Sonora). Refs. – Osten Sacken, 1887: 195; Aldrich, 1905: 279 (cat.); Kertész, 1909: 216 (cat.); Hine, 1911a: 168; Hull, 1962 (2): 461; Martin & Wilcox, 1965: 400 (cat.); Martin & Papavero, 1970: 79 (cat.).

anceps Osten Sacken, 1887: 194. Type-locality: “Mexico”, and Panama, Bugaba, 800-1500 ft. Refs. – Williston, 1901: 321; Aldrich, 1905: 279 (cat.); Kertész, 1909: 216 (cat.); Hull, 1962 (2): 462; Martin & Papavero, 1970: 79 (cat.); Artigas & Papavero, 1997c: 105; Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Syntypes, BMNH.

fuscipennis Macquart of Bellardi, 1861: 24, pl. 11, fig. 1 as var. A), misident.

bellardii Martin, 1965b: 132 (nom. nov. for *quadratus* Bellardi, 1861). Type-locality: “Mexico”. Distr. – Mexico (Veracruz). Refs. – Martin & Papavero, 1970: 79 (cat.); Artigas & Papavero, 1997c: 105.

quadratus Bellardi, 1861: 127 (27), pl. 2, fig. 3 (preocc. Wiedemann, 1828). Type-locality: “Mexico”. Refs. – Osten

- Sacken, 1878: 79, 234 (cat.), 1887: 184; Kertész, 1909: 220 (cat.; as syn. of *forfex* Osten Sacken, 1887); Hull, 1962 (2): 462; Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, TORO.
- captans** (Walker), 1851: 117 (*Trupanea*). Type-locality: "South America". Refs. – Walker, 1855: 590; Williston, 1891: 84 (cat.); Kertész, 1909: 218 (cat.); Hull, 1962 (2): 462; Martin & Papavero, 1970: 79 (cat.).
- cinctus** Bellardi, 1861: 125 (25), pl. 2, fig. 2. Type-locality: "Mexico". Distr. – Mexico (Nuevo León, Veracruz), Guatemala, Honduras, Nicaragua. Refs. – Osten Sacken, 1887: 193; Williston, 1901: 321; Aldrich, 1905: 280 (cat.); Kertész, 1909: 218 (cat.); Carrera & d'Andretta, 1950: 169, figs. 1, 14; 59–61; Hull, 1962 (2): 462; Martin & Papavero, 1970: 80 (cat.); Artigas & Papavero, 1997c: 105; Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, TORO.
- forfex** Osten Sacken, 1887: 194. Type-locality: Costa Rica, Irazu, 6000–7000 m. Distr. – Mexico (Veracruz), Guatemala, Costa Rica, Guyana. Refs. – Williston, 1901: 321; Aldrich, 1905: 280 (cat.); Kertész, 1909: 220 (cat.); Hine, 1911a: 167; Hull, 1962 (2): 462; Martin & Papavero, 1970: 80 (cat.); Artigas & Papavero, 1997: 105. Type, BMNH.
- fuscipennis** (Macquart), 1846: 209 (1846: 81), pl. 8, fig. 4 (*Trupanea*). Type-locality: "Colombia, warm region". Distr. – Colombia, Venezuela (Aragua). Refs. – Walker, 1855: 589 (*Trupanea*); Schiner, 1868: 177; Osten Sacken, 1878: 78, 234 (cat.), 1887: 195; Williston, 1891: 84 (cat.); Aldrich, 1905: 280 (cat.); Kertész, 1909: 221 (cat.); Hull, 1962 (2): 462; Carrera & Machado-Allison, 1963: 252; Martin, 1965a: 32; Martin & Papavero, 1970: 80 (cat.); Artigas & Papavero, 1997c: 105. Syntypes, MNHNP, OXF.
- giganteus** Hine, 1911a: 172. Type-locality: U. S. A., Texas, El Paso. Distr. – U. S. A. (Texas), Mexico (Coahuila). Refs. – Bromley, 1935c: 3; Baker, 1939: 168; Hull, 1962 (2): 461; Martin & Wilcox, 1965: 400 (cat.); Morgan & Shelly, 1988: 419 (body temperature regulation). Type, OSU.
- magnus** Bellardi, 1861: 126 (26). Type-locality: "Mexico". Distr. – U. S. A. (Texas); Mexico (Coahuila, Nuevo León). Refs. – Aldrich, 1905: 280 (cat.); Kertész, 1909: 133 (cat.); Hine, 1918: 6; Baker, 1939: 168; Carrera & d'Andretta, 1950: 171; Hull, 1962 (2): 462; Martin & Wilcox, 1965: 400 (cat.); Martin & Papavero, 1970: 80 (cat.); Artigas & Papavero, 1997c: 105; Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, TORO.
- minusculus** Hine, 1911: 171. Type-locality: U. S. A., New Mexico, near Albuquerque. Distr. – U. S. A. (New Mexico, Texas), Mexico. Ref. – Martin & Wilcox, 1965: 400 (cat.). Type, OSU.
- nigrialbus** Martin in Martin & Papavero, 1970: 80 (*nom. nov.* for *nigripes* Hine, 1911). Type-locality: U. S. A., New Mexico, Albuquerque. Distr. – U. S. A. (New Mexico), Mexico (Nuevo León).
- nigripes* Hine, 1911a: 170 (preocc. Macquart, 1838). Type-locality: U. S. A., New Mexico, Albuquerque. Ref. – Brown, 1929: 315, pl. 29, figs. 30–31, pl. 30, figs. 50–53. Type, OSU.
- nobilis** Osten Sacken, 1887: 196. Type-locality: Costa Rica, Caché. Distr. – Costa Rica, Venezuela (D. F., Mérida). Refs. – Aldrich, 1905: 280 (cat.); Kertész, 1909: 223 (cat.); Hull, 1962 (2): 462; Carrera & Machado-Allison, 1963: 252; Martin & Papavero, 1970: 80 (cat.); Artigas & Papavero, 1997c: 105. Type, BMNH.
- oklahomensis** Pritchard, 1935: 12. Type-locality: U. S. A., Oklahoma, Kenton. Distr. – U. S. A. (Oklahoma), Mexico (Coahuila). Refs. – Bromley, 1935c: 3; Baker, 1939: 168.
- painteri** Bromley, 1934a: 93. Type-locality: U. S. A., Texas, Austin. Distr. – U. S. A. (Texas), Mexico (Coahuila). Ref. – Baker, 1939: 168; Hull, 1962 (2): 461; Martin & Wilcox, 1965: 400 (cat.).
- sackeni** Hine, 1911: 166. Type-locality: U. S. A., southern Arizona. Distr. – U. S. A. (Arizona, Texas), Mexico (Michoacán, northern Sonora). Refs. – Bromley, 1935c: 1; Carrera & d'Andretta, 1950: 171; Hull, 1962 (2): 461; Martin & Wilcox, 1965: 400 (cat.). Type, OSU.
- truquii** Bellardi, 1861: 129 (29), pl. 2, fig. 6. Type-locality: "Mexico". Distr. – U. S. A. (Arizona), Mexico (Veracruz, Hidalgo, ?Nuevo León). Refs. – Williston, 1901: 322; Aldrich, 1905: 280 (cat.); Kertész, 1909: 237 (cat.); Hine, 1911a: 170; Hull, 1962 (2): 462; Martin & Wilcox, 1965: 400 (cat.); Martin & Papavero, 1970: 80 (cat.); Artigas & Papavero, 1997c: 105. Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol.). Type, TORO.
- venustus** Carrera & d'Andretta, 1950: 169, fig. 10. Type-locality: Mexico, San Luís Potosí, Valles El Bonito. Distr. – Mexico (San Luís Potosí). Refs. – Hull, 1962 (2): 462; Artigas & Papavero, 1997c: 105. Type, CMNH.

Myaptex-group

Genus *Atractocoma* Artigas, 1970

Atractocoma Artigas, 1970a: 259. Type-species, *Atractocoma nivosa* Artigas, 1970 (orig. des.).

nivosa Artigas, 1970a: 260, figs. 220–222, 488. Type-locality: Chile, Aysén, Chile Chico, laguna Buenos Aires. Distr. – Chile (Aysén). Ref. – Artigas & Papavero, 1995f: 57, figs. 32–41. Type, DZUC.

Genus **Martintella** Artigas, 1996

Martinella Artigas & Papavero, 1995f: 57 (preocc. Jousseau, 1887). Type-species, *Asilus lestes* Williston, 1901 (orig. des.).
Martintella Artigas, 1996: 75 (*nom. nov.* for *Martinella* Artigas & Papavero, 1995). Type-species, *Asilus lestes* Williston, 1901 (aut.).

lestes (Williston), 1901: 331 (*Asilus*). Type-locality: Mexico, Guerrero, Chilpancingo. Distr. – Mexico (Guerrero, Morelos). Refs. – Aldrich, 1905: 282 (cat.; *Asilus*); Kertész, 1909: 258 (cat.; *Asilus*); Hull, 1962 (2): 481 (*Lecania*); Martin, 1975c: 74, figs. 50, 65 (*Wilcoxius*); Artigas & Papavero, 1995f: 57, figs. 66-73 (*Martinella*); Artigas, 1996: 75 (n. comb.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.

Genus **Myaptex** Hull, 1962

Myaptex Hull, 1962 (2): 508. Type-species, *Myaptex hermanni* Hull, 1962 (orig. des.).

brachyptera (Philippi), 1965: 698 (*Asilus*). Type-locality: Chile, Colchagua. Neotype-locality: Chile, Malleco, Angol. Distr. – Chile (Arauco, Cautín, Concepción, Curicó, Linares, Maule, Malleco, Ñuble, Santiago, Talca, Valparaíso). Refs. – Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 252 (cat.; *Asilus*); Bromley, 1932: 281 (*Asilus*); Stuardo Ortiz, 1946: 84 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*), fig. 2445 [*Myaptex*]; Artigas, 1970a: 324, figs. 317-322, 434 (n. comb.), 1971: 43, figs. 109-110; Artigas & Papavero, 1995f: 57, figs. 12-17. Type, lost. NT DZUC (des. by Artigas, 1970a: 324).
hermanni Hull, 1962 (2): 509, figs. 195, 688, 1499, 2267, 2313, 2437, 2445. Type-locality: Chile, Concepción. Distr. – Chile (Biobío, Concepción, Curicó, Linares, Malleco, Maule, O'Higgins, Santiago, Valparaíso). Refs. – Artigas, 1970a: 327, figs. 312-316, 433, 1971: 43, figs. 107-108; Artigas & Papavero, 1995f: 57, figs. 18-20. Type, DEI.

Genus **Myaptexaria** Artigas & Papavero, 1995

Myaptexaria Artigas & Papavero, 1995f: 58. Type-species, *Myaptex vexillaria* Artigas, 1970 (orig. des.).

acuta (Artigas), 1980: 365, figs. 1-13 (*Myaptex*). Type-locality: Chile, Atacama, Castilla. Distr. – Chile (Atacama). Ref. – Artigas & Papavero, 1995f: 58 (n. comb.). Type, DZUC.
vexillaria (Artigas), 1970a: 329, figs. 323-328, 454 (*Myaptex*). Type-locality: Chile, Coquimbo, Vicuña. Distr. – Chile (Aconcagua, Coquimbo, Santiago, Valparaíso). Refs. – Artigas, 1971: 44, figs. 111-112 (*Myaptex*); Artigas & Papavero, 1995f: 58, figs. 21-31 (n. comb.). Type, DZUC.
virilis (Artigas), 1970: 331, figs. 329-333, 449 (*Myaptex*). Type-locality: Chile, Atacama, Zapata, Termas Socos. Distr. – Chile (Aconcagua, Atacama, Coquimbo). Refs. – Artigas, 1971: 44, figs. 105-106 (*Myaptex*); Artigas & Papavero, 1995f: 58 (n. comb.). Type, DZUC.

Genus **Rhadinosoma** Artigas, 1970

Rhadinosoma Artigas, 1970a: 346. Type-species, *Rhadinosoma calderense* Artigas, 1970 (orig. des.).

calderense Artigas, 1970a: 347, figs. 378-384, 469. Type-locality: Chile, Atacama, 40 km southeast of Caldera. Distr. – Chile (Atacama). Refs. – Artigas, 1971: 48, figs. 118-119; Artigas & Papavero, 1995f: 58, figs. 42-53. Type, DZUC.

Genus **Scarboroughia**, nom. nov.

Furcilla Martin, 1975c: 76 (preocc. Stokes, 1890). Type-species, *Furcilla dorothyae* Martin, 1975 (orig. des.).
Scarboroughia, nom. nov. for *Furcilla* Martin, 1975. Type-species, *Furcilla dorothyae* Martin, 1975 (aut.). [Dedicated to Dr. Aubrey G. Scarborough, Visiting Scholar of the University of Arizona, Tucson].

dorothyae (Martin), 1975c: 76, fig. 48 (*Furcilla*). Type-locality: Mexico, Sonora, 11 miles south of Navajoa, highway 15, km post 1766. Distr. – Mexico (Sonora). Ref. – Artigas & Papavero, 1995f: 57, figs. 54-65 (*Furcilla*). Type, CAS. N. comb.

petila (Martin), 1975c: 77, fig. 53 (*Furcilla*). Type-locality: Mexico, Sonora, 5 miles east of Navajoa. Distr. – Mexico (Sonora). Ref. – Artigas & Papavero, 1995f: 57 (*Furcilla*). Type, CAS. N. **comb.**

Genus **Wilcoxius** Martin, 1975

Wilcoxius Martin, 1975c: 71. Type-species, *Wilcoxius truncus* Martin, 1975 (orig. des.).

Ref. – Scarbrough & Perez-Gelabert, 2005 (rev.).

acutulus Martin, 1975c: 73, fig. 47. Type-locality: Nicaragua, Condega, 2025 ft. Distr. – Nicaragua, El Salvador. Refs. – Artigas & Papavero, 1995f: 58, figs. 9-11; Scarbrough & Perez-Gelabert, 2005: 49. Type, CAS.

bullatus (Bromley), 1929a: 288, fig. 10, pl. 1, fig. 14 (*Asilus*). Type-locality: Cuba, Santiago de las Vegas. Distr. – Cuba. Refs. – Hull, 1962 (2): 545 (*Asilus*); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.; *Pachychaeta*); Scarbrough & Perez-Gelabert, 2005: 42 (n. comb.). PT MCZ.

caputitis Scarbrough & Perez-Gelabert, 2005: 42, figs. 1-3. Type-locality: Honduras, Francisco Morazán, 25 km south of Talanga (Hacienda Havada). Distr. – Honduras. Type, FSCA.

crenus Martin, 1975c: 74, figs. 49, 67. Type-locality: Mexico, Chiapas, Naching. Distr. – Mexico (Chiapas). Refs. – Artigas & Papavero, 1995f: 58; Scarbrough & Perez-Gelabert, 2005: 49. Type, COR.

juventus Scarbrough & Perez-Gelabert, 2005: 44, figs. 4-7. Type-locality: Cuba, Isla de la Juventud, Nueva Gerona. Distr. – Cuba (Isla de la Juventud). Type, CMNH.

loewi (Bromley), 1929a: 289 (*Asilus*). Type-locality: “Cuba”. Distr. – Cuba. Refs. – Hull, 1962 (2): 545 (*Asilus*); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.; *Pachychaeta*); Scarbrough & Perez-Gelabert, 2005: 45 (n. comb.). Type, MCZ.

planus Scarbrough & Perez-Gelabert, 2005: 45, figs. 8-11. Type-locality: Dominican Republic, Azua, rodovía 229, 2 km road to San Juan, near bridge on Viajama river, 194 m. Distr. – Dominican Republic. Ref. – Scarbrough & Perez-Gelabert, 2006: 16. Type, USNM.

ramsdeni (Bromley), 1929a: 289 (*Asilus*). Type-locality: Cuba, Guantánamo, Tiguabos. Refs. – Hull, 1962 (2): 545 (*Asilus*); Alayo Dalmau & Garcia Ávila, 1983: 49 (cat.; *Pachychaeta*); Scarbrough & Perez-Gelabert, 2005: 46 (n. comb.). Type, USNM.

similis Scarbrough & Perez-Gelabert, 2005: 46, figs. 12-14. Type-locality: Guatemala, Zacapa, northwest of Teculután, road to Santerzo Marmal mines, 500 m. Distr. – Guatemala (Izabal, Zacapa). Type, FSCA.

truncus Martin, 1975c: 75, figs. 46, 66. Type-locality: Mexico, Veracruz, Veracruz. Distr. – Mexico (Veracruz, Oaxaca). Ref. – Artigas & Papavero, 1995f: 58, figs. 1-8. Type, CAS.

tumidus Scarbrough & Perez-Gelabert, 1995: 47, figs. 15-17. Type-locality: Dominican Republic, San Juan, road 237, 2 km road Belencio Ingenito, Presa de Sabaneta. Distr. – Dominican Republic (Azua, Pedernales, San Juan, Santo Domingo). Ref. – Scarbrough & Perez-Gelabert, 2006: 16. Type, USNM.

Ommatius-group

Ommatiini Hardy, 1927a: 384, 1927b: 388, 390. Type-genus, *Ommatius* Wiedemann, 1821.

Ommatiinae Ricardo, 1929: 120.

Ommatini Hull, 1962 (2): 433.

Genus **Ommatius** Wiedemann, 1821

Ommatius Wiedemann, 1821: 213. Type-species, *Asilus marginellus* Fabricius, 1781 (sub. des., Coquillett, 1910: 579). *Emphysomera* Schiner of authors, misident.

Ref. – Scarbrough, 2008: 2-4 (key to species-groups).

abana Curran, 1953: 2. Type-locality: Bahamas, South Bimini Island. Distr. – Bahamas. Refs. – Martin, 1957b: 7; Hull, 1962 (2): 435; Martin & Papavero, 1970: 58 (cat.); Scarbrough, 1985a: 1226 (in key). Type, AMNH.

achaetus Scarbrough, 1993: 732 (in key to males), 752, figs. 143-150. Type-locality: Mexico, Oaxaca, 7 miles north of Juchatengo, 3700 ft. Distr. – Mexico (Oaxaca). Ref. – Vieira, Castro & Bravo, 2005: 20 (in key). Type, ?

acus Scarbrough, 1990: 70 (in key), 95, figs. 1H, 2H, 17A-F, 18A-C. Type-locality: Mexico, 29 miles south of Chilpancingo,

- highway 90. Distr. – Mexico (Chiapas, Jalisco, Morelos, Nayarit, Oaxaca). Type, FSCA.
- alexanderi** Farr, 1965: 24, figs. 16, 20. Type-locality: Jamaica, St. Andrew, Ferry. Distr. – Jamaica. Refs. – Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1993: 732 (in key to males), 733 (in key to females), 734, figs. 1-11; Vieira, Castro & Bravo, 2005: 21 (in key). Type, USNM.
- ampliatus** Scarbrough, 2002: 138 (in key), 140, figs. 1, 18-26. Type-locality: Colombia, Cauca, 20 km west of Silvia. Distr. – Colombia (Cauca, Medellín). Type, CNC.
- amula** Curran, 1928a: 1 (in key), 4. Type-locality: Mexico, Guerrero, Amula, 6000 ft. Distr. – Mexico (Chiapas, Chihuahua, Guerrero, Jalisco, Michoacán, Morelos, Nayarit, Oaxaca, Sinaloa, Veracruz), Belize, Guatemala, El Salvador. Refs. – Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1993: 732 (in key to males), 733 (in key to females), 735, figs. 12-2; Vieira, Castro & Bravo, 2005: 20 (in key). Type, AMNH.
- aemula* Hull, 1962 (2): 435, error.
- angulosus** Scarbrough, 2002: 139 (in key), 145, figs. 2, 27-35. Type-locality: Colombia, Valle, 6 miles west of Cali. Distr. – Colombia (Valle), Ecuador. Type, CAS.
- angustatus** Scarbrough, 2002: 140 (in key), 149, figs. 3, 36-44. Type-locality: Costa Rica, Puntarenas, F. Las Cruces, 6 km south of San Vito, 1200-1400 m. Distr. – Guatemala, Costa Rica, Panama. Type, CAS.
- apicalis** (Bellardi), 1861: 157 (57) (*Asilus*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Osten Sacken, 1878: 82, 235 (cat.; *Asilus*); Aldrich, 1905: 282 (cat.; *Asilus*); Kertész, 1909: 251 (cat.; *Asilus*); Hull, 1962 (2): 435, 545 (*Asilus*); Martin & Papavero, 1970: 59 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- aridus** Scarbrough, 2002: 201, figs. 167, 173-181. Type-locality: Ecuador, Galápagos Islands, Floreana Island, 3 km east of Black Beach, 120 m. Distr. – Ecuador (Galápagos Islands). Type, CNC.
- atrosus** Scarbrough, 1997a: 11, figs. 1-7. Type-locality: Dominican Republic, Pedernales, 26 km north of Cabo Rojo. Distr. – Dominican Republic (Pedernales). Ref. – Scarbrough & Perez-Gelabert, 2003: 43, fig. 50. Type, CMNH.
- ayalai** Scarbrough, 2002: 138 (in key), 154, figs. 4, 45-53. Type-locality: Venezuela, Táchira, Delicias, 1800 m. Distr. – Venezuela. Type, IEAUCV.
- bastardoanus** Scarbrough in Scarbrough & Perez-Gelabert, 2003: 43, figs. 1-8, 51. Type-locality: Dominican Republic, La Vega, Arroyazo, Reserva Científica Ébano Verde. Distr. – Dominican Republic (La Vega, Santiago), Hispaniola. Ref. – Scarbrough & Perez-Gelabert, 2006: 46. Type, USNM.
- beameri** Wilcox, 1936a: 174 (in key), 3, pl. 10. Type-locality: U. S. A., Arizona, Patagonia. Distr. – U. S. A. (New Mexico, Arizona), Mexico (Nayarit, Jalisco). Refs. – Hull, 1962 (2): 435; Martin & Wilcox, 1965: 398; Scarbrough, 1990: 70 (in key), 79, figs. 1C, 2C, 7A-F, 8A-C. Type, KANSAS.
- bicolor** (Bigot), 1875: 244 (*Emphysomera*). Type-locality: “Mexico”. Distr. – Mexico. Ref. – Martin & Papavero, 1970: 59 (cat.). Type, OXF.
- bipartitus** Scarbrough, 1985a: 1226 (in key), 1232, figs. 10-14. Type-locality: Cuba, Loma del Gato, Cobre Range, 3000 ft. Distr. – Cuba. Type, MCZ.
- bullatus** Scarbrough, 2002: 138 (in key), 157, figs. 5, 54-62. Type-locality: Venezuela, Notigrito, Parque Nacional El Ávila. Distr. – Venezuela. Type, IZAUCV.
- cinnamomeus** Scarbrough, 1984b: 132 (in key), 147, fig. 14. Type-locality: Haiti, Nordest District, La Hotte, 200-400 ft. Distr. – Haiti (Nordest District), Dominican Republic (Independencia, Pedernales). Refs. – Scarbrough, 1997a: 11, figs. 8-10; Scarbrough & Perez-Gelabert, 2003: 44, fig. 49, 2006: 46, figs. 40-44. NT CMNH [des. by Scarbrough, 1997a: 11].
- cinthiae** Vieira, Castro & Bravo, 2004: 2, figs. 1-6. Type-locality: Brazil, Bahia, Santa Terezinha, Serra da Jibóia. Distr. – Brazil (Bahia). Ref. – Vieira, Castro & Bravo, 2005: 20 (in key). Type, UEFS.
- complanatus** Scarbrough, 1993: 732 (in key to males), 734 (in key to females), 737, figs. 23-33. Type-locality: Brazil, São Paulo, Salesópolis (Estação Biológica de Boracéia), 850 m. Distr. – Brazil (Rio de Janeiro, São Paulo). Ref. – Vieira, Castro & Bravo, 2005: 20 (in key). Type, MZUSP.
- constrictus** Scarbrough, 2002: 137 (in key), 160, figs. 63-65. Type-locality: Guyana, Orenoque and New, River Head. Distr. – Guyana. Type, BMNH.
- conus** Scarbrough, 2002: 204, figs. 168, 182-190. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo (Muri). Distr. – Brazil (Rio de Janeiro). Type, MZUSP.
- costatus** Rondani, 1850a: 188. Type-locality: Brazil, São Paulo, ilha de São Sebastião. Distr. – Trinidad and South America, except Chile and Uruguay. Refs. – Osten Sacken, 1891: 421; Kertész, 1909: 307 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1993: 732 (in key to males), 733 (in key to females), 738, figs. 34-46; Vieira, Castro & Bravo, 2005: 21 (in key). Lectotype TORO [des. by Scarbrough, 1993: 739].
- barbiellini* Curran, 1934a: 18. Type-locality: Brazil, São Paulo. Distr. – Brazil (São Paulo). Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1993: 738 (syn.). Type, AMNH.
- cubanus** Scarbrough, 1985a: 1227 (in key), 1229, figs. 6-7 (as *bubansus*, error). Type-locality: Cuba, Buenos Aires. Distr. – Cuba. Type, MCZ.
- dentatus** Scarbrough, 1993: 732 (in key to males), 733 (in key to females), 739, figs. 47-57. Type-locality: Brazil, Amazonas, Manaus. Distr. – French Guiana, Brazil (Amazonas, Pará, Amapá). Ref. – Vieira, Castro & Bravo, 2005: 21 (in key). Type, INPA.
- depressus** Scarbrough, 2002: 137 (in key), 161, figs. 66-68. Type-locality: Ecuador, San Francisco, río Pastaza, 1300 m. Distr. – Ecuador. Type, AMNH.

- destitutus** Scarbrough, 2002: 140 (in key), 163, figs. 6, 69-77. Type-locality: Peru, Cajamarca. Distr. – Peru. Type, USNM.
- didymus** Scarbrough, 1993: 732 (in key to males), 741, figs. 58-65. Type-locality: Brazil, Amazonas, Manaus. Distr. – Brazil (Amazonas). Ref. – Vieira, Castro & Bravo, 2005: 21 (in key). Type, INPA.
- dignus** Scarbrough, 2000: 917 (*nom. nov.* for *dimidiatus* Scarbrough, 1985). Type-locality: Dominica, Clarke Hall, Cocoa Trail. Distr. – Dominica, Lucia and Martinique.
- dimidiatus* Scarbrough, 1985b: 647, figs. 9-14 (preocc. Macquart, 1850). Type-locality: Dominica, Clarke Hall, Cocoa Trail. Ref. – Scarbrough & Knutson, 1989: 524. Type, USNM.
- dolabrigeriformis** Scarbrough, 2002: 139 (in key), 166, figs. 7, 78-83. Type-locality: Colombia, San Sebastián de Rabago, Sierra Nevada de Santa María, 2000 m. Distr. – Colombia. Type, FMNH.
- elusivus** Scarbrough *in* Scarbrough & Perez-Gelabert, 2003: 46, figs. 9-16, 49. Type-locality: Dominican Republic, Santiago, Casetas Parque Diferencia, Parque Nacional Armando Bermúdez. Distr. – Dominican Republic (La Vega, Santiago), Hispaniola. Ref. – Scarbrough & Perez-Gelabert, 2006: 49, 71 (in key), figs. 45, 52. Type, USNM.
- emarginatus** Scarbrough, 1985b: 646, figs. 7-8. Type-locality: St. Vincent Island. Distr. – St. Vincent Island. Type, BMNH.
- erythropus** Schiner, 1867: 411. Type-locality: “South America”. Refs. – Williston, 1891: 90 (cat.); Kertész, 1909: 308 (cat.); Engel, 1930: 464, fig. 5 (*Ommatinus*); Carrera & Vulcano, 1961: 77 (prey); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.). Type, NMW.
- exilis** Curran, 1928a: 2 (in key), 5. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Distr. – Peru (Loreto), Brazil (Rondônia, Mato Grosso, Tocantins, Rio de Janeiro, São Paulo), Argentina (Salta). Refs. – Bromley, 1946: 112 (cat.); Carrera, 1960a: 167; Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Lamas M., 1972: 312 (cat.); Scarbrough, 2002: 207, figs. 169, 191-199. Type, AMNH.
- falcatus** Scarbrough, 1984a: 625, fig. 6. Type-locality: Puerto Rico, Maricao. Distr. – Puerto Rico. Type, USNM.
- fernandezi** Scarbrough, 2002: 140 (in key), 168, figs. 8, 84-92. Type-locality: Venezuela, Raragua, La Trilla. Distr. – Venezuela. Type, IZAUCV.
- †**fimbrillus** Scarbrough, 2000: 917 (*nom. nov.* for *fimbriatus* Scarbrough & Poinar, 1992). Type-locality: Dominican Republic (Oligocene/Miocene).
- †**fimbriatus** Scarbrough & Poinar Jr., 1992: 13 (preocc. G. H. Hardy, 1949). Type-locality: Dominican Republic (Oligocene/Miocene). Type, GPAC.
- flavescens** Scarbrough *in* Scarbrough & Perez-Gelabert, 2003: 48, figs. 17-21, 49. Type-locality: Dominican Republic, Pedernales, 23.5 km north of Cabo Rojo. Distr. – Dominican Republic (Pedernales). Type, CMNH. Ref. – Scarbrough & Perez-Gelabert, 2006: 72 (in key).
- flexus** Scarbrough, 2002: 138, 139 (in key), 171, figs. 9, 93-101. Type-locality: Venezuela, Táchira, La Parada, San Cristóbal, 850 m. Distr. – Venezuela. Type, IZAUCV.
- fuscipennis** Bellardi, 1862: 220 (23). Type-locality: Mexico, Veracruz, Tuxpango near Orizaba. Distr. – Mexico (Veracruz). Refs. – Aldrich, 1905: 273 (cat.); Kertész, 1909: 308 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Papavero & Ibáñez Bernal, 2001: 110 (hist. Mexican dipteron). Type, TORO.
- geminus** Scarbrough & Perez-Gelabert, 2006: 51, 71 (in key), figs. 46, 53, 59-66. Type-locality: Dominican Republic, La Vega, Parque Nacional Armando Bermúdez, km 1.3 along trail west of La Ciénaga, 900-1100 m. Distr. – Dominican Republic (La Vega). Type, USNM.
- gladiatus** Scarbrough, 2002: 138 (in key), 174, figs. 10, 102-111. Type-locality: Mexico, Veracruz, Jalapa. Distr. – Mexico Nuevo León, Veracruz, Hidalgo, Oaxaca, Chiapas), Guatemala, Costa Rica. Type, USNM.
- gwenae** Scarbrough, 1984b: 132 (in key), 139, figs. 9-10. Type-locality: Dominican Republic, Constanza. Lectotype-locality: Dominican Republic, La Toma, north of San Cristóbal. Distr. – Dominican Republic (Distrito Nacional, Hato Mayor, Independencia, Isla Beata La Altagracia, La Vega, M. T. Sánchez, Pedernales, Peravia, Sánchez Ramírez, San Cristóbal, San Pedro de Macorís, Santiago). Refs. – Scarbrough, 1997a: 12, figs. 11-20; Scarbrough & Perez-Gelabert, 2003: 48, fig. 49, 2006: 46. Lectotype USNM [des. by Scarbrough, 1997a: 12].
- haitiensis** Scarbrough, 1984b: 132, figs. 1-3. Type-locality: Haiti, La Viste and Vie La Salle Range. Distr. – Haiti (La Visite and Vie La Selle Range), Dominican Republic (Bahoruco, Pedernales, Independencia). Refs. – Scarbrough, 1997a: 12, figs. 21-29; Scarbrough & Perez-Gelabert, 2003: 49, fig. 50. Type, MCZ.
- hanebrinki** Scarbrough & Rutkauskas, 1983: 144, figs. 1-9. Type-locality: Bahamas Islands, San Salvador Island, College Center of the Finger Lakes, Bahamian Campus. Distr. – Bahamas Islands. Ref. – Scarbrough, 1985a: 1227 (in key). Type, USNM.
- hierroi** Scarbrough *in* Scarbrough & Perez-Gelabert, 2003: 49, figs. 22-29, 49. Type-locality: Dominican Republic, Duarte, 8 miles north of San Francisco de Macoris, Loma Quita Espuela, 800 m. Distr. – Dominican Republic (Duarte). Type, AMNH.
- hispaniolae** Scarbrough, 1984b: 132 (in key), 137, figs. 7-8. Type-locality: Dominican Republic, Barahona. Distr. – Haiti (Ouest District), Dominican Republic (Azua, Barahona, Independencia, Isla Beata, Montecristi, M. T. Sánchez, Peravia, Pedernales, San Cristóbal, San Juan, Santiago). Refs. – Scarbrough, 1997a: 13, figs. 30-30; Scarbrough & Perez-Gelabert, 2003: 52, fig. 50, 2006: 46. Type, MCZ.
- hispidus** Scarbrough, 1985a: 1226 (in key), 1235, figs. 15-20. Type-locality: Cuba, Santiago de las Vegas. Distr. – Cuba. Type, USNM.
- holosericeus** Schiner, 1867: 411. Type-locality: “Brazil [Rio de Janeiro]”. Distr. – Brazil (Espírito Santo, Rio de Janeiro, São Paulo, Santa Catarina). Refs. – Williston, 1891: 90 (cat.); Kertész, 1909: 308 (cat.); Bromley, 1946: 112 (cat.); Hull, 1962

- (2): 224, 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 2002: 210, figs. 170, 200-208. Lectotype NMW [des. by Scarbrough, 2002: 214].
- humatus** Scarbrough, 1993: 741, figs. 66-76. Type-locality: Costa Rica, Heredia, F. La Selva, 3 km south of Puerto Viejo. Distr. – Mexico, Mesoamerica, Ecuador. Ref. – Vieira, Castro & Bravo, 2005: 20 (in key). Type, USNM.
- incurvatus** Scarbrough, 1993: 731 (in key to males), 733 (in key to females), 744, figs. 77-87. Type-locality: Guatemala, 31 km northeast of Guatemala, 915 m. Distr. – Mexico to Costa Rica. Refs. – Vieira, Castro & Bravo, 2005: 21 (in key); Scarbrough, 2007: 470 (in key). Type, EMF.
- inflatus** Scarbrough in Scarbrough & Perez-Gelabert, 2003: 50, figs. 30-37, 49. Type-locality: Dominican Republic, Montecristi, 9 km north of Montecristi. Distr. – Dominican Republic (Barahona, Montecristi, Peravia, San Juan, Santiago). Ref. – Scarbrough & Perez-Gelabert, 2006: 46, 71 (in key), figs. 48, 54. Type, FSCA.
- infuscatus** Scarbrough, 1990: 70 (in key), 86, figs. 1E, 2E, 11A-F, 12A-C. Type-locality: Mexico, Michoacán, 9 miles west of Morelia. Distr. – Mexico (Jalisco, Michoacán, Morelos). Type, FSCA.
- integerrimus** Scarbrough, 1990: 70 (in key), 82, figs. 1D, 2D, 9A-F, 10A-C. Type-locality: Mexico, Morelos, 2 mi. South of Tepoztlán, 5300 ft. Distr. – Mexico (Guerrero, Michoacán, Morelos, Nayarit). Type, FSCA.
- jamaicensis** Farr, 1965: 23, figs. 15, 19. Type-locality: Jamaica, St. Thomas, Morant Bay road, 14.5 miles east of Kingston. Distr. – Jamaica. Ref. – Martin & Papavero, 1970: 59 (cat.). Type, USNM.
- laticerus** Scarbrough & Perez-Gelabert, 2006: 58, 72 (in key), figs. 49, 55, 67-74. Type-locality: Dominican Republic, Peravia, southern slopes of Cerro Gordo, Bani, on way to Salinas. Distr. – Dominican Republic (Azua, Barahona, Peravia, Santiago). Type, USNM.
- lineolatus** Scarbrough, 1988b: 91, figs. 2a-d. Type-locality: Cuba, Herradura. Distr. – Bahamas, Cuba. Ref. – Scarbrough, 2000: 917. Type, USNM.
- lucidatus** Scarbrough, 1997a: 13, figs. 40-49. Type-locality: Dominican Republic, Pedernales, north of Cabo Rojo. Distr. – Dominican Republic (Azua, Bahoruco, Independencia, Montecristi, Pedernales, Peravia, Santiago). Ref. – Scarbrough & Perez-Gelabert, 2003: 53, 72 (in key), fig. 51, 2006: 62, figs. 50, 56. Type, CMNH.
- lunatus** Scarbrough, 2002: 137, 138 (in key), 178, figs. 11, 112-121. Type-locality: Costa Rica, Heredia, F. La Solva, 3 km south of Puerto Viejo. Distr. – Costa Rica. Type, USNM.
- maculosus** Scarbrough & Perez-Gelabert, 2006: 62, 72 (in key), figs. 47, 57, 75-82. Type-locality: Dominican Republic, Pedernales, road-125, 2 km on trail to Carlitos, Parque Nacional Jaragua, 124 m. Distr. – Dominican Republic (Pedernales). Type, USNM.
- marginellus** (Fabricius), 1781: 474 (*Asilus*). Type-locality: Virgin Islands, St. Croix. Distr. – Virgin Islands, Cuba, St. Thomas, Venezuela, Guyana, Galapagos Islands, Brazil (Pará, Espírito Santo, São Paulo), Paraguay. Refs. – Fabricius, 1787: 360 (*Asilus*); Olivier, 1789: 267 (*Asilus*); Gmelin, 1792: 2898 (*Asilus*); Fabricius, 1794: 384 (*Asilus*), 1805: 170 (*Dasyopogon*); Wiedemann, 1821: 213, 1828: 421, pl. 6, fig. 5; Macquart, 1834: 314, 1838: 134, pl. 11, fig. 4; Walker, 1849: 474, 1855: 755; Schiner, 1866: 682; Osten Sacken, 1878: 83, 235 (cat.); Roeder, 1885: 339; Williston, 1891: 90 (cat.), 1896a: 304; Aldrich, 1905: 273 (cat.); Kertész, 1909: 308 (cat.); Coquillett, 1910: 579; Gowdey, 1926: 77 (in error); Curran, 1928a: 1 (in key). 1928b: 23; Bromley, 1929a: 286; Carrera, 1947a: 48; Wolcott, 1948: 453; Carrera, 1958a: 152, 1960a: 167; Carrera & Vulcano, 1961: 77 (preys); Hull, 1962 (2): 435; Carrera & Machado-Allison, 1963: 263; Farr, 1965: 25 (not in Jamaica); Linsley & Usinger, 1966: 166 (Gallapagos Islands); Martin & Papavero, 1970: 59 (cat.); Alayo Dalmau & García Ávila, 1983: 48 (cat.); Scarbrough, 1984a: 620, figs. 1-4, 1985a: 1237, 1985b: 652. Type, KIEL (without head and abdomen) [cf. Zimsen, 1964: 470].
marginalis Bromley, 1946: 112 (cat.), error.
- mariae** Scarbrough, 2000: 913, figs. 1-9. Type-locality: Bahamas, Eleuthera, Rainbow Bay. Distr. – Bahamas. Type, FSCA.
- megacephalus** (Bellardi), 1861: 158 (58), pl. 2, fig. 14 (*Asilus*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Aldrich, 1905: 282 (cat.; *Asilus*); Kertész, 1909: 260 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 59 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- membranosus** Scarbrough, 1985a: 1227 (in key), 1231, figs. 8-9. Type-locality: Bahamas, Rum Cay. Distr. – Bahamas. Type, MCZ.
- monensis** Scarbrough, 1984a: 623, fig. 5. Type-locality: Puerto Rico, Mona Island, Camp Capuasi. Distr. – Puerto Rico (Mona Island). Type, USNM.
- narrius** Scarbrough, 2002: 214, figs. 171, 209-217. Type-locality: Brazil, São Paulo, São Paulo (Horto Florestal). Distr. – Brazil (Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo). Type, MZUSP.
- nebulosus** Scarbrough, 2008: 4, figs. 1, 6-14. Type-locality: Venezuela, Amazonas, Cerro de la Neblina base camp, 140 m, 0°05'N, 6610'W. Distr. – Venezuela (Amazonas). Type, USNM.
- neotropicus** Curran, 1928a: 1 (in key), 3. Type-locality: “Guyana”. Distr. – Guyana, Brazil (Minas Gerais). Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 2008: 6, figs. 2, 15-23. Type, AMNH.
- nigellus** Scarbrough, 1984b: 132 (in key), 144, fig. 12. Type-locality: Haiti, Furcy. Distr. – Haiti (Ouest District), Dominican Republic (Barahona, Pedernales). Refs. – Scarbrough, 1997a: 15, figs. 50-59; Scarbrough & Perez-Gelabert, 2003: 54, fig. 51, 2006: 46. Type, AMNH.
- norma** Curran, 1928a: 1 (in key), 2. Type-locality: “Guyana”. Distr. – Venezuela (Amazonas, Bolívar), Guyana, Surinam, French Guiana, Brazil (Amazonas, Pará, Rondônia, Mato Grosso do Sul, Minas Gerais, São Paulo), Paraguay (Cordillera). Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 2008: 7, figs. 3, 24-32 (as *normus*, error). Type, AMNH.

- orenoquensis** Bigot, 1876: lxxxv. Type-locality: "Guyana". Distr. – Costa Rica and St. Vincent Islands southward, South America except Uruguay and Chile. Refs. – Osten Sacken, 1891: 421; Kertész, 1909: 310 (cat.); Curran, 1928a: 1 (in key); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1993: 732 (in key to males), 733 (in key to females), 746, figs. 88-98; Vieira, Castro & Bravo, 2005: 20 (in key). Lectotype OXF [des. by Scarbrough, 1993: 746].
- infractus** Scarbrough, 1985b: 643. Type-locality: St. Vincent Island. Ref. – Scarbrough, 1993: 746 (syn.). Type, USNM.
- oreophilus** Farr, 1965: 20, figs. 14, 18, 22. Type-locality: Jamaica, St. Andrew, Hardwar Gap. Distr. – Jamaica. Refs. – Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1993: 732 (in key to males), 733 (in key to females), 747, figs. 99-109; Vieira, Castro & Bravo, 2005: 20 (in key). Type, USNM.
- ovatus** Scarbrough, 2002: 139 (in key), 181, figs. 122-124. Type-locality: Peru, San Martín. Distr. – Peru. Type, USNM.
- parvulus** Schaeffer, 1916: 69. Type-locality: U. S. A., Arizona, Huachuca Mts.. Distr. – U. S. A. (Arizona). Refs. – Pritchard, 1935: 8; Hull, 1962 (2): 435; Martin & Wilcox, 1965: 398 (cat.).
- parvus** Bigot, 1875: 247. Type-locality: "Mexico". Distr. – Mexico (Chiapas, Colima, Guerrero, Hidalgo, Jalisco, Michoacán, Morelos, Nayarit, Oaxaca, Sinaloa, Veracruz), Guatemala, Honduras, El Salvador, Nicaragua, Costa Rica. Refs. – Williston, 1901: 332; Aldrich, 1905: 273 (cat.); Kertész, 1909: 310 (cat.); Curran, 1928a: 2 (in key); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1990: 71, figs. 1A, 2A, 3A-F, 4A-C. Lectotype OXF [des. by Scarbrough, 1990: 71].
- pumilus* Macquart of Bellardi, 1861: 59, misident.
- perangustimus** Scarbrough, 1990: 70 (in key), 94, figs. 1G, 2G, 15A-F, 16A-C. Type-locality: Mexico, Oaxaca, 27 miles south of Juchatengo, 4200 ft. Distr. – Mexico (Chiapas, Oaxaca, Sinaloa). Type, EMF.
- peregrinus** Osten Sacken, 1887: 210. Type-locality: Panama, Bugaba, 800-1500 ft. Distr. – Panama. Refs. – Williston, 1901: 331; Aldrich, 1909: 311 (cat.); Kertész, 1909: 311 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.). Type, BMNH.
- piliferous** [sic] Scarbrough, 1985a: 1226 (in key), 1227, figs. 1-5. Type-locality: "Cuba". Distr. – Cuba. Ref. – Scarbrough, 1993: 732 (in key to males), 733 (in key to females), 748, figs. 110-120; Vieira, Castro & Bravo, 2005: 21 (in key). Type, MCZ.
- pilosulus** (Bigot), 1875: 243 (*Emphysomera*). Type-locality: "Mexico". Distr. – Mexico. Refs. – Williston, 1901: 332; Aldrich, 1905: 273 (cat.); Kertész, 1909: 311 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.). Type, OXF.
- pilosus* Curran, 1928a: 1 (in key), error.
- praelongus** Scarbrough & Perez-Gelabert, 2006: 66, 71 (in key), figs. 51, 58, 83-90. Type-locality: Dominican Republic, Pedernales, road-266, Las Abejas, Parque Nacional Sierra de Bahoruco, 1310 m. Distr. – Dominican Republic (Pedernales). Type, USNM.
- praestigiatus** Scarbrough, 1990: 71 (in key), 76, figs. 1B, 2B, 5A-F, 6A-C. Type-locality: Venezuela, Miranda, Los Ocumitos. Distr. – Venezuela. Type, CAS.
- pretiosus** Banks, 1911: 129. Type-locality: U. S. A., Arizona. Distr. – U. S. A. (Arizona), Mexico. Refs. – Curran, 1928: 1 (in key); Hull, 1962 (2): 435; Martin & Wilcox, 1965: 398 (cat.).
- prolongatus** Scarbrough, 1985b: 653, figs. 18-19. Type-locality:Montserrat Island, Plymouth. Distr. – Montserrat Island. Type, USNM.
- pulcher** (Engel), 1885: 1885: 146 (*Emphysomera*). Type-locality: Brazil, Minas Gerais, São João del Rei [Sello]. Distr. – Brazil (Alagoas, Mato Grosso do Sul, D. F., Goiás, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Santa Catarina), Paraguay (Amambay, Ybycui). Refs. – Curran, 1928a: 1 (in key; as *pulchra*, error); Engel, 1930: 465 (*Emphysomera*); Bromley, 1946: 112 (cat.); [*Ommatius* (*Emphysomera*)]; Hull, 1962 u(2): 437 (*Emphysomera*); Martin & Papavero, 1970: 60 (cat.); Scarbrough, 2008: 8, figs. 4, 33-41.
- pulverius** Scarbrough, 1997a: 16, figs. 60-66. Type-locality: Dominican Republic, Pedernales, Cabo Rojo. Distr. – Dominican Republic (Pedernales, Isla Beata). Ref. – Scarbrough & Perez-Gelabert, 2003: 54, fig. 50. Type, USNM.
- pumilus** Macquart, 1847: 58 (1847: 42), pl. 1, fig. 10. Type-locality: "Mexico". Distr. – Mexico (Chiapas, Guerrero, Oaxaca, Puebla, Sinaloa, Veracruz). Refs. – Walker, 1855: 755; Aldrich, 1905: 274 (cat.); Kertész, 1909: 311 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 1990: 70 (in key), 89, figs. 1F, 2F, 13A-F, 14A-C. Lectotype OXF [des. by Scarbrough, 1990: 92].
- pumillus* Papavero & Ibáñez Bernal, 2001: 86 (hist. Mexican dipterol.), error.
- quadratus** Scarbrough, 2002: 138 (in key), 182, figs. 125-127. Type-locality: Peru, Huaoapiatana, río Tarma. Distr. – Peru. Type, CU.
- riali** Vieira, Castro & Bravo, 2005: 21 (in key and text), figs. 1-7. Type-locality: Brazil, Bahia, Cachoeira. Distr. – Brazil (Bahia). Type, UEFS.
- ruficauda** Curran, 1928a: 2 (in key), 5. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Distr. – Brazil (Mato Grosso). Refs. – Bromley, 1946: 112 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 59 (cat.); Scarbrough, 2008: 11, fig. 5 (as *ruficaudus*, error) TP AMNH.
- rufipes** Macquart, 1846: 216 (1846: 88) (preocc. Macquart, 1838). Type-locality: "Colombia". Refs. – Walker, 1855: 756; Williston, 1891: 90 (cat.); Kertész, 1909: 311 (cat.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 60 (cat.). Type, lost.
- russelli** Scarbrough, 1984b: 132 (in key), 142 fig. 11. Type-locality: Dominican Republic, La Palma, 12 km east of El Río. Distr. – Dominican Republic (Azua, Independencia, La Vega, San Cristóbal). Refs. – Scarbrough, 1997a: 16; Scarbrough & Perez-Gelabert, 2003: 55, fig. 50. Type, USNM.
- saccas** Walker, 1849: 474. Type-locality: "Jamaica". Distr. – St. Croix, Jamaica. Refs. – Johnson, 1894: 274, 1919: 430; Hull, 1962 (2): 435 (as syn. of *marginellus* Fabricius, 1781); Farr, 1965: 18, figs. 13, 17, 21; Martin & Papavero, 1970: 60

- (cat.). Type, BMNH.
- schineri** Martin, 1965b: 130 (*nom. nov.* for *apicalis* Schiner, 1868). Type-locality: "South America". Ref. – Martin & Papavero, 1970: 60 (cat.).
apicalis Schiner, 1868: 194 (preocc. Bellardi, 1861). Type-locality: "South America". Refs. – Williston, 1891: 90 (cat.); Kertész, 1909: 306 (cat.); Curran, 1928a: 1 (in key). Type, NMW.
- scopifer** Schiner, 1868: 194. Type-locality: "Colombia". Distr. – Colombia. Refs. – Williston, 1891: 90 (cat.); Kertész, 1909: 311 (cat.); Carrera & d'Andretta, 1950: 181 (misident.); Hull, 1962 (2): 435; Martin & Papavero, 1970: 60 (cat.). Type, NMW.
- serrajiboensis** Vieira, Castro & Bravo, 2004: 3, figs. 7-13. Type-locality: Brazil, Bahia, Santa Terezinha, Serra da Jibóia. Distr. – Brazil (Bahia). Ref. – Vieira, Castro & Bravo, 2005: 20 (in key). Type, UEFS.
- setiferous** [sic] Scarbrough, 1988b: 90, figs. 1a-g. Type-locality: Bahama Islands, Mayaguana Island, Abraham Bay. Distr. – Bahama Islands (Mayaguana Island). Type, USNM.
- simulans** Scarbrough, 2002: 217, figs. 172, 218-226. Type-locality: Brazil, Goiás, Corumbá de Goiás (Fazenda Monjolinho). Distr. – Brazil (Goiás, Minas Gerais, São Paulo). Type, MZUSP.
- spatulatus** Curran, 1928a: 2 (in key), 4. Type-locality: unknown [?Rio de Janeiro]. Distr. – Brazil (Amazonas, Goiás, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Paraná), Argentina (Jujuy). Refs. – Carrera, 1945a: 165 (preys); Bromley, 1946: 112 (cat.); Carrera, 1960a: 167; Hull, 1962 (2): 435; Martin & Papavero, 1970: 60 (cat.); Scarbrough, 2007: 470. Type, AMNH.
- spinosus** Scarbrough, 1993: 732 (in key to males), 734 (in key to females), 749, figs. 121-131. Type-locality: Peru, Huánuco, San Domingo. Distr. – Ecuador, Peru, Brazil (Amazonas, Pará, Maranhão, Goiás, Rio de Janeiro), Bolivia. Ref. – Vieira, Castro & Bravo, 2005: 21 (in key). Type, USNM.
- stramineus** Scarbrough, 1984b: 132 (in key), 145, fig. 13. Type-locality: Haiti, Mt. Bouretta, 5000 ft. Distr. – Haiti, Dominican Republic (La Vega, Peravia). Refs. – Scarbrough, 1997a: 17, figs. 67-76; Scarbrough & Perez-Gelabert, 2003: 55, fig. 50. Type, USNM.
- †**subtus** Scarbrough & Poinar Jr., 1992: 13. Type-locality: Dominican Republic (Oligocene/Miocene).
- tandapiensis** Scarbrough, 2002: 184, figs. 12, 128-136. Type-locality: Ecuador, Tandapi, 40 km southwest of Quito, 1300-1500 m. Distr. – Ecuador. Type, CNC.
tanpadiensis Scarbrough, 2002: 139 (in key), error.
- tepui** Scarbrough, 2008: 11, figs. 42-48. Type-locality: Venezuela, Amazonas, Puerto Ayacucho, 40 km south of Tibogán, Canocoromoto. Distr. – Venezuela (Amazonas). Type, USNM.
- tractus** Scarbrough, 2007: 466, figs. 1-10, p. 470 (in key). Type-locality: Brazil, São Paulo, São Paulo (Santo Amaro). Distr. – Brazil (São Paulo). Type, MZUSP.
- triangularis** Scarbrough, 2002: 140 (in key), 187, figs. 13, 137-145. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. – Venezuela. Type, USNM.
- tropidus** Scarbrough, 2002: 139 (in key), 190, figs. 146-148. Type-locality: Bolivia, Chapare, Cristal. Distr. – Bolivia. Type, IZAUCV.
- tucumanensis** Scarbrough, 2002: 140 (in key), 192, figs. 14, 149-157. Type-locality: Argentina, Trucumán, Tafi Viejo, Quebrada Cainza. Distr. – Bolivia, Argentina (Tucumán). Type, MZUSP.
- uncatus** Scarbrough, 1993: 731 (in key to males), 733 (in key to females), 750, figs. 132-142. Type-locality: Peru, Huánuco, San Domingo. Distr. – Colombia, Peru, Brazil Pará). Refs. – Vieira, Castro & Bravo, 2005: 20 (in key); Scarbrough, 2007: 469 (in key). Type, USNM.
- unguiculatus** Scarbrough, 2002: 140 (in key), 195, figs. 15, 158-166. Type-locality: Venezuela, Distrito Federal, El Junquito. Distr. – Venezuela. Type, IZAUCV.
- villosus** Scarbrough, 1985b: 650, figs. 15-17. Type-locality: Dominica, Clarke Hall. Distr. – Dominica. Ref. – Scarbrough & Knutson, 1989: 524. Type, USNM.
- vitreus** Bigot, 1875: 246. Type-locality: "Haiti". Distr. – Hispabiola, Puerto Rico (Mona Island), Haiti (Ouest District), Dominican Republic (Pedernales, La Vega). Refs. – Kertész, 1909: 310 (cat.; as syn. of *marginellus* Fabricius, 1791); Hull, 1962 (2): 435 (as syn. of *marginellus* Fabricius, 1791); Martin & Papavero, 1970: 60 (cat.); Scarbrough, 1984a: 627, 1984b: 132 (in key), 135, figs. 4-6, 1997a: 17, figs. 77-86; Scarbrough & Perez-Gelabert, 2003: 55, fig. 51. Lectotype OXF [des. by Scarbrough, 1984: 135].
- vivus** Scarbrough, 1997a: 17, figs. 87-96. Type-locality: Dominican Republic, Pedernales, 35 km north of north of Cabo Rojo, Las Abejas, 1250 m. Distr. – Dominican Republic (Pedernales). Ref. – Scarbrough & Perez-Gelabert, 2003: 55, fig. 51. Type, USNM.
- willistoni** Curran, 1928a: 2 (in key), 3. Type-locality: Mexico, Guerrero, Tepetlapa, 3000 ft. Distr. – Mexico (Guerrero). Refs. – Hull, 1962 (2): 435; Martin & Papavero, 1970: 60 (cat.). Type, AMNH.

Proctacanthus-group

Genus **Eccritosia** Schiner, 1866

Eccritosia Schiner, 1866: 674. Type-species, *Asilus barbatus* Fabricius, 1787 (orig. des.).
Ecoritosia Bromley, 1951: 25, error.

Ref. – Lamas M., 1973c (rev.).

- barbata** (Fabricius), 1787: 358 (*Asilus*). Type-locality: “French Guiana, Cayenne (Rohr)” [cf. Papavero, 1971e: 20]. Distr. – U. S. A. (Texas), Mexico, Honduras, Costa Rica, Panama, Venezuela (Bolívar), Guyana, Peru (Loreto), Brazil, Bolivia, Paraguay, Argentina. Refs. – Olivier, 1789: 264 (*Asilus*); Gmelin, 1792: 2896 (*Asilus*); Fabricius, 1794: 379 (*Asilus*), 1805: 155 (*Asilus*); Wiedemann, 1821: 187 (*Asilus*), 1828: 439 (*Asilus*); Macquart, 1838: 122 (*Proctacanthus*); Walker, 1849: 652 (*Proctacanthus*), 1855: 703 (*Asilus*); Schiner, 1866: 703; Wulp, 1882b: 112; Hansen, 1883: pl. 4, figs. 1-3; Williston, 1891: 88 (cat.); Kertész, 1909: 249 (cat.); Curran, 1934a: 13 (in key), 14; Bromley, 1935c: 4 (*Erax*), 1946a: 116 (cat.); Carrera, 1947a: 48; Carrera & d’Andretta, 1953: 68; Carrera, 1960a: 164; Carrera & Vulcano, 1961: 74 (prey); Hull, 1962 (2): 489; Carrera & Machado-Allison, 1963: 253; Martin & Papavero, 1970: 61 (cat.); Lamas M., 1972: 313 (cat.), 1973c: 54, figs. 2, 8-10; Fisher, 1984: 34; Artigas & Papavero, 1995d: 147, figs. 33-34, 39. Type, KIEL [cf. Zimsen, 1964: 468].
- amphinome** Walker, 1849: 387 (*Asilus*), part. Type-locality: “Honduras”. Refs. – Osten Sacken, 1887: 207; Williston, 1901: 327; Aldrich, 1905: 275 (cat.); Kertész, 1909: 249 (cat.); Schaeffer, 1916: 66; Engel, 1930: 460; Curran, 1934a: 13 (in key), 14; Hull, 1962 (2): 489, figs. 390, 728, 1370, 1379, 2285, 2334, 2479, 2482; Martin & Wilcox, 1965: 393 (cat.); Martin & Papavero, 1970: 61 (cat.). Syntypes, BMNH.
- tricolor** Walker, 1850: xcixii (*Mallophora*). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Walker, 1855: 583 (*Mallophora*); Williston, 1891: 88 (cat.; *Mallophora*); Curran, 1934a: 14 (may be syn. of *barbata* Fabricius, 1787); Bromley, 1946: 116 (cat.); Carrera & d’Andretta, 1952: 258 (prey). Type, BMNH.
- plinthopyga** (Wiedemann), 1821: 184 (*Asilus*). Type-locality: “Cuba” (possibly in error). Distr. – Brazil (Rio Grande do Sul), Uruguay (Montevideo, San José). Refs. – Wiedemann, 1828: 432 (*Asilus*); Walker, 1855: 695 (*Asilus*); Bigot, 1857a: 331 (*Asilus*); Schiner, 1866: 683, 1867: 399; Aldrich, 1905: 275 (cat.); Kertész, 1909: 250 (cat.); Curran, 1934a: 15; Hull, 1962 (2): 489; Martin & Papavero, 1970: 61 (cat.); Lamas M., 1973c: 62, figs. 4, 14-16; Alayo Dalmau & Garcia Ávila, 1983: 48 (cat.); Artigas & Papavero, 1995d: 147, fig. 37. Type, NMW.
- vetustus** Walker, 1837: 340 (*Asilus*). Type-locality: Uruguay, Gorriti Island. Refs. – Walker, 1849: 420 (*Proctacanthus*), 1855: 653 (*Proctacanthus*); Lynch Arribálzaga, 1880: 260 (*Proctacanthus*), 1883a: 148 (cat.; *Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 248 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*). Type, BMNH.
- ruberiventris** (Macquart), 1850: 391 (1850: 87), pl. 8, fig. 3 (*Proctacanthus*). Type-locality: “Brazil”. Distr. – Brazil (São Paulo, Santa Catarina, Rio Grande do Sul), Uruguay (Canelones), Argentina (Salta, Tucumán, Mendoza), Chile (Arauco, Cautín, Concepción, Malleco, Ñuble, Santiago, Tarapacá); Australia (New South Wales, introduced). Refs. – Schiner, 1866: 714 (*Proctacanthus*); Rondani, 1868: 33 (*Proctacanthus*); Lynch Arribálzaga, 1880: 259 (*Proctacanthus*); Wulp, 1882b: 108 (*Proctacanthus*); Lynch Arribálzaga, 1883a: 147 (cat.; *Proctacanthus*), 1883b: 12 (*Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 248 (cat.; *Proctacanthus*); Bromley, 1946: 116 (cat.; *Proctacanthus*); Carrera, 1960b: 50 (n. comb.); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas, 1970a: 279, figs. 371-377, 410, 1971: 35, figs. 83-84; Martin & Papavero, 1970: 61 (cat.); Lamas M., 1973c: 59, figs. 3, 11-13; Artigas & Reyes, 1975: 201, figs. 3, 7-9, 11, 13; Artigas & Papavero, 1995d: 147, figs. 15-18, 32, 35, 38. Type, MNHNP.
- xanthopogon** Burmeister, 1861 (vol. I): 317, (vol. II): 171 (*Asilus (Proctacanthus)*). Type-locality: Argentina, Mendoza and Tucumán, Monteros TP MACN.
- speciosus** Philippi, 1865: 193, pl. 26, fig. 28 (*Erax*). Type-locality: Chile, Colchagua, near Llico. Refs. – Schiner, 1866: 713 (*Erax*); Stuardo Ortiz, 1946: 84 (cat.; n. comb.). Type, lost.
- barbiellini** Curran, 1934a: 13. Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 116 (cat.); Hull, 1962 (2): 489. Type, AMNH.
- wirthi** Paramonov, 1964: 157. Type-locality: Australia, New South Wales, Dee Why, near Sydney. Type, CSIRO.
- zamon** (Townsend), 1895: 600 (*Proctacanthus*). Lectotype-locality: Mexico, Baja California Sur, San José del Cabo. Distr. – U. S. A. (Arizona, Texas), Mexico (Baja California, Chihuahua, Guerrero, Jalisco, Oaxaca), Guatemala, Honduras, Nicaragua, Costa Rica.. Refs. – Kertész, 1909: 249 (cat.; as syn. of *amphinome*); Curran, 1934a: 13 (in key), 14; Wilcox, 1937: 45; Bromley, 1951: 35, figs. 2, 8; Hull, 1962 (2): 489; Martin & Wilcox, 1965: 393 (cat.); Lamas M., 1973c: 53, figs. 1, 5-7; Lavigne, 1977: 626 (ethology); Artigas & Papavero, 1995d: 147, fig. 36. Lectotype AMNH [des. by Bromley, 1951: 26].
- amphinome** Walker, 1849: 387 (*Asilus*), part. Type-locality: “Honduras”. Refs. – Osten Sacken, 1887: 207; Williston, 1901: 327; Aldrich, 1905: 275 (cat.); Kertész, 1909: 249 (cat.); Bromley, 1934a: 106; Martin & Papavero, 1970: 61 (cat.). Syntypes, BMNH.

Genus **Proctacanthella** Bromley, 1934

Proctacanthella Bromley, 1934a: 96. Type-species, *Asilus cacopilodus* Hine, 1909 (orig. des.).

Ref. – Artigas & Papavero, 1995d: 153, figs. 1-8, 154, figs. 9-14.

exquisita (Osten Sacken), 1887: 206, pl. 3, fig. 12 (*Proctacanthus*). Type-locality: Mexico, northern Sonora. Distr. – U. S. A., Mexico (Sonora). Refs. – Aldrich, 1905: 274 (cat.; *Proctacanthus*); Kertész, 1909: 246 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Martin & Wilcox, 1965: 398 (cat.); Wilcox, 1965a: 108 (in key), fig. 1; Martin & Papavero, 1970: 82 (cat.); Artigas & Papavero, 1995d: 147; Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.

mexicanus Cole, 1923: 463, fig. 6 (*Stenopogon*). Type-locality: Mexico, Sonora, Guaymas. Ref. – Hull, 1962 (1): 124 (*Scleropogon*). Type, CAS.

jamesi Pritchard, 1935: 13. Type-locality: U. S. A., Oklahoma, Kenton. Ref. – Hull, 1962 (2): 499. Type, AMNH.

robusta Bromley, 1951: 26, fig. 3. Type-locality: U. S. A., Texas, Port Lavaca. Distr. – U. S. A. (Texas), Mexico. Ref. – Hull, 1962 (2): 499; Wilcox, 1965a: 107 (in key), 108, fig. 1. Type, OSU.

taina Scarbrough & Perez-Gelabert, 2006: 12, figs. 1-2, 188. Type-locality: Dominican Republic, Montecristi, 4 km north of Botoncillo. Distr. – Dominican Republic (Bahoruco, Montecristi). Type, FSCA.

tolandi Wilcox, 1965a: 107 (in key), 109, fig. 1. Type-locality: U. S. A., California, Riverside Co., White Water. Distr. – U. S. A. (California), Mexico. Type, CAS.

Genus **Proctacanthus** Macquart, 1838

Proctacanthus Macquart, 1838: 120 (1839: 236). Type-species, *Proctacanthus philadelphicus* Macquart, 1838 (sub. des.; Coquillett, 1910: 595).

Acanthodelphia Bigot, 1857b: 545 (in key). No species included.

Proctacantha Scudder, 1884: 262, emend.

Ref. – Lavigne, Nelson & Schreiber, 1994 (preys and prey choice).

arno Townsend, 1895: 599. Lectotype-locality: Mexico, Baja California Sur, San José del Cabo. Distr. – U. S. A. (Texas, Utah); Mexico (Baja California). Refs. – Aldrich, 1905: 274 (cat.); Kertész, 1909: 245 (cat.); Hine, 1911: 161, 1918: 3; Brown, 1929: 309; Bromley, 1935c: 3; Hull, 1962 (2): 488; Martin, 1962a: 186; Martin & Papavero, 1970: 77 (cat.); Artigas & Papavero, 1995d: 148. Lectotype KU [des. by Martin, 1962a: 186].

basifascia Walker, 1855: 654. Type-locality: Brazil, Pará, Santarém. Distr. – Brazil (Pará). Refs. – Williston, 1891: 87 (cat.); Kertész, 1909: 246 (cat.); Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148.

bifascia Bromley, 1946: 115 (cat.), error.

caudatus Hine, 1911: 159. Type-locality: Mexico, Veracruz. Distr. – Mexico (Veracruz). Refs. – Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148. Type, OSU.

coprates Walker, 1848: 409. Type-locality: unknown. Refs. – Walker, 1855: 652; Williston, 1891: 88 (cat.); Kertész, 1909: 246 (cat.); Bromley, 1946: 115 (cat.); Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.).

coquilletti Hine, 1911: 160. Type-locality: U. S. A., California, Los Angeles. Distr. – U. S. A. (California), Mexico. Refs. – Hull, 1962 (2): 488; Martin & Wilcox, 1965: 399 (cat.). Type, OSU.

craverii Bellardi, 1861: 150 (50). Type-locality: Mexico, D. F., surroundings of Mexico City. Distr. – Mexico (D. F., Jalisco, Veracruz). Refs. – Williston, 1901: 327; Aldrich, 1905: 274 (cat.); Kertész, 1909: 274 (cat.); Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148; Papavero & Ibáñez Bernal, 2001: 106 (hist. Mexican dipterol). Type, TORO.

danforthi Curran, 1951: 5 (in key), 8. Type-locality: Puerto Rico, Arecibo. Distr. – Puerto Rico. Refs. – Hull, 1962 (2): 488; Maldonado-Capriles & Navarro, 1967: 52; Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148. Type, AMNH.

darlingtoni Curran, 1951: 5 (in key), 8. Type-locality: Dominican Republic, Barahona. Distr. – Dominican Republic (Barahona, Peravia). Refs. – Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148; Scarbrough & Perez-Gelabert, 2006: 14. Type, MCZ.

dominicarus Curran, 1951: 5 (in key), 9 (as *dominicana*, error). Type-locality: Dominican Republic, Samaná, Sanchez. Distr. – Dominican Republic (Azua, La Altagracia, Pedernales, Peravia, San Cristóbal, Santo Domingo). Refs. – Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148; Scarbrough & Perez-Gelabert, 2006: 15 (as *dominicana*, error). Type, AMNH.

lernerii Curran, 1951: 5. Type-locality: Bahama Islands, South Bimini Island. Distr. – Bahamas Islands South Bimini Island). Refs. – Martin, 1957b: 7; Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148, figs.

- 49-51, 52-56. Type, AMNH.
- micans** Schiner, 1867: 397. Type-locality: "North America". Distr. – U. S. A. (Arizona, Colorado, New Mexico), Mexico. Refs. – Kertész, 1909: 247 (cat.); Hine, 1911: 160; Hull, 1962 (2): 488; Dennis & Lavigne, 1976 (pupa); Artigas & Papavero, 1995d: 157, figs. 40-42. Type, NMW.
- nearno** Martin, 1962a: 187. Type-locality: U. S. A., Arizona, Baboquivari Mountains. Distr. – U. S. A. (Arizona, California, New Mexico, Texas, Utah). Ref. – Lavigne & Dennis, 1979: 438 (ethology). Type, KU.
- nigrimanus** Curran, 1951: 5 (in key), 7. Type-locality: Cuba, Oriente, Maisi. Distr. – Cuba. Ref. – Hull, 1962 (2): 488; Martin & Papavero, 1970: 78 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 50 (cat.); Artigas & Papavero, 1995d: 148. Type, MCZ.
- nigrofemoratus** Hine, 1911: 161. Type-locality: Mexico, San José. Distr. – U. S. A. (Texas), Mexico. Refs. – Hull, 1962 (2): 488; Martin & Wilcox, 1965: 399 (cat.); Martin & Papavero, 1970: 78 (cat.); Artigas & Papavero, 1995d: 148. Type, OSU.
- vittatus** (Olivier), 1789: 263 (*Asilus*). Type-locality: "Saint Domingue". Distr. – Dominican Republic (La Altagracia, La Vega, National District), Haiti. Refs. – Aldrich, 1905: 283 (cat.; *Asilus*); Kertész, 1909: 265 (cat.; *Asilus*); Bromley, 1929a: 293; Curran, 1934a: 8 (in key), 1951: 5 (in key), 7; Hull, 1962 (2): 488; Scarbrough & Perez-Gelabert, 2006: 15. Type, lost.
- rufiventris** Macquart, 1838: 123 (1839: 239), pl. 10, fig. 2. Type-locality: "Dominican Republic". Refs. – Walker, 1849: 409, 1855: 650; Roeder, 1885: 339; Aldrich, 1905: 274 (cat.); Kertész, 1909: 248 (cat.); Hine, 1911: 158; Curran, 1928b: 23, 1951: 7; Martin & Papavero, 1970: 79 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 50 (cat.); Artigas & Papavero, 1995d: 148. Type, MNHNP.

Genus *Taurhynchus* Artigas & Papavero, 1995

Taurhynchus Artigas & Papavero, 1995d: 148. Type-species, *Asilus xanthopterus* Wiedemann, 1828 (orig. des.).

- antidomus** (Walker), 1849: 408 (*Asilus*). Type-locality: "Brazil". Refs. – Walker, 1855: 652; Williston, 1891: 87 (cat.; *Proctacanthus*); Kertész, 1909: 245 (cat.; *Proctacanthus*); Curran, 1934a: 14 (?*Eccritosia*); Bromley, 1946: 115 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Martin & Papavero, 1970: 61 (?*Eccritosia*); Artigas & Papavero, 1995d: 149 (n. comb.).
- aurolineatus** (Macquart), 1846: 214 (1846: 86), pl. 8, fig. 9 (*Proctacanthus*). Type-locality: "Colombia" (as "Nouvelle Grenade"). Distr. – Colombia. Refs. – Walker, 1855: 653 (*Proctacanthus*); Williston, 1891: 87 (cat.; *Proctacanthus*); Kertész, 1909: 245 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Martin & Papavero, 1970: 77 (cat.); Artigas & Papavero, 1995d: 149 (n. comb.). Type, lost.
- barbiventris** (Rondani), 1850: 187 (*Asilus*). Type-locality: Brazil, São Paulo, Ilha de São Sebastião. Refs. – Osten Sacken, 1891: 421 (*Asilus*); Kertész, 1909: 252 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, TORO.
- bromleyi** (Curran), 1931b: 22 (*Proctacanthus*). Type-locality: Colombia, Magdalena, Rio Frio. Distr. – Colombia (Magdalena), Venezuela (Aragua) Refs. – Curran, 1934a: 8 (in key); Hull, 1962 (2): 488 (*Proctacanthus*); Carrera & Machado-Allison, 1963: 253 (*Proctacanthus*); Artigas & Papavero, 1995d: 149, figs. 19-21 (n. comb.). Type, AMNH.
- camposi** (Curran), 1934a: 9 (in key), 10 (*Proctacanthus*). Type-locality: Ecuador, Guayas, Posorja. Refs. – Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, AMNH.
- cruentus** (Lynch Arribálzaga), 1880: 264 (*Proctacanthus*). Type-locality: Argentina, Misiones. Refs. – Lynch Arribálzaga, 1883a: 147 (cat.; *Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 246 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.).
- daraps** (Walker), 1849: 416 (*Asilus*). Type-locality: "Brazil". Refs. – Walker, 1855: 653 (*Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 246 (cat.; *Proctacanthus*); Bromley, 1946: 115 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, BMNH (tip of abdomen missing. Walker did not declare the sex in the original description, so probably the specimen was broken since that time).
- dina** (Curran), 1934a: 9 (*Proctacanthus*). Type-locality: Brazil, Amazonas, rio Caiari, Uaupés. Distr. – Venezuela (Carabobo, Distrito Federal), Brazil (Amazonas). Refs. – Bromley, 1946: 115 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Carrera & Machado-Allison, 1963: 253 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, AMNH.
- fervidus** (Curran), 1934a: 9 (in key), 11 (*Proctacanthus*). Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 115 (cat.; *Proctacanthus*); Carrera, 1947d: 271 (*Proctacanthus*) (preys); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149, figs. 22-24, 25-31 (n. comb.). Type, AMNH.
- flavipennis** (Macquart), 1846: 213 (1846: 85), pl. 8, fig. 11 (*Proctacanthus*). Type-locality: "Colombia". Refs. – Walker, 1855: 653 (*Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 246 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, lost.
- guianicus** (Curran), 1934a: 9 (in key), 12 (*Proctacanthus*; as *guianica*, error). Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Refs. – Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, AMNH.
- hagno** Walker, 1849: 417 (*Asilus*). Type-locality: "Brazil". Refs. – Walker, 1855: 653 (*Proctacanthus*); Williston, 1891: 88

- (cat.; *Proctacanthus*); Kertész, 1909: 246 (cat.; *Proctacanthus*); Bromley, 1946: 115 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, BMNH (a male, according to the original description. Now tip of abdomen missing).
- leonides** (Walker), 1851: 150 (*Asilus*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Refs. – Walker, 1855: 705 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 258 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Artigas & Papavero, 1995d: 149 (n. comb.).
- leucopogon** (Wiedemann), 1828: 430 (*Asilus*). Type-locality: Uruguay, Montevideo. Distr. – Uruguay. Refs. – Schiner, 1866: 683 (*Proctacanthus*), 1867: 395 (*Proctacanthus*); Lynch Arribálzaga, 1880: 261 (*Proctacanthus*, 1883a: 147 (cat.; *Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 247 (cat.; *Proctacanthus*); Hine, 1918: 7; Hull, 1962 (2): 488 (*Proctacanthus*), 499 [*Proctacanthella*], fig. 377, 2303, 2314; Martin & Papavero, 1970: 78 (cat.); Lavigne & Dennis, 1980: 160 (*Proctacanthus*; ethology); Shelly & Pearson, 1980: 7 (*Proctacanthella*; predatory behavior), 1983: 520 (diurnal variation in predatory behavior). Type, NMW. N. comb.
- ?**macrotelus** Walker, 1837: 340 (*Asilus*). Type-locality: Uruguay, Gorriti Island. Refs. – Walker, 1849: 420 (*Asilus*), 1855: 653 (*Proctacanthus*); Lynch Arribálzaga, 1880: 260 (*Proctacanthus*), 1883a: 147 (cat.; *Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 247 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*).
- mystaceus** (Macquart), 1846: 214 (1846: 86) (*Proctacanthus*). Type-locality: “Colombia”. Refs. – Walker, 1855: 652 (*Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 244 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, lost.
- rubricornis** (Macquart), 1838: 122 (1839: 238) (*Proctacanthus*). Type-locality: Brazil, Rio de Janeiro. Distr. – Brazil (Rio de Janeiro), Paraguay. Refs. – Walker, 1855: 652 (*Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 248 (cat.; *Proctacanthus*); Bromley, 1946: 115 (cat.; *Proctacanthus*); Carrera, 1947a: 47 (*Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, MNHNP.
- salti** (Curran), 1934a: 9 (in key), 11 (*Proctacanthus*). Type-locality: Colombia, Magdalena, Río Frío. Refs. – Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, AMNH.
- tibialis** (Macquart), 1850: 392 (1850: 88) (*Proctacanthus*). Type-locality: “South America”. Refs. – Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 248 (cat.; *Proctacanthus*); Curran, 1934a: 10 (considers unrecognizable); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.). Type, lost.
- vittatus** (Lynch Arribálzaga), 1880: 261 (*Proctacanthus*). Type-locality: Argentina, Misiones. Refs. – Lynch Arribálzaga, 1883a: 148 (cat.; *Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 249 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Artigas & Papavero, 1995d: 149 (n. comb.).
- xanthopterus** (Wiedemann), 1828: 427 (*Asilus*). Type-locality: “Brazil” [Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-102]. Refs. – Macquart, 1838: 121 (*Proctacanthus*); Schiner, 1866: 682 (*Proctacanthus*); Williston, 1891: 88 (cat.; *Proctacanthus*); Kertész, 1909: 249 (cat.; *Proctacanthus*); Bromley, 1946: 116 (cat.; *Proctacanthus*); Hull, 1962 (2): 488 (*Proctacanthus*); Martin & Papavero, 1970: 79 (cat.); Artigas & Papavero, 1995d: 149. Syntypes, BZM [no. 738; cf. Schumann, 1973: 170].

Unplaced or unrecognized species of Asilinae

- albisetosus** Wulp, 1882b: 116 (*Epitriptus*). Type-locality: “Argentina (Weyenbergh)” [cf. Papavero, 1973d: 357]. Refs. – Lynch Arribálzaga, 1883b: 17 (*Epitriptus*); Williston, 1891: 89 (cat.; *Epitriptus*); Kertész, 1909: 294 (cat.; *Epitriptus*); Hull, 1962 (2): 5673 (*Epitriptus*). Type, ZMA.
- americanus** Macquart, 1846: 215 (1846: 87) (*Lophonotus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 701 (*Asilus*); Williston, 1891: 89 (cat.; *Lophonotus*); Kertész, 1909: 274 (cat.; *Lophonotus*); Bromley, 1946: 117 (cat.; *Neolophonotus* (*Dysmachus*)); Hull, 1962 (2): 567 (*Dysmachus*); Martin & Papavero, 1970: 87 (cat.; *Dysmachus*). Type, OXF.
- aurivestitus** Hine, 1919: 142, fig. 42 (*Erax*). Type-locality: Mexico, Morelos. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 63 (cat.; *Efferia*); Artigas & Papavero, 1997b: 90 (*Efferia*-group). Type, OSU.
- flavescens** James, 1953: 46, fig. 1 (*Eicherax*). Type-locality: Honduras, Olanchito. Refs. – Hull, 1962 (2): 475 (*Eicherax*); Martin & Papavero, 1970: 70 (consider genus misident.). Type, MICH.
- pavidus** Williston, 1901: 326 (*Erax*). Type-locality: Mexico, Guerrero, Xucumanatlán. Refs. – Aldrich, 1905: 277 (cat.; *Erax*); Kertész, 1909: 241 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 67 (*Efferia*); Artigas & Papavero, 1997b: 90; Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.
- schadei** Bromley, 1951: 30, fig. 5 (*Erax*). Type-locality: Paraguay, Hape [?]. Refs. – Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 68 (*Efferia*); Artigas & Papavero, 1997b: 90. Type, AMNH.
- singularis** Macquart, 1838: 111 (1839: 227), pl. 9, fig. 10 (*Erax*). Type-locality: Western part of Uruguay, along Uruguay river (as “Uruguay, depuis l’embouchure de l’Uruguay jusqu’aux Missions”). Refs. – Walker, 1855: 634 (*Erax*); Lynch Arribálzaga, 1880: 111 (*Erax*), 1883b: 150 (*Erax*); Williston, 1891: 87 (cat.; *Erax*); Kertész, 1909: 242 (cat.; *Erax*); Hull, 1962 (2): 478 (*Nerax*); Martin & Papavero, 1970: 68 (*Efferia*); Artigas & Papavero, 1997b: 90. Type, MNHNP.
- spissibarbis** Macquart, 1846: 208 (1846: 80) (*Trupanea*). Type-locality: “Chile”. Refs. – Walker, 1866: 589 (*Trupanea*);

Williston, 18791: 84 (cat.; *Promachus*); Kertész, 1909: 226 (cat.; *Promachus*); Hull, 1962 (2): 462 (*Promachus*); Martin & Papavero, 1970: 80 (cat.; *Promachus*).

strigitibia Curran, 1931b: 25 (*Dysmachus*). Type-locality: Ecuador, Riobamba. Refs. – Hull, 1962 (2): 567 (*Dysmachus*); Martin & Papavero, 1970: 87 (cat.; *Dysmachus*). Type, AMNH.

Unrecognized genera of Asilinae

Genus **Labromyia** Hull, 1962

Labromyia Hull, 1962 (2): 533. Type-species, *Labromyia albobarbis* Hull, 1962 (orig. des.).

albobarbis Hull, 1962 (2): 534, figs. 341, 722, 1466, 2266, 2327, 2425, 2435. Type-locality: Paraguay, Cordillera, San Bernardino. Ref. – Martin & Papavero, 1970: 83 (cat.). Type, ZSBS.

Genus **Pediophoneus** Lynch Arribálzaga, 1883

Pediophoneus Lynch Arribálzaga, 1883b: 87. No species included. Refs. – Kertész, 1909: 267 (cat.); Hull, 1962 (2): 519; Martin & Papavero, 1970: 81 (cat.).

Subfamily Dasypogoninae Macquart, 1838

Dasypogonitae Macquart, 1838: 22 (1839: 138). Type-genus, *Dasypogon* Meigen, 1803.

Dasypogonidae Bigot, 1852: 485.

Dasypogonites Macquart, 1850: 472.

Dasypogonidii Bigot, 1853: 300.

Dasypogonina Loew, 1858: 342.

Dasipogonina Rondani, 1863: 52, 156.

Dasypoginæ Wulp, 1896: 75.

Dasypogomnia, *Dasipogonia* Valdés-Raguès, 1908: 309-310.

Saropogonini G. H. Hardy, 1926: 306. Type-genus, *Saropogon* Loew, 1847.

Dasipogoninae Karl, 1935: 115.

Saroponini Martin & Papavero, 1970: 23 (cat.), error.

Tribe Dasypogonini Macquart, 1838

Genus **Aczelia** Carrera, 1955

Aczelia Carrera, 1955c: 297. Type-species, *Laparus argentinus* Wulp, 1882 (orig. des.).

argentina (Wulp), 1882b: 95 (*Laparus*). Type-locality: “Argentina”. Distr. – Argentina (La Rioja, Mendoza, Salta, Tucumán).

Refs. – Williston, 1891: 74 (cat.; *Neolaparus*); Brèthes, 1908: 287 (*Neolaparus*); Kertész, 1909: 120 (cat., *Neolaparus*); Carrera, 1955c: 299, figs. 1-3 (n. comb.); Carrera & Papavero, 1962: 54; Hull, 1962 (1): 279, fig. 558; Martin & Papavero, 1970: 36 (cat.); Artigas & Papavero, 1988b: 240, figs. 108-110, 242, fig. 116. Type, ZMA.

infumata (Lynch Arribálzaga), 1880: 183 (*Allopogon*). Type-locality: Argentina, Buenos Aires, San José de Flores. Distr. – Argentina (Buenos Aires). Refs. – Lynch Arribálzaga, 1883a: 137 (cat.; *Allopogon*); Williston, 1891: 73 (cat.; *Allopogon*); Kertész, 1909: 118 (cat.; *Allopogon*); Hull, 1962 (1): 279, fig. 76, 1126, 1135, 1707-1716-1717, 1880, 2031; Martin & Papavero, 1970: 37 (cat.).

tsacasi Papavero, 1971d: 23, figs. 1-3. Type-locality: Argentina, Santiago del Estero, Averias, rio Salado. Distr. – Argentina (Santiago del Estero). Ref. – Artigas & Papavero, 1988b: 240, figs. 105-107. Type, MNHNP.

Genus **Amorimius**, nom. nov.

Macrocolus Engel, 1930: 470 (preocc. Wagner, 1846). Type-species, *Macrocolus bicolor* Engel, 1930 (orig. des., as gen. n., sp. n.).
Amorimius, nom. nov. for *Macrocolus* Engel, 1930. Type-species, *Macrocolus bicolor* Engel, 1930 (aut.). [Dedicated to Prof. Dr. Dalton de Souza Amorim, Departamento de Biologia, Faculdade de Filosofia, Ciências e Letras, Universidade de São Paulo, Ribeirão Preto].

Refs. – Hull, 1962 (1): 132, figs. 125, 477, 900, 1702, 1847; Artigas & Papavero, 1988b: 211 (key to spp.).

barrettoi (Carrera), 1949: 120, fig. 180 (*Macrocolus*). Type-locality: Brazil, Minas Gerais, Araçuaí. Distr. – Brazil (Minas Gerais). Refs. – Hull, 1962 (1): 133 (*Macrocolus*); Martin & Papavero, 1970: 23 (cat.; *Macrocolus*). Type, MZUSP. N. comb.

bicolor (Engel), 1930: 471, fig. 8 (*Macrocolus*). Type-locality: Bolivia, Santa Cruz, Santa Cruz de la Sierra and Ipitás; Paraguay, Cordillera, San Bernardino. Distr. – Bolivia (Santa Cruz), Brazil (São Paulo), Paraguay (Cordillera). Refs. – Carrera, 1949: 119, figs. 8, 61, 91, 119, 137-138, 179 (*Macrocolus*); Carrera & Papavero, 1962: 55 (*Macrocolus*); Hull, 1962 (1): 133 (*Macrocolus*); Martin & Papavero, 1970: 23 (cat.; *Macrocolus*); Artigas & Papavero, 1988b: 244, figs. 124-125 e 127, 245, figs. 128-132 (*Macrocolus*). N. comb.

martinorum (Artigas & Papavero), 1988b: 211, 244, fig. 126 (*Macrocolus*). Type-locality: Mexico, Guerrero, 24 miles south of Iguala. Distr. – Mexico (Guerrero). Type, MZUSP. N. comb.

rubripes (Carrera & Papavero), 1962: 55, fig. 2 (*Macrocolus*). Type-locality: Brazil, São Paulo, Campos do Jordão (Eugênio Lefevre). Distr. – Brazil (São Paulo). Type, MZUSP. N. comb.

Genus **Araucopogon** Artigas & Papavero, 1988

Araucopogon Artigas & Papavero, 1988b: 209. Type-species, *Dasypogon* (*Saropogon*) *cyanogaster* Loew, 1851 (orig. des.).

cyanogaster (Loew), 1851: 5 (*Dasypogon* (*Saropogon*)). Type-locality: “Chile (Eschscholtz)” [cf. Papavero, 1971e: 51]. Distr. – Chile (Aconcagua, Arauco, Cautín, Colchagua, Linares, Malleco, Maule, Ñuble, O’Higgins, Santiago, Talca, Valparaíso). Refs. – Williston, 1891: 74 (cat.; *Sarapogon*); Kertész, 1909: 137 (cat.); Carrera, 1953a: 275 (*Araiopogon*); Hull, 1962: 282 (*Araiopogon*); Artigas, 1970a: 68, figs. 168, 170-171, 438 (*Araiopogon*); Martin & Papavero, 1970: 36 (cat.; *Araiopogon*); Artigas, 1971: 9, figs. 15, 16 (*Araiopogon*); Artigas & Papavero, 1988b: 241, figs. 111-115, 242, fig. 117. Type, BZM [no. 433; cf. Schumann, 1973: 100].

pictus Philippi, 1865: 638 (*Dasypogon*). Type-locality: Chile, Colchagua, Colchagua. Refs. – Williston, 1891: 74 (cat.; *Sarapogon*); Kertész, 1909: 139 (cat.; *Saropogon*); Bromley, 1932: 266 (*Saropogon*); Stuardo Ortiz, 1946: 82 (cat.; *Saropogon*); Carrera, 1953a: 275 (syn.). Type, MNHNS (headless).

Genus **Austenmyia** Carrera, 1955

Austenmyia Carrera, 1955a: 104. Type-species, *Austenmyia amazona* Carrera, 1955 (orig. des.).

amazona Carrera, 1955a: 105, fig. 1. Type-locality: Brazil, Amazonas, Manaus. Distr. – Brazil (Amazonas). Refs. – Hull, 1962 (1): 281, figs. 519, 1123, 1132; Martin & Papavero, 1970: 37 (cat.); Artigas & Papavero, 1988b: 239, fig. 98. Type, BMNH.

Genus **Cleptomyia** Carrera, 1949

Cleptomyia Carrera, 1949: 124. Type-species, *Cleptomyia bacillifera* Carrera, 1949 (orig. des.) = *Cleptomyia tripartita* (Walker, 1854).

tripartita (Walker), 1854: 437 (*Dasypogon*). Type-locality: Brazil, Pará, Santarém. Distr. – Brazil (Pará, Ceará, Rio Grande do Norte), Paraguay, Argentina. Ref. – Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 132 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Papavero, 1971d: 21 (n. comb.); Artigas & Papavero, 1988: 239, figs. 99 e 102, 242, fig. 118, 243, figs. 110-123. Type, BMNH.

bacillifera Carrera, 1949: 125, figs. 5, 63, 99, 155, 157, 184, pl. 16. Type-locality: Brazil, Ceará, Russas. Refs. – Carrera, 1959: 2; Carrera & Papavero, 1962: 61; Hull, 1962 (1): 260, figs. 577, 1144, 1153; Martin & Papavero, 1970: 32 (cat.);

Papavero

Papavero, 1971d: 21 (n. syn.). Type, MZUSP.

Genus **Neoderomyia** Artigas, 1971

Neoderomyia Artigas, 1971: 12. Type-species, *Deromyia fulvipes* Philippi, 1865 (orig. des.).

fulvipes (Philippi), 1865: 706 (*Deromyia*). Type-locality: Chile, Santiago. Distr. – Chile (Biobío, Maule, Santiago, Valparaíso). Refs. – Williston, 1891: 75 (cat.; *Deromyia*); Kertész, 1909: 116 (cat.; *Deromyia*); Stuardo Ortiz, 1946: 82 (cat.; *Deromyia*); Hull, 1962 (1): 250 (*Deromyia*); Artigas, 1970a: 154, figs. 135–137, 139–141, 435 (*Oberon*); Martin & Papavero, 1970: 37 (cat.; *Oberon*); Artigas, 1971: 12, figs. 17–18 (n. comb.); Artigas & Papavero, 1988b: 246, figs. 133–137.

Genus **Parataracticus** Cole, 1924

Parataracticus Cole, 1924: 11. Type-species, *Parataracticus rubidus* Cole, 1924 (orig. des.).

Ref. – Artigas & Papavero, 1988b: 247, figs. 138–142, 248, fig. 143.

arenicolus Martin, 1968d: 182. Type-locality: Mexico, Baja California, Los Friales. Distr. – Mexico (Baja California). Refs. – Martin & Papavero, 1970: 35 (cat.); Artigas & Papavero, 1988b: 239, fig. 103. Type, CAS.

niger Martin, 1955b: 118. Type-locality: U. S. A., California, Riverside County, Perris. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Hull, 1962 (1): 272; Martin & Wilcox, 1965: 381 (cat.); Martin & Papavero, 1970: 35 (cat.). Type, CAS.

Genus **Saropogon** Loew, 1847

Dasypogon, subg. **Saropogon** Loew, 1847: 439. Type-species, *Dasypogon luctuosus* Wiedemann, 1820 (sub. des., Coquillett, 1910: 603).

Sarapogon Williston, 1891: 74 (cat.), error.

Araiopogon Carrera, 1949: 122. Type-species, *Dasypogon gayi* Macquart, 1838 (orig. des.).

Oberon Carrera & Papavero, 1962: 57. Type-species, *Oberon velutinus* Carrera & Papavero, 1962 (orig. des.).

One species known as fossil: *Saropogon oblitescens* Cockerell (1914: 102) (Oligocene; U. S. A., Colorado).

albifrons Back, 1904: 291. Type-locality: U. S. A., Arizona, Bill Williams Fork. Distr. – U. S. A. (Arizona), Mexico (Baja California). Refs. – Kertész, 1909: 136 (cat.); Hull, 1962 (1): 278 (as syn. of *semiustus* Coquillett, 1904); Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 130 (in key); Martin & Papavero, 1970: 35 (cat.).

combustus Loew of Back, 1909: 351, part, misident.

bryanti Wilcox, 1966a: 130 (in key), 132. Type-locality: U. S. A., Arizona, Baboquivari Canyon, west side of Baboquivari Mountains. Distr. – U. S. A. (Arizona, New Mexico), Mexico. Type, CAS.

carbonarius (Philippi), 1865: 687 (*Dasypogon*). Type-locality: Chile, Colchagua and Valdivia. Distr. – Chile. Refs. – Bigot, 1878: 221; Williston, 1891: 68 (cat.; *Sarapogon*); Kertész, 1909: 136 (cat.); Bromley, 1932: 267 (*Dasypogon*); Hull, 1962 (1): 278; Martin & Papavero, 1970: 36 (cat.; *Araiopogon*).

choapensis (Artigas), 1970a: 67, fig. 442 (*Araiopogon*). Type-locality: Chile, Coquimbo, Choapa. Distr. – Chile (Coquimbo). Type, DZUC.

combustus Loew, 1874: 373. Type-locality: U. S. A., Texas. Distr. – U. S. A. (Colorado, Nebraska, Kansas, Oklahoma, Texas), Mexico (Coahuila). Refs. – Osten Sacken, 1874: 185; Williston, 1883: 23; Bromley, 1935: 3; Baker, 1939: 168; Hull, 1962 (1): 277; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 129, 130 (in key). Type, MCZ.

adustus Loew, 1874: 375. Type-locality: U. S. A., Texas. Ref. – Osten Sacken, 1874: 185.

coquillettii Back, 1909: 348. Type-locality: U. S. A., New Mexico. Distr. – U. S. A. (Arizona, New Mexico), Mexico (Chihuahua). Refs. – Curran, 1930b: 3 (in key), 1931b: 2 (in key); Hull, 1962 (1): 277; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 129 (in key); Martin & Papavero, 1970: 35 (cat.). Type, USNM.

dispar Coquillett, 1902: 139. Type-locality: U. S. A., Texas. Distr. – U. S. A. (Georgia, Oklahoma, Texas), Mexico (Tamaulipas).

Refs. – Aldrich, 1905: 263 (cat.); Kertész, 1909: 137 (cat.); Hine, 1918: 3; Curran, 1930b: 2, 3 (in key), 1931b: 2; Fattig, 1945: 17; Hull, 1962 (1): 277; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 129 (in key); Martin & Papavero, 1970: 35 (cat.); Artigas & Papavero, 1988b: 248, fig. 144, 249, figs. 146–150. Type, USNM.

fraternus Bigot, 1878: 414. Type-locality: “Chile”. Neotype-locality: Chile, Choapa, Illapel. Distr. – Chile (Arauco, Coquimbo, Santiago). Refs. – Williston, 1891: 74 (cat.; *Sarapogon*); Bromley, 1932: 266; Stuardo Ortiz, 1946: 82 (cat.); Hull, 1962 (1): 278; Artigas, 1970a: 72, figs. 166–167, 432 (*Araiopogon*), 1971: 9, figs. 5–6 (*Araiopogon*); Martin & Papavero, 1970: 36 (cat.; *Araiopogon*). Type, lost. NT DZUC [des. by Artigas, 1970a: 72].

fulvicornis (Macquart), 1850: 372 (1850: 68), pl. 6, fig. 12 (*Dasypogon*). Type-locality: Chile, Coquimbo. Distr. – Chile (Coquimbo, Santiago). Refs. – Blanchard, 1852: 366 (*Dasypogon*); Philippi, 1865: 685 (*Dasypogon*); Schiner, 1868: 163 (*Scylaticus*); Bigot, 1878: 221 (*Triclis*); Brauer, 1885: 385, pl. 7a–c (*Scylaticus*); Williston, 1891: 72 (cat.; *Scylaticus*); Kertész, 1909: 103 (cat.; *Scylaticus*); Stuardo Ortiz, 1946: 80 (cat.; *Pritchardia*); Carrera, 1953a: 276 (*Araiopogon*); Hull, 1962: 282 (*Araiopogon*); Carrera & Papavero, 1962: 60 (*Oberon*); Hull, 1962 (1): 145 (*Scylaticus*), 282 (*Araiopogon*); Artigas, 1970a: 152, figs. 133–134, 138, 469 (*Oberon*), 1971: 10, figs. 7, 9–10 (*Araiopogon*); Martin & Papavero, 1970: 37 (cat.; *Oberon*). Syntypes, MNHNP.

fulvipes (Philippi), 1865: 706 (*Deromyia*). Type-locality: Chile, Santiago. Neotype-locality: Chile, Santiago, Peñalolén. Distr. – Chile (Biobío, Maule, Santiago, Valparaíso). Refs. – Stuardo Ortiz, 1946: 82 (cat.; *Deromyia*); Artigas, 1970a: 154, figs. 135–137, 139–141, 435 (*Oberon*); Martin & Papavero, 1970: 37 (cat.; *Oberon*). Type, lost. NT DZUC [des. by Artigas, 1970a: 154].

gayi (Macquart), 1838: 37 (1839: 153) (*Dasypogon*). Type-locality: “Chile”. Distr. – Chile (Aconcagua, Arauco, Atacama, Coquimbo, O’Higgins, Santiago, Talca, Valparaíso). Refs. – Blanchard, 1852: 365, pl. 3, figs. 1a–d (*Dasypogon*); Walker, 1854: 434 (*Dasypogon*); Philippi, 1865: 685 (*Dasypogon*); Williston, 1891: 74 (cat.); Kertész, 1909: 138 (cat.); Brèthes, 1918: 49 (*Dasypogon*); Stuardo Ortiz, 1946: 82 (cat.); Carrera, 1949: 123 (*Araiopogon*), 1953a: 275 (*Araiopogon*); Hull, 1962 (1): 282, figs. 555, 1157, 1166, 1762, 1853, 1858 (*Araiopogon*); Martin & Papavero, 1970: 36 (cat.; *Araiopogon*); Knutson, 1971: 248 (*Trachysphyrus nigricornis* (Brûlé) (Hymenoptera: Ichneumonidae) as prey). Syntypes, MNHNP.

chalybeiventris Loew, 1851: 5 (*Dasypogon (Saropogon)*). Type-locality: Chile”. Ref. – Carrera, 1949: 123 (*Araiopogon*, syn.). Type, BZM [no. 434; cf. Schumann, 1973: 96].

hyacinthinus Bigot, 1878: 413. Type-locality: “Chile”. Refs. – Williston, 1891: 74 (cat.; *Sarapogon*); Kertész, 1909: 138 (cat.); Bromley, 1932: 266; Stuardo Ortiz, 1946: 82 (cat.); Hull, 1962 (1): 278. Type, OXF.

hypomelas (Loew), 1866: 24 (Cent. 7, no. 42) (*Diogmites*). Type-locality: U. S. A., ?New Mexico. Distr. – U. S. A. (New Mexico, Texas), Mexico (Nuevo León). Refs. – Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 116 (cat.; *Deromyia*); Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 130 (in key); Martin & Papavero, 1970: 35 (cat.). Type, MCZ.

melisoma (Carrera & Papavero), 1962: 60 (*Araiopogon*). Type-locality: Argentina, Tucumán. Distr. – Argentina (Tucumán). Ref. – Martin & Papavero, 1970: 36 (cat.; *Araiopogon*). Type, MZUSP.

mellipes Bromley, 1934b: 336. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Distr. – Guyana. Refs. – Carrera, 1949: 125 (says it may belong in *Cleptomyia*); Martin & Papavero, 1970: 36 (cat.). Type, AMNH.

nigrinasutus Bigot, 1878: 414. Type-locality: “Chile”. Distr. – Chile. Refs. – Williston, 1891: 74 (cat.; *Sarapogon*); Kertész, 1909: 139 (cat.); Bromley, 1932: 266; Stuardo Ortiz, 1946: 82 (cat.); Hull, 1962 (1): 278; Artigas, 1970a: 168, figs. 181–182; Martin & Papavero, 1970: 36 (cat.). Type, OX.

nitidus Wilcox, 1966a: 130 (in key), 135. Type-locality: U. S. A., Texas, Brewster County, Lajitas. Distr. – U. S. A. (Texas), Mexico (Coahuila). Ref. – Martin & Papavero, 1970: 36 (cat.). Type, CAS.

perniger Schiner, 1868: 166. Type-locality: “Chile”. Distr. – Chile. Refs. – Williston, 1891: 74 (cat.); Kertész, 1909: 139 (cat.); Stuardo Ortiz, 1946: 82 (cat.); Hull, 1962 (1): 278; Artigas, 1970a: 76 (*Araiopogon*); Martin & Papavero, 1970: 36 (cat.; *Araiopogon*). Type, NMW.

pulcherimus Williston, 1901: 312, pl. 6, fig. 3 (as *pulcherrima*, error). Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 263 (cat.); Kertész, 1909: 139 (cat.); Curran, 1930b: 2 (in key; as *pulcherrima*, error), 1931b: 1 (in key); Hull, 1962 (1): 278; Martin & Papavero, 1970: 36 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Syntypes, BMNH, AMNH.

purus Curran, 1930b: 3. Type-locality: U. S. A., Arizona, Baboquivari Mountains, Kits Peak Rincon. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Curran, 1931b: 1, 2 (in key); Hull, 1962 (1): 277; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 129, 130 (in key); Martin & Papavero, 1970: 36 (cat.). Type, AMNH.

semiustus Coquillett, 1904b: 186. Type-locality: U. S. A., California. Distr. – U. S. A. (California), Mexico (Sonora). Refs. – Kertész, 1909: 140 (cat.); Curran, 1930b: 3 (in key), 1931b: 2 (in key); Hull, 1962 (1): 278; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 130 (in key); Martin & Papavero, 1970: 36 (cat.). Type, USNM.

senex Osten Sacken, 1887: 179. Type-locality: Mexico, Sinaloa, Presio, near Mazatlán. Distr. – U. S. A., Mexico (Sinaloa, Durango, Jalisco, Sonora). Refs. – Aldrich, 1905: 263 (cat.); Kertész, 1909: 140 (cat.); Hull, 1962 (1): 278; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 128 (in key); Martin & Papavero, 1970: 36 (cat.); Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.

aridus Curran, 1930b: 3. Type-locality: U. S. A., Arizona, Sierras, Back Dike Prospect. Type, AMNH.

solutus Bromley, 1951: 15. Type-locality: U. S. A., Texas, Hidalgo County. Distr. – U. S. A. (Texas). Refs. – Hull, 1962 (1): 278; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1966a: 128 (in key). Type, USNM.

velutinus (Carrera & Papavero), 1962: 58, figs. 3-5 (*Oberon*). Type-locality: Argentina, Misiones, Loreto. Distr. – Argentina (Misiones). Type, MZUP.

wilcoxi Papavero, 1971b: 161, figs. 1-3. Type-locality: Mexico, Morelos, Cuernavaca. Distr. – Mexico (Morelos). Type, CAS.

Genus **Taracticus** Loew, 1872

Taracticus Loew, 1872a: 64 (Cent. 10, no. 24). Type-species, *Dioctria octopunctata* Say, 1823 (mon.). Ref. – Artigas & Papavero, 1988: 239, fig. 104, 250, figs. 151-156.

Dioctrodes Coquillett, 1904b: 181. Type-species, *Dioctrodes flavipes* Coquillett, 1904 (orig. des.) = *Taracticus octopunctatus* (Say, 1823).

Ceraturgus Wiedemann of authors, misident.

Two species known as fossils: *Taracticus contusus* Cockerell (1910: 282) and *Taracticus renovatus* Cockerell (1911: 80) (Oligocene; U. S. A., Colorado).

aciculatus Pritchard, 1938b: 179 (in key), 180. Type-locality: Mexico, Guerrero, Chilpancingo. Distr. – Mexico (Guerrero, Jalisco). Refs. – Hull, 1962 (1): 272; Martin & Papavero, 1970: 35 (cat.). Type, MINN.

argentifacies James, 1953: 55. Type-locality: Honduras, Morazán. Distr. – Honduras. Refs. – Hull, 1962 (1): 272; Martin & Papavero, 1970: 35 (cat.). Type, MICH.

dimidiatus (Macquart), 1847: 51 (1847: 35) (*Dasyptogon*). Type-locality: “Mexico”. Distr. – Mexico (Veracruz). Refs. – Walker, 1847: 428 (*Dasyptogon*); Bellardi, 1861: 61 (*Ceraturgus*); Osten Sacken, 1874: 177 (*Ceraturgus*); Aldrich, 1905: 258 (cat.; *Ceraturgus*); Kertész, 1909: 97 (cat.; *Ceraturgus*); Hull, 1962 (1): 169 (?*Ceraturgus*); Martin & Papavero, 1970: 35 (cat.); Papavero & Ibáñez Bernal, 2001: 85 (hist. Mexican dipterol.). Type, OXF.

geniculatus (Bigot), 1878: 443 (*Ceraturgus*). Type-locality: “Mexico”. Refs. – Williston, 1883: 33 (*Ceraturgus*); Aldrich, 1905: 258 (cat.; *Ceraturgus*); Kertész, 1909: 97 (cat.; *Ceraturgus*); Hull, 1962 (1): 169; Martin & Papavero, 1970: 35 (cat.); Papavero & Ibáñez Bernal, 2001: 122 (hist. Mexican dipterol.). Type, OXF.

guerrerensis Pritchard, 1938b: 179 (in key), 184. Type-locality: Mexico, Guerrero, Chilpancingo. Distr. – Mexico (Guerrero). Refs. – Hull, 1962 (1): 272; Martin & Papavero, 1970: 35 (cat.). Type, MINN.

nigrimystaceus Williston, 1901: 313, pl. 5, fig. 23. Type-locality: Mexico, Guerrero, Zucumanatlán, 7000 ft. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 266 (cat.); Kertész, 1909: 143 (cat.); Hull, 1962 (1): 272; Martin & Papavero, 1970: 35 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol.). Type, BMNH.

nigripes Williston, 1901: 313. Type-locality: Mexico, Guerrero, Xucumatlán, Omiltemi and Chilpancingo. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 258 (cat.); Kertész, 1909: 97 (cat.); Pritchard, 1938b: 180 (in key), 185; Hull, 1962 (1): 272; Martin & Papavero, 1970: 35 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol.). Syntypes, BMNH, SMTD [cf. Kallweit, 1992: 196].

rufipennis (Macquart), 1847: 48 (1847: 32) (*Ceraturgus*). Type-locality: “Mexico”. Refs. – Walker, 1854: 378 (*Ceraturgus*); Bellardi, 1861: 59 (*Ceraturgus*); Osten Sacken, 1874: 177 (*Ceraturgus*); Aldrich, 1905: 258 (cat.; *Ceraturegus*); Kertész, 1909: 97 (cat.; *Ceraturgus*); Hull, 1962 (1): 169; Martin & Papavero, 1970: 35 (cat.; n. comb.); Papavero & Ibáñez Bernal, 2001: 85 (hist. Mexican dipterol.). Type, OXF.

similis Williston, 1901: 312. Type-locality: Mexico, Guerrero, Omiltemi and Sierra de las Aguas Escondidas, 5000-6000 ft. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 266 (cat.); Kertész, 1909: 143 (cat.); Pritchard, 1938b: 180 (in key), 187; Hull, 1962 (1): 272; Martin & Papavero, 1970: 35 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol.). Type, BMNH.

vitripennis (Bellardi), 1861: 160 (60) (*Ceraturgus*). Type-locality: Mexico, Morelos, Cuautla. Distr. – Mexico (Jalisco, Guerrero, Chiapas). Refs. – Osten Sacken, 1874: 177 (*Ceraturgus*); Snow, 1896: 179; Williston, 1901: 313; Aldrich, 1905: 266 (cat.); Kertész, 1909: 143 (cat.); Pritchard, 1938b: 179 (in key), 181; Hull, 1962 (1): 272; Martin & Papavero, 1970: 35 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol.). Type, TORO.

Genus **Tocantinia** Carrera, 1955

Tocantinia Carrera, 1955a: 102. Type-species, *Dasyptogon miser* Walker, 1854 (orig. des.).

misera (Walker), 1854: 433 (*Dasyptogon*). Type-locality: Brazil, Pará, rio Tapajós. Distr. – Brazil (Pará). Refs. – Williston, 1891: 76 (cat.; *Dasyptogon*); Kertész, 1909: 230 (cat.; *Dasyptogon*); Bromley, 1946: 107 (cat.; *Dasyptogon*); Carrera, 1955a: 103 (n. comb.); Hull, 1962 (1): 281, figs. 564, 1128, 1137; Martin & Papavero, 1970: 37 (cat.); Artigas & Papavero, 1988b: 239, fig. 101. Type, BMNH.

Tribe Isopogonini G. H. Hardy, 1948

Isopogonini G. H. Hardy, 1948: 118. Type-genus, *Isopogon* Loew, 1847 = *Leptarthrus* Stephens, 1829. Ref. – Dikow, 2003: 463.

Genus **Alvarenga** Carrera, 1960

Alvarenga Carrera, 1960a: 159. Type-species, *Alvarenga icarius* Carrera, 1960 (orig. des.).

icarius Carrera, 1960a: 160, figs. 7-10. Type-locality: Brazil, Paraíba, Soledade, Juazeirinho. Distr. – Brazil (Ceará, Paraíba). Refs. – Hull, 1962 (2): 596; Martin & Papavero, 1970: 33 (cat.); Artigas & Papavero, 1988b: 221, figs. 1-9. Type, MNRJ.

matilei Papavero, 1971d: 24. Type-locality: Argentina, Chaco de Santiago del Estero, margins of the rio Salado, near Icaño. Distr. – Argentina (Santiago del Estero). Ref. – Artigas & Papavero, 1988b: 222, figs. 10-13. Type, MNHNP.

Genus **Annamyia** Pritchard, 1941

Annamyia Pritchard, 1941a: 131. Type-species, *Annamyia maren* Pritchard, 1941 (orig. des.).

maren Pritchard, 1941a: 132, fig. 1. Type-locality: Brazil, Minas Gerais, Diamantina. Distr. – Brazil (Minas Gerais). Refs. – Carrera, 1949: 131; Hull, 1962 (1): 266, figs. 172, 562, 1146, 1155, 1582, 1583, 1700, 1836; Martin & Papavero, 1970: 33 (cat.). Type, USNM.

Genus **Aphamartania** Schiner, 1866

Aphamartania Schiner, 1866: 671. Type-species, *Aphamartania frauenfeldi* Schiner, 1866 (orig. des.).

breviventris (Macquart), 1848: 181 (1848: 21), pl. 1, fig. 13 (*Dasypogon*). Type-locality: “Rio Negro”. Refs. – Walker, 1854: 435 (*Dasypogon*); Williston, 1891: 73 (*Theromyia*); Hull, 1962 (1): 263; Papavero, 1971: 21; Martin & Papavero, 1970: 32 (cat.). Type, OXF.

beviventris Kertész, 1909: 143 (cat.), error.

rufiventris Bromley, 1946: 108, error.

digna Pritchard, 1941a: 137, fig. 2. Type-locality: Peru, Huánuco, Shishmay. Distr. – Peru (Huánuco). Refs. – Carrera, 1949: 128; Hull, 1962 (1): 263; Martin & Papavero, 1970: 32 (cat.). Type, USNM.

flavipennis (Macquart), 1846: 192 (1846: 64), pl. 7, fig. 4 (*Dasypogon*). Type-locality: “Colombia” (as “Nouvelle Grenade”). Ref. – Papavero, 1971d: 21 (n. comb.). Type, OXF.

frauenfeldi Schiner, 1866: 671 (only name cited, but species not described), 1868: 372 (descr.). Type-locality: “Venezuela”. Distr. – Panama, Venezuela, ?Bolivia. Refs. – Williston, 1891: 73 (cat.); Kertész, 1909: 144 (cat.); Engel, 1930: 468; Pritchard, 1941a: 136; Carrera, 1949: 127; Hull, 1962 (1): 263, 183, 554, 1143, 1152; Martin & Papavero, 1970: 32 (cat.). Type, NMW.

panamensis Curran, 1931b: 5 (in key), 6 (*Cophura*). Type-locality: Panama, Canal Zoner, Barro Colorado I. Type, AMNH.

knutsoni Papavero, 197d: 53, figs. 1-3. Type-locality: Panama, Potrerillos. Distr. – Panama. Type, USNM.

maculipennis (Macquart), 1838: 37 (1839: 133), pl. 3, fig. 8 (*Dasypogon*). Type-locality: Brazil, western Minas Gerais. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo, Paraná), Argentina (Misiones). Refs. – Walker, 1854: 436 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 129 (cat.; *Dasypogon*); Carrera, 1945b: 170; Bromley, 1946: 107 (cat.; *Dasypogon*); Carrera, 1949: 128; Hull, 1962 (1): 228 (*Dasypogon*), 263, figs. 174, 551, 1145, 1154, 1856, 1951 [*Aphamartania*]; Martin & Papavero, 1970: 32 (cat.). Type, MNHNP.

pritchardi Carrera, 1943: 120, pl. 10. Type-locality: Brazil, Paraná, Curitiba. Refs. – Carrera, 1949: 129, figs. 64, 93, 153, 158, 183, 1960: 161; Carrera & Papavero, 1962: 62; Hull, 1962 (1): 263; Martin & Papavero, 1970: 32 (cat.); Papavero, 1971d: 23 (n. syn.); Martin & Papavero, 1988: 224, fig. 20. Type, MZUSP.

marga Pritchard, 1941a: 138, fig. 4. Type-locality: Argentina, Tucumán. Distr. – Argentina (Tucumán). Refs. – Carrera, 1949: 130; Carrera & Papavero, 1962: 62; Hull, 1962 (1): 263; Martin & Papavero, 1970: 32 (cat.). Type, USNM.

Genus *Aspidopyga* Carrera, 1949

Aspidopyga Carrera, 1949: 132. Type-species, *Aspidopyga cophurooides* Carrera, 1949 (orig. des.).

cophurooides Carrera, 1949: 133, figs. 4, 65, 92, 121, 127-131, 182. Type-locality: Brazil, Rio de Janeiro, Palmeiras. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Carrera, 1960a: 161; Carrera & Papavero, 1962: 62; Hull, 1962 (1): 268, figs. 479, 1141, 1150, 1725, 1729; Martin & Papavero, 1970: 33 (cat.); Artigas & Papavero, 1988b: 223, figs. 14-19, 224, fig. 21. Type, IOC.

Genus *Cophura* Osten Sacken, 1887

Blax Loew, 1872a: 65 (Cent. 10, no. 24) (preocc. Koch, 1840). Type-species, *Blax bellus* Loew, 1872 (mon.).

Blacodes Loew, 1874a: 377 (*nom. nov.* for *Blax* Loew, 1872, but preocc. Dejean, 1859). Type-species, *Blax bellus* Loew, 1872 (aut.).

Cophura Osten Sacken, 1887: 181. Type-species, *Cophura sodalis* Osten Sacken, 1887 (mon.).

Loewiella Williston, 1896b: 57 (*nom. nov.* for *Blacodes* Loew, 1874, but preocc. Meunier, 1894). Type-species, *Blax bellus* Loew, 1872 (aut.).

Buckellia Curran, 1925b: 156. Type-species, *Cophura albosetosa* Hine, 1908 (orig. des.).

One species known as fossil: *Cophura antiquella* Cockerell (1913: 213, Oligocene; U. S. A., Colorado).

acapulcae Pritchard, 1943: 285 (in key), 302. Type-locality: Mexico, Guerrero, Acapulco. Distr. – Mexico (Guerrero). Refs. – Hull, 1962 (1): 270, fig. 529; Martin & Papavero, 1970: 33 (cat.). Type, MINN.

apotma Pritchard, 1943: 284 (in key), 301. Type-locality: Mexico, Puebla, Tehuacán. Distr. – Mexico (Puebla). Refs. – Martin & Papavero, 1970: 33 (cat.). Type, MINN.

apatma Hull, 1962 (1): 270, error.

arizonensis (Schaeffer), 1916: 65 (*Lasiocnemus*). Type-locality: U. S. A., Arizona. Distr. – U. S. A. (Arizona, New Mexico), Mexico (Sonora). Refs. – Pritchard, 1943: 284 (in key), 299; Hull, 1962 (1): 270; Martin & Wilcox, 1965: 364 (cat.); Wilcox, 1965b: 809 (in key), fig. 1; Artigas & Papavero, 1988b: 226, figs. 29-33, 227, fig. 34.

drakei Pritchard, 1935: 6 (*Buckellia*). Type-locality: U. S. A., Arizona.

atypha Pritchard, 1943: 284 (in key), 301. Type-locality: Mexico, Puebla, Tehuacán. Distr. – Mexico (Guerrero, Puebla). Refs. – Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). Type, MINN.

bella (Loew), 1872a: 63 (Cent. 10, no. 24) (*Blax*). Type-locality: U. S. A., Texas. Distr. – U. S. A. (Arizona, New Mexico, Texas), Mexico (Jalisco, Nayarit, Sinaloa). Refs. – Coquillett, 1893b: 33 (*Blacodes*); Aldrich, 1905: 266 (cat.); Back, 1909: 379; Curran, 1930b: 5 (in key); Bromley, 1934a: 102; Pritchard, 1943: 285 (in key), 291, fig. 1; Hull, 1962 (1): 270; Martin & Wilcox, 1965: 367 (cat.); Wilcox, 1965b: 807 (in key), fig. 1; Martin & Papavero, 1970: 34 (cat.); Artigas & Papavero, 1988: 227, fig. 35. Type, MCZ.

calla Pritchard, 1943: 285 (in key), 303. Type-locality: Mexico, Guerrero, Iguala. Distr. – Mexico (Guerrero, Puebla). Refs. – Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). Type, MINN.

cora Pritchard, 1943: 284 (in key), 299. Type-locality: Mexico, Puebla, Tehuacán. Distr. – Mexico (Puebla). Refs. – Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). Type, MINN.

dammersi Wilcox, 1965b: 807, fig. 1. Type-locality: U. S. A., California, Imperial County, 25 miles south of Oasis. Distr. – U. S. A. (California), Mexico (Baja California). Type, CAS.

daphne Pritchard, 1943: 283 (in key), 290. Type-locality: Mexico, Morelos, Cuernavaca. Distr. – U. S. A. (Arizona), Mexico (Morelos). Refs. – Hull, 1962 (1): 270; Martin & Wilcox, 1965: 367 (cat.); Martin & Papavero, 1970: 34 (cat.). Type, USNM.

feigei Kaletta, 1983: 302. Type-locality: Venezuela, Miranda, Los Moralistos. Distr. – Venezuela (Miranda). Type, IZAUCV.

gellergrimmi Papavero, *nom. nov.* for *wilcoxi* Kaletta, 1983. Type-locality: Venezuela, Guárico, San Francisco de Tiznados. Distr. – Venezuela (Guárico). [Dedicated to Dr. Fritz Geller-Grimm, Museum of Wiesbaden].

wilcoxi Kaletta, 1983: 301 (preocc. Pritchard, 1935). Type-locality: Venezuela, Guárico, San Francisco de Tiznados. Type, IZAUCV.

getzendaneri Wilcox, 1959: 123, fig. (p. 126), 127 (in key). Type-locality: U. S. A., California, Indio. Distr. – U. S. A. (California, Arizona), Mexico (Baja California). Refs. – Martin & Wilcox, 1965: 367 (cat.); Wilcox, 1965b: 806 (in key), fig. 1; Martin & Papavero, 1970: 34 (cat.). Type, CAS.

humilis (Bellardi), 1861: 177 (77) (?*Dasyopogon*). Type-locality: Mexico, Morelos, Cuautla. Distr. – Mexico (Guerrero, Jalisco). Refs. – Osten Sacken, 1874: 180 (*Anisopogon*); Aldrich, 1905: 263 (cat.; *Heteropogon*); Kertész, 1909: 65 (cat.; *Heteropogon*); Curran, 1930b: 5 (in key); Pritchard, 1943: 284 (in key), 289; Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.

sodalis Osten Sacken of Williston, 1901: 314, misident.

igualae Pritchard, 1943: 284 (in key), 300. Type-locality: Mexico, Guerrero, Iguala. Distr. – Mexico (Guerrero). Refs. – Hull,

- 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). Type, Pritchard Coll.
- incerta** (Becker), 1919: 165 (*Metapogon*). Type-locality: Ecuador, Casitagua, 3500 m; Loja, 2220 m; Cuenca, 2532 m. Lectotype-locality: Ecuador, Casitagua, 3500 m. Distr. – Ecuador. Refs. – Hull, 1962 (1): 203 (*Metapogon*); Martin & Papavero, 1970: 89 (*Metapogon*); Papavero, 1971: 23 (n. comb.). Lectotype MNHNP [des. by Papavero, 1971d: 23].
- sundra** Pritchard, 1943: 294 (in key), 295. Type-locality: Ecuador, Ponda. Refs. – Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). Type, MINN.
- nephressa** Pritchard, 1943: 285 (in key), 292. Type-locality: Mexico, Nuevo León, Galeana, 5000-6000 ft. Distr. – Mexico (Nuevo León). Refs. – Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). Type, MCZ.
- painteri** Pritchard, 1943: 285 (in key), 293. Type-locality: U. S. A., Arizona, Safford. Distr. – U. S. A. (California, Arizona), Mexico (Baja California, Sonora). Refs. – Hull, 1962 (1): 270; Martin & Wilcox, 1965: 367 (cat.); Wilcox, 1965b: 812, fig. 1. Type, Painter coll.
- picta** Carrera, 1955a: 106, fig. 2. Type-locality: “Ecuador”. Distr. – Ecuador. Refs. – Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). Type, BMNH.
- pollinosa** Curran, 1930a: 10. Type-locality: U. S. A., Arizona, Baboquivari Mountains, Kits Peak Rincon, 4050 ft. Distr. – U. S. A. (Arizona, Colorado, Utah), Mexico (Coahuila, Sonora). Refs. – Pritchard, 1935: 6 (in key), 7 (*Buckellia*), 1943: 285 (in key), 304; Hull, 1962 (1): 270, fig. 1981; Martin & Wilcox, 1965: 367 (cat.); Wilcox, 1965b: 809 (in key), fig. 1; Martin & Papavero, 1970: 34 (cat.); Nelson, 1987: 61. Type, AMNH.
- lutzi** Curran, 1931b: 5 (in key), 7. Type-locality: U. S. A., Colorado, Regnier. Refs. – Bromley, 1935: 3; Pritchard, 1935c: 6 (in key), 7 (*Buckellia*); Baker, 1939: 170 (*Buckellia*); Pritchard, 1943: 285 (in key), 305. Type, AMNH.
- wilcoxi** Pritchard, 1935: 8 (*Buckellia*). Type-locality: U. S. A., Arizona. Ref. – Martin & Wilcox, 1965: 367 (cat.; syn.).
- pulchella** Williston, 1901: 314, pl. 5, fig. 14. Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. – U. S. A. (Utah), Mexico (Guerrero). Refs. – Aldrich, 1905: 266 (cat.); Kertész, 1909: 142 (cat.); Curran, 1930b: 5 (in key); Pritchard, 1943: 284 (in key), 297; Hull, 1962 (1): 270; Martin & Wilcox, 1965: 367 (cat.); Martin & Papavero, 1970: 34 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Syntypes, AMNH.
- sodalis** Osten Sacken, 1887: 181, pl. 3, fig. 13. Type-locality: Mexico, Sinaloa, Presio, near Mazatlán. Distr. – Mexico (Sinaloa, Sonora). Refs. – Williston, 1901: 314; Aldrich, 1905: 266 (cat.); Kertész, 1909: 142 (cat.); Curran, 1930b: 5 (in key); Pritchard, 1943: 283 (in key), 289, figs. 3, 5; Hull, 1962 (1): 270, figs. 149, 1122, 1131, 1821, 1854; Martin & Wilcox, 1965: 367 (cat.); Wilcox, 1965b: 807; Martin & Papavero, 1970: 34 (cat.); Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.
- tanbarki** Wilcox, 1965b: 811. Type-locality: U. S. A., California, Los Angeles County, Tanbark Flat. Distr. – U. S. A. (California), Mexico (Baja California). Type, CAS.
- tolandi** Wilcox, 1959: 125, fig. (p. 126), 127 (in key). Type-locality: U. S. A., California, Riverside County, White Water. Distr. – U. S. A. (California, Arizona), Mexico (Baja California). Ref. – Martin & Wilcox, 1965: 367 (cat.). Type, CAS.
- willistoni** Pritchard, 1943: 284 (in key), 290 (*nom. nov.* for *humilis* Williston, 1901). Type-locality: Mexico, Guerrero, Tepetlapa. Distr. – Mexico (Guerrero, Veracruz). Refs. – Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.). g
- humilis** Williston, 1901: 315 (preocc. Bellardi, 1861). Type-locality: Mexico, Guerrero, Tepetlapa. Refs. – Aldrich, 1905: 266 (cat.); Kertész, 1909: 142 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Type, BMNH.
- zandra** Pritchard, 1943: 284 (in key), 294. Type-locality: Peru, Cajamarca, Celendia. Distr. – Peru (Cajamarca), Argentina. Refs. – Carrera & Papavero, 1962: 63; Hull, 1962 (1): 270; Hull, 1962 (1): 270; Martin & Papavero, 1970: 34 (cat.); Lamas M., 1972: 308. Type, MINN.

Genus *Hodophylax* James, 1933

Hodophylax James, 1933: 1. Type-locality: *Hodophylax aridus* James, 1933 (orig. des.). Ref. – Artigas & Papavero, 1988b: 230, fig. 49.

- basingeri** Pritchard, 1938c: 130, fig. Type-locality: U. S. A., California, San Bernardino County, Quail Spring. Distr. – U. S. A. (California), Mexico (Baja California, Sonora). Refs. – Wilcox, 1961: 113 (in key and text); Hull, 1962 (1): 267; Martin & Wilcox, 1965: 374 (cat.); Martin & Papavero, 1970: 33 (cat.); Artigas & Papavero, 1988b: 228, figs. 37-43. Type, Basinger Coll.
- tolandi** Wilcox, 1961: 113 (in key), 114. Type-locality: U. S. A., Arizona, Shumway. Distr. – U. S. A. (Arizona), Mexico (Chihuahua, Nuevo León, Zacatecas). Refs. – Martin & Wilcox, 1965: 374 (cat.); Martin & Papavero, 1970: 33 (cat.). Type, CAS.

Genus *Nicocles* Jaennicke, 1867

Pygostylus Loew, 1866: 16 (Cent. 7, no. 28) (preocc. Haliday, 1833). Type-species, *Pygostolus argentifer* Loew, 1866 (sub.

des., Coquillett, 1910: 598) = *Nicocles politus* (Say, 1823).
Nicocles Jaennicke, 1867: 355. Type-species, *Nicocles analis* Jaennicke, 1867 (mon.).

Ref. – Artigas & Papavero, 1988b: 229, figs. 44–48, 230, fig. 50.

One species known as fossil: *Nicocles miocenicus* Cockerell (1909: 155) (Oligocene; U. S. A., Colorado).

analis Jaennicke, 1867: 355, pl. 43, fig. 13. Type-locality: “Mexico”. Distr. – Mexico (Distrito Federal, Durango, Jalisco, Hidalgo, Oaxaca, Puebla). Refs. – Aldrich, 1905: 267 (cat.); Kertész, 1909: 141 (cat.); Hull, 1962 (1): 259; Martin & Papavero, 1970: 31 (cat.); Papavero & Ibáñez Bernal, 2001: 114 (hist. Mexican dipterol).

bromleyi D. E. Hardy, 1943: 28, figs. 4a-d. Type-locality: U. S. A., Arizona, Huachuca Mountains, Sunnyside Canyon. Distr. – U. S. A. (Arizona), Mexico (Aguascalientes, Chihuahua, Durango, Jalisco, Zacatecas). Refs. – Hull, 1962 (1): 259; Martin & Wilcox, 1965: 380 (cat.); Martin & Papavero, 1970: 31 (cat.). Type, KU.

Genus **Theromyia** Williston, 1891

Cylindrophora Philippi, 1865: 704 (preocc. Solier, 1849). Type-species, *Cylindrophora murina* Philippi, 1865 (mon.).
Lynchia Williston, 1889: 225 (*nom. nov.* for *Cylindrophora* Philippi, 1865, but preocc. Weyenbergh, 1882). Type-species, *Cylindrophora murina* Philippi, 1865 aut.).

Myiothera Williston, 1889: 229 (*nom. nov.* for *Lynchia* Williston, 1889, but preocc. Illiger, 1811). Type-species, *Cylindrophora murina* Philippi, 1865 (aut.).

Theromyia Williston, 1891: 73 (cat.; *nom. nov.* for *Myiothera* Williston, 1889). Type-species, *Cylindrophora murina* Philippi, 1865 (aut.).

murina (Philippi), 1865: 704 (*Cylindrophora*). Type-locality: “Chile”. Neotype-locality: Chile, Valparaíso, Algarrobo. Distr. – Chile (Aconcagua, Arauco, Cautín, Concepción, Coquimbo, Curicó, Linares, Malleco, Maule, Santiago, Valparaíso). Refs. – Williston, 1889: 255 (*Lynchia*), 159 (*Myoothera*), 1891: 73 (*Theromyia*); Kertész, 1909: 144 (*Aphamartania*); Bromley, 1932: 266 (*Aphamartania*); Pritchard, 1941a: 135, fig. 3 (*Aphamartania*); Stuardo Ortiz, 1946: 83 (cat.; *Aphamartania*); Carrera, 1949: 130, figs. 98, 135–136, 1953a: 276; Hull, 1962 (1): 262, figs. 170, 1142, 1151, 1874, 1908; Artigas, 1970a: 184, figs. 184–191, 462; Martin & Papavero, 1970: 32 (cat.); Artigas, 1971: 13, figs. 21–22. Type, lost. NT DZUC [des. by Artigas, 1970a: 184].

calopyga Schiner, 1868: 166 (*Cylindrophora*). Type-locality: “Chile”. Refs. – Williston, 1891: 73 (*Theromyia*); Kertész, 1909: 144 (cat.; *Aphamartania*). Type, lost.

nana (Pritchard), 1941a: 135 (*Aphamartania*). Type-locality: Peru, Lima, Verrugas Canyon, 7000 ft. Distr. – Peru (Huánuco, Lima). Refs. – Hull, 1962 (1): 263; Martin & Papavero, 1970: 32 (cat.); Lamas M., 1972: 308. Type, USNM.

pegnai Artigas, 1970a: 187, figs. 192–196, 443. Type-locality: Chile, Colchagua, Lolol. Distr. – Chile (Colchagua, Santiago). Refs. – Artigas, 1971: 14, figs. 19–20; Artigas & Papavero, 1988: 231, figs. 52–57. Type, DZUC.

Tribe **Lastaurini** Papavero, 1973

Lastaurini Papavero, 1973b: 277. Type-genus, *Lastaurus* Loew, 1851.
Blepharepiini Papavero, 1973b: 276, 277. Type-genus, *Blepharepium* Rondani, 1848.
Lastauracini Sabrosky, 1999: 172, error.

Genus **Apolastaurooides** Artigas & Papavero, 1988

Apolastaurooides Artigas & Papavero, 1988b: 208. Type-species, *Apolastaurooides kamakusa* Papavero & Artigas, 1988 (orig. des.).

kamakusa Papavero & Artigas, 1988b: 208. Type-locality: Guyana, Mazaruni, Kamakusa. Distr. – Guyana. Type, MZUSP.

Genus **Blepharepium** Rondani, 1848

Blepharepium Rondani, 1848: 89. Type-species, *Blepharepium luridum* Rondani (mon.).

Senobasis Macquart of aurthors, part, misident.

Planetolestes Lynch Arribálzaga, 1879: 147. Type-species, *Laphria coarctata* Perty, 1833 (orig. des.).

Ref. – Papavero & Bernardi, 1973: 168 (key to spp.).

One species known as fossil: *Blepharepium borealis* (James) (1939: 45; as *Senobasis*) (Oligocene; U. S. A., Colorado).

annulatum (Bigot), 1857a: 331, pl. 20, figs. 3, 3a (*Senobasis*). Type-locality: “Cuba”. Distr. – Mexico, Guatemala, Honduras, El Salvador, Costa Rica, Belize, Cuba, Panama, Trinidad, St. Eustatius, Aruba, Bonaire, Colombia and Venezuela.

Refs. – Osten Sacken, 1874: 184 (*Diogmites*), 1878: 72, 233 (cat.; *Diogmites*); Williston, 1891: 75 (cat.; *Diogmites*); Kertész, 1909: 124 (cat.; *Stenobasis*, as syn. of *coarctatum* Perty, 1833); Bromley, 1929a: 279, pl. 1, fig. 21; Curran, 1942: 55; Carrera, 1949: 46 (note); Hull, 1962 (1): 233; Papavero & Bernardi, 1973: 172, figs. 4-6, map 3. Type, MHNTP.

secabilis Walker, 1860: 276 (*Dasypogon*). Type-locality: “Mexico”. Refs. – Bellardi, 1861: 63, pl. 1, fig. 4 (?*Saropogon*); Schiner, 1867: 371 (*Stenobasis*); Lynch Arribálzaga, 1879: 150 (*Planetolestes*; note), 1881: 24; Osten Sacken, 1874:

184 (*Diogmites*, as syn. of *annulatus* Bigot, 1857), 1887: 171 (*Planetolestes*); Williston, 1891: 76 (cat.); Kertész, 1909: 124 (cat.; *Stenobasis*, as syn. of *coarctatum* Perty, 1833); Bromley, 1929: 279 (in key); Curran, 1942: 55;

Carrera, 1949: 53 (part), 1953a: 272; Carrera & d’Andretta, 1961: 68 (prey); Carrera & Papavero, 1962: 47; Hull, 1962

(1): 233; Carrera & Machado-Allison, 1963: 237; Martin, 1965a: 20; Wilcox & Martin, 1965: 265 (cat.); Martin &

Papavero, 1970: 28 (cat.); Papavero & Bernardi, 1973: 172 (syn.); Alayo Dalmau & Garcia Ávila, 1983: 47 (cat.; *Senobasis*); Papavero & Ibáñez Bernal, 2001: 95 (hist. Mexican dipterol.). Type, BMNH.

cajennense (Fabricius), 1787: 360 (*Asilus*). Type-locality: French Guiana, Cayenne (Rohr) [cf. Papavero, 1971e: 20]. Neotype-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Distr. – Colombia, Venezuela, Trinidad, Guyana, Surinam, Peru, Bolivia, Brazil (Acre, Amazonas). Refs. – Olivier, 1789: 269 (*Asilus*); Fabricius, 1794: 386 (*Asilus*), 1805: 162 (*Laphria*); Wiedemann, 1821: 220 (*Laphria*), 1828: 383 (*Dasypogon*); Walker, 1854: 442 (*Dasypogon*); Schiner, 1866: 677 (*Dasypogon*); Williston, 1891: 67 (cat.; *Dasypogon*); Kertész, 1909: 126 (cat.; *Dasypogon*); Curran, 1934c: 334; Hull, 1962 (1): 233; Martin & Papavero, 1970: 27 (cat.); Lamas M., 1972: 307 (cat.); Papavero & Bernardi, 1973: 178, figs. 1-17, map 1; Fisher, 1984: 27; Artigas & Papavero, 1988b: 260, fig. 216. Type, KIEL (only indeterminable remains; cf. Zimsen, 1964: 469). NT MZUSP [des. by Papavero & Bernardi, 1973: 178, 179].

cayennensis Gmelin, 1792: 162 (*Asilus*), error. Refs. – Bromley, 1929a: 279 (in key); Curran., 1934c: 334, 1942: 54; Carrera, 1949: 50; Carrera & Machado-Allison, 1963: 237.

inserens Walker, 1851: 91. Type-locality: unknown. Ref. – Papavero & Bernardi, 1973: 178 (syn.). Type, BMNH.

auricincta Schiner, 1867: 371 (*Senobasis*). Type-locality: “Surinam”. Refs. – Osten Sacken, 1874: 184 (*Diogmites*; as syn. of *auricinctus* Bigot, 1857); Lynch Arribálzaga, 1881: 26 (n. comb.); Williston, 1891: 75 (cat.); Kertész, 1909: 124 (cat.; *Stenobasis*, as syn. of *coarctatum* Perty, 1833); Curran, 1942: 55; Hull, 1962 (1): 233 (cat.); Martin & Papavero, 1970: 27 (cat.); Papavero & Bernardi, 1973: 178 (syn.). Type, NMW.

inca Curran, 1942: 54. Type-locality: Peru, Loreto, Iquitos. Refs. – Carrera, 1949: 57; Hull, 1962 (1): 233; Martin & Papavero, 1970: 28 (cat.); Papavero & Bernardi, 1970: 178 (syn.). Type, AMNH.

vorax Curran, 1942: 53. Type-locality: Peru, La Merced. Refs. – Carrera, 1949: 56; Carrera & d’Andretta, 1953: 65; Carrera & Papavero, 1962: 48; Hull, 1962 (1): 233; Martin & Papavero, 1970: 28 (cat.); Papavero & Bernardi, 1973: 178 (syn.). Type, AMNH.

ssp. **coarctatum** (Perty), 1833: 181, pl. 36, fig. 4 (*Laphria*). Type-locality: Brazil, Amazonas, rio Negro. Distr. – Peru, Bolivia, Brazil (Pernambuco, Mato Grosso, Goiás, Distrito Federal, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Rio Grande do Sul), Paraguay, Uruguay, Argentina (Santiago del Estero, Chaco, Entre Ríos, Santa Fe, Misiones, Formosa, Córdoba, Mendoza, Tucumán, Buenos Aires). Refs. – Walker, 1854: 504 (*Dasypogon*); Lynch Arribálzaga, 1879: 150 (*Planetolestes*), 1881: 24 (n. comb.); Wulp, 1882b: 88; Lynch Arribálzaga, 1883a: 139 (cat.); Williston, 1891: 75 (cat.); Kertész, 1909: 124 (cat.; *Stenobasis*); Snow, 1896: 177; Williston, 1901: 310; Aldrich, 1905: 236 (cat.); Kertész, 1909: 124 (cat.; *Stenobasis*); Bromley, 1919: 279 (in key); Lindner, 1929: 170 (biol. notes); Engel, 1930: 469; Curran, 1942: 55; Bromley, 1946: 107 (cat.); Carrera, 1947a: 43, 1949: 55, figs. 15, 70, 118; Carrera & d’Andretta, 1961: 68 (prey); Carrera & Papavero, 1962: 47 (part); Hull, 1962 (1): 233, figs. 543, 547, 1067, 1076, 1920; Martin, 1965a: 19; Papavero & Bernardi, 1973: 182, figs. 21-28, map 1 (n. stat.). Type, ZSBS.

bonariensis Macquart, 1838: 194 (1839: 310) (*Dasypogon*). Type-locality: Argentina, Buenos Aires. Refs. – Walker, 1854: 439 (*Dasypogon*); Lynch Arribálzaga, 1879: 150 (n. comb.; as syn. of *coarctatum* Perty, 1833); Kertész, 1909: 124 (cat.; *Stenobasis*, as syn. of *coarctatum* Perty, 1833); Carrera, 1949: 47; Martin & Papavero, 1970: 27 (cat.); Papavero & Bernardi, 1973: 182, figs. 21-28 (n. status). Type, MHNTP.

occidens Walker, 1851: 91 (*Dasypogon*). Type-locality: unknown (“South America” in type label). Refs. – Walker, 1854: 485 (*Dasypogon*); Kertész, 1909: 139 (cat.; *Dasypogon*); Papavero & Bernardi, 1973: 182 (syn.). Type, BMNH.

lynchi Carrera, 1949: 51, figs. 71, 171. Type-locality: Argentina, Delta, Abra Vieja. Type, MZUSP. Refs. – Carrera & Papavero, 1962: 49; Hull, 1962 (1): 233; Martin & Papavero, 1970: 28 (cat.); Papavero & Bernardi, 1973: 182

- (syn.). Type, MZUSP.
- secabile* Walker of Carrera, 1949: 54, part, misident.
- ssp. *cunctabundum* Papavero & Bernardi, 1973: 185, figs. 18-20, map 1. Type-locality: Brazil, São Paulo, Santo André, Distr. – Brazil ((São Paulo, Paraná, Santa Catarina, Rio Grande do Sul). Type, MZUSP.
- fuscipennis** (Macquart), 1834: 294 (*Dasypogon*; preocc. Wiedemann, 1821). Type-locality: French Guiana, Cayenne. Distr. – Surinam, French Guiana, Peru, Brazil (Pará, Amapá, Rio Grande do Norte, Espírito Santo, Rio de Janeiro). Refs. – Walker, 1854: 439 (*Dasypogon*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 128 (cat.; *Dasypogon*); Carrera & d'Andretta, 1953: 66; Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 89 (cat.; *Dasypogon*); Papavero & Bernardi, 1973: 186, figs. 29-31, map 2 (n. comb.); Artigas & Papavero, 1988b: 252, figs. 168-172. Type, MHNTP.
- insertus* Walker, 1851: 94 (*Dasypogon*). Type-locality: Brazil, Pará (i. e., Pará, Belém). Refs. – Walker, 1854: 431 (*Dasypogon*); Williston, 1891: 68, 76 (cat.; *Dasypogon*); Kertész, 1909: 129 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Martin & Papavero, 1970: 89 (cat.); Papavero & Bernardi, 1973: 187 (syn.). Type, BMNH.
- maculipennis* Macquart, 1855: 71 (1855: 51) (*Senobasis*). Type-locality: unknown. Refs. – Lynch Arribálzaga, 1881: 26; Kertész, 1909: 124 (cat.; *Stenobasis*); Carrera, 1949: 49, 1953a: 273; Carrera & Papavero, 1962: 47; Hull, 1962 (1): 233; Martin & Papavero, 1970: 28 (cat.); Papavero & Bernardi, 1973: 124 (syn.). Type, lost.
- luridum** Rondani, 1848: 89. Type-locality: “Brazil”[i. e., Pará, Belém, cf. Papavero, 1973d: 343]. Neotype-locality: Brazil, Pará, Oriximiná. Distr. – Guyana, Surinam, French Guiana, Peru (Loreto), Brazil (Amazonas, Pará, Mato Grosso). Refs. – Lynch Arribálzaga, 1881: 24 (as syn. of *coarctatum* Perty, 1833); Williston, 1891: 75 (cat.; as syn. of *coarctatum* Perty, 1833); Kertész, 1909: 124 (cat.; *Stenobasis*, as syn. of *coarctatum* Perty, 1833); Carrera, 1949: 47, fig. 60 (revalidation of sp.); Carrera & d'Andretta, 1953: 66; Carrera & Papavero, 1962: 47; Hull, 1962 (1): 233; Martin & Papavero, 1970: 28 (cat.); Lamas M., 1972: 307 (cat.); Papavero & Bernardi, 1973: 188, figs. 32-34, map 3; Artigas & Papavero, 1988b: 256, fig. 196. Type, lost. NT MZUSP [des. by Papavero & Bernardi, 1973: 188].
- modesta* Bigot, 1878: 472 (*Senobasis*). Type-locality: “Amazon”. Refs. – Williston, 1891: 75 (cat.; *Senobasis*); Kertész, 1909: 124 (cat.; *Stenobasis*); Bromley, 1946: 106 (cat.); Hull, 1962 (1): 233; Martin & Papavero, 1970: 58 (cat.; *Senobasis*); Papavero & Bernardi, 1973: 188 syn.). Type, OXF.
- sonorense** Papavero & Bernardi, 1973: 175, figs. 7-9, map 3 (as *sonorensis*, error). Type-locality: U. S. A., Arizona, Palos Verdes. Distr. – U. S. A. (Utah, Arizona, Texas), Mexico (Tamaulipas, Coahuila, Nayarit, Durango, Chihuahua), Cuba. Ref. – Nelson, 1987: 59. Type, MZUSP.
- coarctatum* Perty of Snow, 1896: 177, Aldrich, 1905: 263 (cat.) and Martin, 1965a: 19, misident.
- secabile* Walker of Wilcox, 1937: 37 and Linsley, 1960 (biol. notes), misident.
- subcontractum** (Walker), 1856: 455 (*Dasypogon*). Type-locality: Brazil, “Amazon district”. Distr. – Costa Rica, Guyana, Peru (Loreto), Brazil (Amazonas, Pará). Refs. – Lynch Arribálzaga, 1881: 24; Williston, 1891: 76 (cat.); Kertész, 1909: 124 (cat.; *Stenobasis*, as syn. of *coarctatum* Perty, 1833); Carrera, 1949: 52, 1953a: 273; Hull, 1962 (1): 233; Martin & Papavero, 1970: 28 (cat.); Lamas M., 1972: 307 (cat.); Papavero & Bernardi, 1973: 190, figs. 38-40, map 2. Type, BMNH.
- bassleri* Curran, 1942: 54. Type-locality: Peru, río Marañoñ. Refs. – Carrera, 1949: 58; Hull, 1962 (1): 233; Martin & Papavero, 1970: 28 (cat.; syn.). Type, AMNH.
- surumu** Papavero & Bernardi, 1973: 177, map 3. Type-locality: Brazil, Roraima, Surumu. Distr. – Venezuela (Bolívar), Brazil (Roraima). Type, MZUSP.

Genus *Caenarolia* Thomson, 1869

- Allopogon* Schiner, 1886: 670 (preocc. Horn, 1880). Type-species, *Dasypogon vittatus* Wiedemann, 1828 (orig. des.).
- Caenarolia** Thomson, 1869: 470. Type-species, *Caenarolia longipennis* Thomson, 1869 (mon.) = *Caenarolia equestris* (Wiedemann, 1828).
- Caenorolia* Williston, 1891: 74 (cat.), error.

Ref. – Artigas & Papavero, 1988b: 205 (as *Allopogon*; key to spp.).

- anomala** Carrera, 1947a: 40. Type-locality: Brazil, São Paulo, São Paulo (Ipiranga). Distr. – Brazil (Mato Grosso, São Paulo), Paraguay, Argentina (Corrientes, Formosa). Refs. – Carrera, 1949: 68, figs. 40-41, 175, 1953b: 186, fig. 10; Carrera & d'Andretta, 1961: 68 (prey); Carrera & Papavero, 1962: 43; Hull, 1962 (1): 232; Martin & Papavero, 1970: 24 (cat.). Type, MZUSP. **N. comb.**

- argyrocincta** (Schiner), 1867: 370 (*Saropogon*). Type-locality: “Brazil”. Distr. – Brazil (Rio de Janeiro). Refs. – Williston, 1891: 74 (cat.; *Saropogon*); Bromley, 1946: 107 (cat.; *Saropogon*); Carrera, 1949: 36, fig. 31; Carrera & Papavero, 1962: 49; Hull, 1962 (1): 231, figs. 146, 1718, 1726, 1730; Martin & Papavero, 1970: 24 (cat.); Artigas & Papavero, 1988: 206 (*Allopogon*). Type, NMW. **N. comb.**

- dimidiatus* Curran, 1935: 3 (*Allopogon*; in key), 4. Type-locality: Brazil, Rio de Janeiro. Refs. – Bromley, 1946: 106 (cat.; *Allopogon*); Carrera, 1949: 36 (*Allopogon*; syn.); Hull, 1962 (1): 229 (*Allopogon*). Type, AMNH.

- basalis** (Curran), 1935: 3 (*Allopogon*). Type-locality: “Brazil”. Distr. – Brazil (Minas Gerais, Paraná). Refs. – Bromley, 1946:

- 106 (*Allopogon*; cat.); Carrera, 1949: 37 163, figs. 19, 36, 1953a: 272; Hull, 1962 (1): 231, fig. 162, 539, 1055, 1064; Martin & Papavero, 1970: 24 (cat.). Type, AMNH.
- castigans** (Walker), 1851: 89 (*Dasypogon*). Type-locality: "South America". Distr. – Brazil (Minas Gerais). Refs. – Walker, 1854: 433 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 126 (cat.; *Dasypogon*); Papavero, 1971: 20 (*Diogmites*); Artigas & Papavero, 1988b: 207 (*Allopogon*). Type, BMNH. **N. comb.**
- spitzi** Carrera, 1949: 39, figs. 33, 161. Type-locality: Brazil, Minas Gerais, Araguari. Refs. – Hull, 1962 (1): 231; Martin & Papavero, 1970: 24 (cat.); Artigas & Papavero, 1988b: 207 (*Allopogon*. syn.). Type, MZUSP.
- equestris** (Wiedemann), 1828: 392 (*Dasypogon*). Type-locality: "Brazil". Distr. – Brazil (Rio de Janeiro). Refs. – Walker, 1854: 445 (*Dasypogon*); Schiner, 1866: 678 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 127 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Artigas & Papavero, 1988: 207 (n. comb.). Type, ? **N. comb.**
- longipennis** Thomson, 1869: 471. Type-locality: Brazil, Rio de Janeiro. Refs. – Williston, 1891: 74 (cat.); Kertész, 1909: 57 (cat.); Bromley, 1946: 105 (cat.); Carrera, 1949: 35, figs. 30, 78, 156, 160; Hull, 1962 (1): 231; Martin & Papavero, 1970: 24 (cat.); Artigas & Papavero, 1988: 207 (*Allopogon*; syn.).
- miles** (Wiedemann), 1828: 393 (*Dasypogon*). Type-locality: "Brazil". Refs. – Walker, 1854: 445 (*Dasypogon*); Schiner, 1866: 678 (*Dasypogon*); Williston, 1891: 74 (cat.); Kertész, 1909: 57 (cat.; *Allopogon*); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 231; Martin & Papavero, 1970: 24 (cat.). Type, NMW.
- necans** (Wiedemann), 1828: 392 (*Dasypogon*). Type-locality: "Brazil". Distr. – Brazil (Minas Gerais). Refs. – Walker, 1854: 174 (*Dasypogon*); Schiner, 1866: 678 (*Allopogon*); Williston, 1891: 73 (cat.; *Allopogon*); Kertész, 1909: 118 (cat.; *Allopogon*); Curran, 1935: 3 (*Allopogon*; in key); Bromley, 1946: 106 (cat.; *Allopogon*); Carrera, 1949: 43, fig. 49 (*Allopogon*); Hull, 1962 (1): 229 (*Allopogon*); Martin & Papavero, 1970: 23 (cat.; *Allopogon*); Papavero, 1971: 20 (*Allopogon*); Artigas & Papavero, 1988: 207 (*Allopogon*). **N. comb.**
- asiloides** Bigot, 1878: 426 (*Lochites*). Type-locality: "Brazil". Refs. – Williston, 1891: 75 (cat.; *Senobasis*); Kertész, 1909: 125 (cat.; *Lochites*); Bromley, 1946: 106 (cat.; *Senobasis*); Hull, 1962: 233 (*Blepharepium*); Papavero, 1971: 20 (*Allopogon*; syn.). Type, OXF.
- placida** (Wulp), 1882b: 94 (*Deromyia*). Type-locality: Argentina. Distr. – Argentina. Refs. – Williston, 1891: 75 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Engel, 1930: 469 (*Diogmites*); Carrera, 1949: 45 (*Allopogon*), 1953b: 183 (*Allopogon*, as syn. of *ferrugineus* Lynch Arribálzaga, 1880); Martin & Papavero, 1970: 23 (cat.; *Allopogon*). Type, ZMA. **N. comb.**
- tessellata** (Wiedemann), 1828: 390 (*Dasypogon*). Type-locality: Uruguay, Montevideo. Distr. – Brazil (Ceará, Rio Grande do Norte, Pernambuco, Espírito Santo, Rio Grande do Sul), Argentina (Tucumán, Rioja), Uruguay. Refs. – Walker, 1854: 447 (*Dasypogon*); Schiner, 1866: 678 (*Allopogon*); Lynch Arribálzaga, 1879: 147 (*Allopogon*), 1883a: 138 (cat.; *Allopogon*); Williston, 1891: 74 (cat.; *Allopogon*); Kertész, 1909: 119 (cat.; *Allopogon*); Curran, 1935: 3 (*Allopogon*; in key); Carrera, 1947d: 266 (*Allopogon*; preys), 1949: 45, figs. 18, 46-47 (*Allopogon*), 1960a: 157 (*Allopogon*); Carrera & d'Andretta, 1961: 68 (*Allopogon*; prey); Carrera & Papavero, 1962: 49 (*Allopogon*); Hull, 1962 (1): 229 (*Allopogon*); Martin & Papavero, 1970: 23 (cat.; *Allopogon*); Artigas & Papavero, 1988b: 207, 251, figs. 164-165 (*Allopogon*). Type, ? **N. comb.**
- weyenberghi** Wulp, 1882b: 93 (*Deromyia*). Type-locality: "Argentina". Refs. – Williston, 1891: 75 (cat.; *Deromyia*); Kertész, 1909: 118 (cat.; *Deromyia*); Carrera, 1949: 45 (*Allopogon*); Carrera & Papavero, 1962: 49 (*Allopogon*); Hull, 1962 (1): 229 (*Allopogon*); Martin & Papavero, 1970: 23 (cat.; *Allopogon*). Type, ZMA.
- vittata** (Wiedemann), 1828: 389 (*Dasypogon*). Type-locality: Uruguay, Montevideo. Distr. – Brazil (Paraná, Santa Catarina, Rio Grande do Sul), Uruguay (Treinta y Tres, Rivadavia), Argentina (Buenos Aires, Misiones). Refs. – Walker, 1854: 446 (*Dasypogon*); Schiner, 1866: 670 (*Allopogon*); Lynch Arribálzaga, 1879: 146 (*Allopogon*), 1883a: 138 (cat.; *Allopogon*); Williston, 1891: 74 (cat.; *Allopogon*); Kertész, 1909: 119 (cat.; *Allopogon*); Curran, 1935: 3 (*Allopogon*; in key); Bromley, 1946: 106 (cat.; *Allopogon*); Carrera, 1949: 41, figs. 17, 38-39, 154, 159 (*Allopogon*), 1953a: 272 (*Allopogon*), 1960a: 157 (*Allopogon*); Carrera & Papavero, 1962: 49 (*Allopogon*); Hull, 1962 (1): 229, figs. 175, 542, 1052, 1061, 1747, 1864, 1870 (*Allopogon*); Martin & Papavero, 1970: 23 (cat.; *Allopogon*); Artigas & Papavero, 1988b: 207, 251, figs. 166-167, 256, fig. 195 (*Allopogon*). Type, ? **N. comb.**
- longiungulatus** Macquart, 1838: 36 (1839: 152), pl. 3, fig. 6 (*Dasypogon*). Type-locality: "Brésil, Missions" [= western part of the state of Rio Grande do Sul; cf. Papavero, 1971e: 121]. Refs. – Walker, 1854: 436 (*Dasypogon*); Osten Sacken, 1891: 418 (*Dasypogon*); Kertész, 1909: 119 (*Dasypogon*).
- annulitarsis** Rondani, 1868: 32 (*Dasipogon*). Type-locality: Argentina, Santa Fe, Córdoba, Río Cuarto and Rosario. Refs. – Lynch Arribálzaga, 1880: 28 (*Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*). Type, MZUN.
- gracilis** Bigot, 1878: 418 (*Dasypogon*). Type-locality: Uruguay, Montevideo. Ref. – Kertész, 1909: 119 (cat.).

Genus *Diogmites* Loew, 1866

Diogmites Loew, 1866: 21 (Cent. 7, no. 36). Type-species, *Diogmites platypterus* Loew, 1866 (sub. des., Coquillett, 1910: 533).

Deromyia Philippi of authors, misident.

- aberrans** (Wiedemann), 1821: 223 (*Dasypogon*). Type-locality: "South America". Distr. – Venezuela, Guyana, Ecuador, Brazil (Pará). Refs. – Wiedemann, 1828: 389 (*Dasypogon*); Walker, 1854: 443 (*Dasypogon*); Schiner, 1866: 678 (*Dasypogon*); Williston, 1891: 67 (cat.; *Dasypogon*); Kertész, 1909: 125 (cat.; *Dasypogon*); Bromley, 1934b: 335 (*Deromyia*), 1946a: 107; Carrera, 1949: 82 (n. comb.), 1953b: 184; Hull, 1962 (1): 232; Martin & Papavero, 1970: 24 (cat.).
- examinans** Walker, 1851: 90 (*Dasypogon*). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Walker, 1854: 433, 504 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 127 (cat.; *Dasypogon*); Bromley, 1934b: 335 (*Deromyia*), 1946: 107 (cat.; *Dasypogon*); Carrera, 1948a: 116 (n. comb.), 1949: 82.
- abberans** Bromley, 1946: 107 (cat.), error.
- affinis** (Bellardi), 1851: 173 (73) (*Dasypogon*). Type-locality: "Mexico". Distr. – Mexico. Refs. – Osten Sacken, 1874: 184, 1887: 178; Aldrich, 1905: 264 (cat.); Hull, 1962 (1): 232; Martin & Papavero, 1970: 24 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- alvesi** Carrera, 1949: 79. Type-locality: Brazil, Rio Grande do Norte, Natal. Distr. – Brasil (Rio Grande do Norte, Paraíba). Refs. – Carrera, 1953b: 185, fig. 8, 1960: 158; Hull, 1962 (1): 232; Martin & Papavero, 1970: 24 (cat.). Type, MZUSP.
- amethystinus** Carrera, 1953b: 178, fig. 2. Type-locality: Colombia, Meta, Restrepo, 500 m. Distr. – Colombia (Meta). Refs. – Hull, 1962 (1): 232; Martin & Papavero, 1970: 24 (cat.). Type, Bomley Coll. (USNM?).
- angustipennis** Loew, 1866: 23 (Cent. 7, no. 41). Type-locality: U. S. A., Kansas. Distr. – U. S. A. (Kansas, Colorado, Illinois, Wyoming, Nebraska, New Mexico, Texas, Oklahoma), Mexico (Chihuahua, Coahuila). Refs. – Osten Sacken, 1874: 183; Williston, 1883: 25 (*Deromyia*); Aldrich, 1905: 264 (cat.; *Deromyia*); Hine, 1906: 30, fig. 15 (*Deromyia*); Kertész, 1909: 114 (cat.; *Deromyia*); Baker, 1939: 168; Hull, 1962 (1): 232; Martin & Wilcox, 1965: 371 (cat.); Martin & Papavero, 1970: 24 (cat.). Type, MCZ.
- angustiventris** (Macquart), 1846: 65, pl. 6, fig. 9 (*Dasypogon*). Type-locality: "Mexico". Distr. – Mexico. Refs. – Walker, 1854: 488 (*Dasypogon*); Williston, 1883: 24 (*Deromyia*; as syn. of *winthemi* Wiedemann, 1821); Kertész, 1909: (cat.; as syn. of *winthemi* Wiedemann, 1821); Carrera, 1953b: 195, fig. 20.
- atriapex** Carrera, 1953b: 192, fig. 18. Type-locality: Argentina, Santa Fé, Tartagal. Distr. – Argentina (Santa Fe). Refs. – Hull, 1962 (1): 232; Martin & Papavero, 1970: 24 (cat.). Type, IML.
- aureolus** Carrera & Papavero, 1962: 44, fig. 1. Type-locality: Argentina, Entre Ríos. Distr. – Argentina (Entre Rios). Ref. – Martin & Papavero, 1970: 24 (cat.). Type, MZUSP.
- bellardii** (Bromley), 1929b: 45 (*Deromyia*). Type-locality: Mexico, Distrito Federal. Distr. – Mexico (Distrito Federal, Morelos, Colima). Refs. – Carrera, 1953b: 175; Hull, 1962 (1): 232; Martin & Papavero, 1970: 24 (cat.). Type, USNM.
- bicolor** (Jaennicke), 1867: 357 (*Saropogon*). Type-locality: "Panama". Distr. – Panama. Refs. – Osten Sacken, 1874: 184; Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 115 (cat.; *Deromyia*); Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.).
- bifasciatus** Carrera, 1949: 72, fig. 87. Type-locality: Brazil, São Paulo, Juquiá. Distr. – Brazil (São Paulo). Refs. – Carrera, 1953b: 192, figs. 16-17; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.). Type, MZUSP.
- bigoti** (Bellardi), 1861: 170 (70) (*Dasypogon*). Type-locality: "Mexico". Distr. – U. S. A. (Utah), Mexico. Refs. – Osten Sacken, 1874: 184, 1887: 178; Aldrich, 1905: 264 (cat.; *Deromyia*); Back, 1909: 357 (*Deromyia*); Kertész, 1909: 115 (cat.; *Deromyia*); Brown, 1929: 307, pl. 28, figs. 7-11 (*Deromyia*); Carrera, 1953b: 175; Hull, 1962 (1): 232, 415 (*Atoniomyia*); Martin & Papavero, 1970: 88 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, lost.
- bilineatus** (Loew), 1866: 23 (Cent. 7, no. 40) (*Deromyia*). Type-locality: "Cuba". Distr. – Cuba. Refs. – Osten Sacken, 1874: 184; Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 115 (cat.; *Deromyia*); Bromley, 1929a: 278 (*Deromyia*); Carrera, 1953b: 175; Fattig, 1945: 13 (?misident; U. S. A., Georgia); Hull, 1962 (1): 232; Martin & Wilcox, 1965: 371 (cat.); Martin & Papavero, 1970: 25 (cat.); Alayo Dalmau & García Ávila, 1983: 45 (cat.). Type, MCZ.
- bimaculatus** (Bromley), 1929b: 46 (*Deromyia*). Type-locality: Mexico, Morelos, Cuernavaca. Distr. – Mexico (Morelos, Nayarit). Refs. – Carrera, 1953b: 196; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.). Type, USNM.
- bromleyi** Carrera, 1949: 78, figs. 81, 173. Type-locality: Brazil, Minas Gerais, Cambuquira. Distr. – Brazil (Goiás, Minas Gerais). Refs. – Carrera, 1953b: 198, fig. 22; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.). Type, IOC.
- brunneus** (Fabricius), 1787: 359 (*Asilus*). Type-locality: French Guiana, Cayenne (Rohr) [cf. Papavero, 1971e: 20]. Distr. – Venezuela, French Guiana. Refs. – Gmelin, 1792: 2899 (*Asilus*); Fabricius, 1794: 382 (*Asilus*), 1805: 165 (*Asilus*); Wiedemann, 1821: 219 (*Dasypogon*), 1828: 382 (*Dasypogon*); Macquart, 1838: 34 (*Dasypogon*); Walker, 1849: 338 (*Dasypogon*), 1854: 421 (*Dasypogon*); Bellardi, 1861: 67 (*Dasypogon*); Schiner, 1868: 677 (*Dasypogon*); Osten Sacken, 1874: 184, 1887: 178; Williston, 1891: 75 (cat.; *Deromyia*); Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 115 (cat.; *Deromyia*); Carrera, 1949: 81; Carrera, 1953b: 175; Hull, 1962 (1): 232; Carrera & Machado-Allison, 1963: 235; Martin, 1963: 9, 1965a: 21; Martin & Papavero, 1970: 25 (cat.). Type, lost [cf. Zimsen, 1964: 470].
- castaneus** (Macquart), 1838: 35 (1839: 151) (*Dasypogon*). Type-locality: Brazil, Pará (i. e., Pará, Belém). Distr. – Colombia, Peru (Loreto), Brazil (Amazonas and Pará to Rio de Janeiro and São Paulo), Bolivia. Refs. – Walker, 1849: 338 (*Dasypogon*), 1854: 431 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 62 (cat.; ?*Oligopogon*); Bromley, 1946: 105 (cat.; *Oligopogon*); Carrera, 1948a: 118, figs. 1-2 (n. comb.), 1949: 85, figs. 48, 176, 1953b: 185, fig. 9; Carrera & d'Andretta, 1953: 66; Carrera, 1960a: 158; Carrera & Papavero, 1962: 43; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Papavero, 1971: 20; Lamas M., 1972: 306 (cat.); Fisher, 1984: 28; Artigas & Papavero,

- 1988b: 256, fig. 197. Type, MNHNP.
- fulvus** Bigot, 1878: 426 (*Dasypogon*). Type-locality: Brazil, Amazonia. Refs. – Williston, 1891: 75 (cat.; *Senobasis*); Kertész, 1909: 125 (cat.; *Lochites*); Hull, 1962: 233 (*Blepharepium*); Martin & Papavero, 1970: 58 (*Senobasis*); Papavero, 1971d: 20 (syn.). Type, OXF.
- fulvis** Bromley, 1946: 106 (cat.; *Senobasis*), error.
- coffeatus** (Wiedemann), 1819: 49 (*Dasypogon*). Type-locality: “Brazil”. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Wiedemann, 1821: 217 (*Dasypogon*), 1828: 374 (*Dasypogon*); Walker, 1854: 444 (*Dasypogon*); Schiner, 1866: 676 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 127 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Carrera, 1949: 65, figs. 16, 45, 174, 1953b: 189, figs. 3-4; Carrera & d’Andretta, 1961: 68 (preys); Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Artigas & Papavero, 1988b: 258, fig. 199. Type, NMW.
- craverii** (Bellardi), 1861: 168 (68) (*Dasypogon*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Osten Sacken, 1874: 184, 1887: 178; Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 115 (cat.; *Deromyia*); Carrera, 1953b: 175; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- cuantlensis** (Bellardi), 1861: 167 (68) (*Dasypogon*). Type-locality: Mexico, Morelos, Cuautla. Distr. – Mexico (Guerrero, Chiapas, Jalisco, Nuevo León, Oaxaca, Puebla, Veracruz, Yucatán). Refs. – Osten Sacken, 1874: 184, 1887: 175; Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 115 (cat.; *Deromyia*); Carrera & d’Andretta, 1950: 163, figs. 35-40; Carrera, 1953b: 189, fig. 26; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- dubius** (Bellardi), 1861: 174 (74) (*Dasypogon*). Type-locality: Mexico, Morelos, Cuautla. Distr. – Mexico (Guerrero), Costa Rica. Refs. – Osten Sacken, 1874: 184, 1887: 178; Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 115 (cat.; *Deromyia*); Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- duillius** (Walker), 1849: 340 (*Dasypogon*). Type-locality: “Honduras”. Distr. – Honduras. Refs. – Walker, 1854: 431 (*Dasypogon*); Osten Sacken, 1874: 184, 1878: 73, 233 (cat.), 1887: 178; Williston, 1891: 75 (cat.; *Deromyia*); Aldrich, 1905: 264 (*Deromyia*, as *duillia*); Kertész, 1909: 116 (cat.; *Deromyia*); Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.). Type, BMNH.
- fasciatus** (Macquart), 1834: 295 (*Dasypogon*). Type-locality: French Guiana, Cayenne. Distr. – French Guiana. Syntypes, MNHNP, OXF. Refs. – Walker, 1854: 440 (*Dasypogon*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 128 (cat.; *Dasypogon*); Bromley, 1934b: 335 (*Deromyia*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 25 (cat.).
- ferrugineus** (Lynch Arribálzaga), 1880: 182 (*Allopogon*). Type-locality: Argentina, Mendoza. Distr. – Brazil, Bolivia, Argentina. Refs. – Lynch Arribálzaga, 1883a: 137 (cat.; *Allopogon*); Williston, 1891: 73 (cat.; *Allopogon*); Kertész, 1909: 118 (cat.; *Allopogon*); Carrera, 1949: 83 (n. comb.), 1953b: 183, fig. 7, 1959: 1; Carrera & Papavero, 1962: 42; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Artigas & Papavero, 1988: 252, figs. 173-177, 157, fig. 198.
- goniostigma** (Bellardi), 1861: 165 (65), pl. 1, fig. 6 (*Dasypogon*). Type-locality: Mexico, Vera Cruz, Playa Vicente. Distr. – México (Vera Cruz). Refs. – Osten Sacken, 1874: 184, 1887: 178; Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 116 (cat.; *Deromyia*); Carrera, 1953b: 175; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- heydenii** (Jaennicke), 1867: 356, pl. 44, fig. 1 (*Dasypogon*). Type-locality: Argentina (Corrientes). Distr. – Argentina (Corrientes). Type, SENCK [cf. Geller-Grimm, 1999: 200]. Refs. – Lynch Arribálzaga, 1880: 137 (*Dasypogon*), 1883a: 137 (cat.; *Dasypogon*); Williston, 1891: 73 (cat.; *Allopogon*); Kertész, 1909: 118 (cat.; *Allopogon*); Carrera, 1953b: 176; Hull, 1962 (1): 229 (*Allopogon*); Martin & Papavero, 1970: 25 (cat.).
- imitator** Carrera, 1953b: 101. Type-locality: Brazil, Pará (i. e., Pará, Belém). Distr. – Brazil (Pará). Refs. – Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.). Type, BMNH.
- inclusus** (Walker), 1851: 95 (*Dasypogon*). Type-locality: Brazil, Pará (i. e., Pará, Belém). Distr. – Venezuela, Peru, Brazil (Amazonas, Pará). Refs. – Walker, 1854: 433 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 128 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Carrera, 1948a: 119 (n. comb.), 1949: 76, 1953b: 199, fig. 25, 1960a: 158; Carrera & Papavero, 1962: 46; Hull, 1962 (1): 232; Martin & Papavero, 1970: 25 (cat.); Fisher, 1984: 28.
- parvus** Carrera, 1948a: 120, fig. 3. Type-locality: Brazil, Amazonas, rio Parauari. Type, IOC.
- intactus** (Wiedemann), 1828: 371 (*Dasypogon*). Type-locality: “South America”. Distr. – Brazil. Refs. – Walker, 1854: 445 (*Dasypogon*); Schiner, 1866: 676 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 129 (cat.; *Dasypogon*); Carrera, 1949: 82 (n. comb.); Carrera & d’Andretta, 1950: 162, figs. 4, 11, 25, 29, 31-33 (misident.); Carrera, 1953b: 187, figs. 11-12 (misident.); Hull, 1962 (1): 232; Martin, 1965a: 22; Martin & Papavero, 1970: 26 (cat.). Type, NMW.
- jalapensis** (Bellardi), 1861: 165 (65), pl. 1, fig. 5 (*Dasypogon*). Type-locality: Mexico, Veracruz, Jalapa. Distr. – México (Veracruz), Honduras, Panama. Refs. – Osten Sacken, 1874: 184, 1887: 177; Williston, 1901: 311 (*Deromyia*); Aldrich, 1905: 264 (cat.; *Deromyia*); Kertész, 1909: 264 (cat.; *Deromyia*); Carrera & d’Andretta, 1950: 163; Carrera, 1953b: 200; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- jalapensis** James, 1953: 56, error.
- lindigi** (Schiner), 1868: 165 (*Dasypogon*). Type-locality: “South America”. Distr. – Costa Rica, Panama, Venezuela, Paraguay, Brazil (Santa Catarina). Refs. – Osten Sacken, 1887: 174; Williston, 1891: 75 (cat.; *Deromyia*); Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 116 (cat.; *Deromyia*); Carrera, 1947a: 39, 1953b, figs. 13-14: 188; Carrera & Papavero,

- 1962: 43; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.). Type, NMW.
- lineola** (Bromley), 1934b: 335, fig. 22 (*Deromyia*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Distr. – Brazil (Mato Grosso). Refs. – Bromley, 1946: 107 (cat.; n. comb.); Carrera, 1949: 83, 1953b: 176; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.). Type, AMNH.
- litoralis** (Curran), 1930b: 1 (*Deromyia*). Type-locality: Panama, Canal Zone, Bruja Point. Distr. – Panama. Refs. – Carrera, 1953b: 182, fig. 6; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.). Type, AMNH.
- maculatus** Curran, 1934a: 1. Type-locality: Ecuador, Guayas, Golfo de Guayaquil, Puerto Grande, Isla Puná. Distr. – Ecuador. Refs. – Carrera, 1949: 81, 1953b: 176; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.). Type, AMNH.
- memnon** Osten Sacken, 1887: 174, pl. 3, fig. 9. Type-locality: Costa Rica; Panama, Volcán de Chiriquí, 2000-3000 ft. Distr. – Costa Rica, Panama. Refs. – Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 116 (cat.; *Deromyia*); Carrera, 1953b: 176; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.). Type, BMNH.
- neoternatus** (Bromley), 1931b: 433 (*Deromyia*). Type-locality: U. S. A., Kansas, Russell County. Distr. – U. S. A. (widespread), Mexico (Sonora). Refs. – Fattig, 1945: 15; Hull, 1962 (1): 232; Martin & Wilcox, 1965: 371 (cat.); Martin & Papavero, 1970: 26 (cat.); Dennis & Gowen, 1978: 366 (“nocturnal” foraging). Type, OSU.
- nigripennis** (Macquart), 1847: 50 (1847: 34), pl. 1, fig. 6 (*Dasypogon*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Walker, 1854: 428 (*Dasypogon*); Bellardi, 1861: 75 (*Dasypogon*); Osten Sacken, 1874: 184; Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 116 (cat.; *Deromyia*); Carrera, 1953b: 176; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.); Papavero & Ibáñez Bernal, 2001: 85 (hist. Mexican dipterol.). Type, OXF.
- nigripes** (Bellardi), 1861: 175 (75) (*Dasypogon*). Type-locality: Mexico, Vera Cruz, Playa Vicente. Distr. – Mexico (Veracruz). Refs. – Osten Sacken, 1874: 184, 1887: 178; Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 116 (cat.; *Deromyia*); Carrera, 1953b: 176; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol.). Type, TORO.
- nigritarsis** (Macquart), 1846: 196 (1846: 68) (*Dasypogon*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Walker, 1854: 428 (*Dasypogon*); Osten Sacken, 1874: 185; Aldrich, 1905: 267 (cat.; *Dasypogon*); Kertész, 1909: 130 (cat.; *Dasypogon*); Martin & Papavero, 1970: 26 (cat.); Papavero & Ibáñez Bernal, 2001: 84 (hist. Mexican dipterol.). Type, MNHNP.
- obscurus** Carrera, 1949: 67, fig. 177. Type-locality: Brazil, Goiás. – Brazil (Goiás), Bolivia, Argentina. Refs. – Carrera, 1953b: 185, 1959: 1; Carrera & d’Andretta, 1961: 68 (preys); Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.). Type, MZUSP.
- perplexus** (Back), 1909: 360 (*Deromyia*). Type-locality: U. S. A., New Mexico. Distr. – U. S. A. (New Mexico, Arizona, Colorado), Mexico. Refs. – Hull, 1962 (1): 232; Martin & Wilcox, 1965: 371 (cat.).
- pseudojalapensis** (Bellardi), 1862: 222 (25) (*Dasypogon*). Type-locality: Mexico, Veracruz, Tuxpan, near Orizaba. Distr. – Mexico (Veracruz). Refs. – Osten Sacken, 1887: 178; Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Carrera, 1953b: 176; Hull, 1962 (1): 232; Martin & Papavero, 1970: 26 (cat.); Papavero & Ibáñez Bernal, 2001: 110 (hist. Mexican dipterol.). Type, TORO.
- pseudojalapensis* Osten Sacken, 1874: 184; Kertész, 1909: 117 (cat.; *Deromyia*), emend.
subialapensis, error.
- reticulatus** (Fabricius), 1805: 160 (*Laphria*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Guyana, Surinam, Brazil (Amazonas, Pará). Refs. – Wiedemann, 1821: 217 (*Dasypogon*), 1828: 378 (*Dasypogon*); Walker, 1854: 445 (*Dasypogon*); Schiner, 1866: 677 (*Dasypogon*); Williston, 1891: 67 (cat.; *Dasypogon*); Kertész, 1909: 131 (cat.; *Dasypogon*); Bromley, 1934b: 335; Carrera, 1953b: 181, fig. 5, 1960a: 158; Carrera & Papavero, 1962: 42; Hull, 1962 (1): 228 (*Dasypogon*), 233 [*Diogmites*]; Carrera & Machado-Allison, 1963: 236; Martin & Papavero, 1970: 26 (cat.). Syntypes, COPN [cf. Zimšen, 1964: 468].
- nigricauda* Curran, 1925a: 259 (*Deromyia*). Type-locality: Guyana, Cuyuni-Mazaruni, Bartica. Refs. – Bromley, 1934b: 335 (*Deromyia*); Carrera, 1949: 66 (n. comb.). Type, AMNH.
- rubescens** (Bellardi), 1861: 171 (71) (*Dasypogon*). Type-locality: Mexico, Distrito Federal, México, and Veracruz, Tuxpan. Distr. – Mexico (Veracruz, Distrito Federal). Refs. – Osten Sacken, 1874: 184, 1887: 176; Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Carrera & d’Andretta, 1950: 163; Carrera, 1953b: 176; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol.). Syntypes, TORO.
- rufibasis** Bigot, 1878: 416. Type-locality: “Brazil”. Refs. – Kertész, 1909: 117 (cat.; *Deromyia*); Bromley, 1946: 107 (cat.); Carrera, 1953b: 177; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.). Type, OXF.
- rufitarsis* Williston, 1891: 75, error.
- rufipalpis** (Macquart), 1838: 38 (1939: 154), pl. 3, fig. 2 (*Dasypogon*). Type-locality: “?French Guiana, Cayenne”. Distr. – Peru, Brazil (Amazonas, Pará). Refs. – Walker, 1854: 432 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 131 (cat.; *Dasypogon*); Carrera, 1953b: 190, fig. 15; Carrera & Papavero, 1962: 43; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Fisher, 1984: 28.
- sallei** (Bellardi), 1861: 170 (70) (*Dasypogon*). Type-locality: Mexico, Distrito Federal, México, and Veracruz, Tuxpan. Distr. – Mexico (Sonora, Chihuahua, Jalisco, Colima, Chiapas). Refs. – Osten Sacken, 1874: 184, 1887: 174, pl. 3, fig. 8; Williston, 1901: 311 (*Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Carrera, 1953b: 177; Carrera & Papavero, 1962: 42; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol.). Syntypes, TORO.
- sallaei* Aldrich, 1905: 117 (cat.; *Deromyia*), error.
- superbus** Carrera, 1953b: 177, fig. 1. Type-locality: Costa Rica, Turrialba, 620 m. Distr. – Costa Rica. Refs. – Hull, 1962 (1):

- 232; Martin & Papavero, 1970: 27 (cat.). Type, MZUSP.
- symmachus** Loew, 1872: 66 (Cent. 10, no. 26). Type-locality: U. S. A., Texas. Distr. – U. S. A., Mexico (Chihuahua). Refs. – Osten Sacken, 1874: 183; Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Bromley, 1935c: 3 (n. comb.); Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Artigas & Papavero, 1988: 258, fig. 202. Type, MCZ.
- tau** Osten Sacken, 1887: 176, pl. 3, fig. 11. Type-locality: Panama, Volcán de Chiriquí, 3000-4000 ft. Distr. – Costa Rica, Panama. Refs. – Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Carrera, 1953b: 197; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.). Type, BMNH.
- teresita** Lamas M., 1972: 306, figs. 1-4. Type-locality: Peru, Ayacucho, La Mar (Hacienda Teresita). Distr. – Peru (Ayacucho). Type, MEUNAL.
- ternatus** (Loew), 1866: 22 (Cent. 7, no. 38) (*Deromyia*). Type-locality: “Cuba”. Distr. – Cuba. Refs. – Osten Sacken, 1874: 184; Williston, 1883: 25 (*Deromyia*); Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Hine, 1918: 3 (*Deromyia*); Bromley, 1929a: 278 (*Deromyia*); Carrera, 1953b: 198; Hull, 1962 (1): 232; Martin & Wilcox, 1965: 371 (cat.); Martin & Papavero, 1970: 27 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 45 (cat.). Type, MCZ.
- tricolor** (Bellardi), 1861: 169 (69) (*Dasypogon*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Osten Sacken, 1874: 184; Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Carrera, 1953b: 177; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- virescens** (Bellardi), 1861: 169 (69) (*Dasypogon*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Osten Sacken, 1874: 184; Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Carrera, 1953b: 182; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- vulgaris** Carrera, 1947d: 266 (descr. and prey). Type-locality: Brazil, São Paulo, Guatapará. Distr. – Peru, Brazil (Mato Grosso, Goiás, Minas Gerais, São Paulo). Refs. – Carrera, 1949: 69, figs. 12, 42-43, 82, 114, 145, 149, 203-210, 1953b: 194, fig. 19; Carrera & Papavero, 1962: 44; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Fisher, 1984: 28; Artigas & Papavero, 1988: 251, figs. 162-163; Dennis & Knutson, 1988: 853, fig. 1 (pupa in ventral, lateral and dorsal views; host: *Dyscinetus rugifrons* (Burmeister, 1847) (Coleoptera, Scarabaeridae, Dynastinae)). Type, MZUSP.
- winthemi** (Wiedemann), 1821: 223 (*Dasypogon*). Type-locality: “South America”. Distr. – Colombia, Brazil (Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo). Refs. – Wiedemann, 1828: 387, pl. 6, fig. 2 (*Dasypogon*); Walker, 1849: 332 (*Dasypogon*), 1854: 432 (*Dasypogon*); Schiner, 1868: 678 (*Dasypogon*); Wulp, 1882b: 93 (*Deromyia*), 1884: 207 (*Deromyia*); Williston, 1883: 24, pl. 2, fig. 6 (*Deromyia*); Osten Sacken, 1887: 177; Williston, 1891: 75 (cat.; *Deromyia*); Aldrich, 1905: 265 (cat.; *Deromyia*); Kertész, 1909: 118 (cat.; *Deromyia*); Malloch, 1915 (pupa; misident.: actually *Diogmites misellus* Loew, 1866, a Nearctic species; cf. Dennis & Knutson, 1988: 852); Bromley, 1946: 107 (cat.); Carrera, 1949: 73, figs. 11-178, 89, 116, 1953b: 196, fig. 21, 1960a: 158; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.); Artigas & Papavero, 1988: 258, fig. 199. Type, NMW.
- wygodzinskyi** Carrera, 1949: 77, figs. 90, 146-147. Type-locality: Brazil, Rio de Janeiro, Seropédica (as km 40 da antiga rodovia Rio-São Paulo). Distr. – Brazil (Rio Grande do Norte, Rio de Janeiro, São Paulo). Refs. – Carrera, 1953b: 198, figs. 23-24; Hull, 1962 (1): 232; Martin & Papavero, 1970: 27 (cat.). Type, UFRuRJ.

Genus *Lastaurina* Curran, 1934

- Lastaurina** Curran, 1934a: 171. Type-species, *Dasypogon ardens* Wiedemann, 1828 (sub. des., Curran, 1935: 5).
- Lastauronia* Carrera, 1949: 104. Type-species, *Lastauronia travassosi* Carrera, 1949 (orig. des.). Ref. – Artigas & Papavero, 1988: 209 (syn.).

Ref. – Artigas & Papavero, 1988b: 209 (key to spp.).

- ardens** (Wiedemann), 1828: 391 (*Dasypogon*). Type-locality: “Brazil”. Distr. – Brazil (São Paulo, Paraná, Rio Grande do Sul), Paraguay, Argentina (Buenos Aires). Refs. – Walker, 1854: 446 (*Dasypogon*); Schiner, 1866: 678 (*Lastaurus*); Lynch Arribálzaga, 1879: 151 (*Lastaurus*), 1883a: 140 (cat.; *Lastaurus*); Williston, 1891: 73 (cat.; *Lastaurus*); Kertész, 1909: 144 (cat.; *Lastaurus*); Engel, 1930: 468 (*Lastaurus*); Curran, 1935: 5; Carrera, 1945a: 160, fig. 2 (*Lastaurus*; prey); Bromley, 1946: 108 (cat.; *Lastaurus*); Carrera, 1947a: 43, 1949: 106, figs. 63, 76, 106, 168, 1953a: 275, 1959: 2; Hull, 1962 (1): 239, figs. 544, 1070, 1079; Martin & Papavero, 1970: 30 (cat.); Artigas & Papavero, 1988b: 253, figs. 178-182, 259, fig. 205. Syntypes, BZM [no. 373; cf. Schumann, 1973: 87].
- biezankoi** (Carrera & Papavero), 1962: 50, fig. 6 (*Lastauropsis*). Type-locality: Brazil, Rio de Janeiro, Duque de Caxias, São Bento. Distr. – Brazil (Espírito Santo, Rio de Janeiro). Refs. – Martin & Papavero, 1970: 30 (cat.; *Lastauropsis*); Artigas & Papavero, 1988b: 210 (n. comb.). Type, MZUSP.
- travassosi** (Carrera), 1949: 105, figs. 52, 69, 104 (*Lastauronia*). Type-locality: Brazil, São Paulo, São Paulo. Distr. – Brazil (São Paulo). Refs. – Hull, 1962 (1): 240; Martin & Papavero, 1970: 29 (cat.; *Lastauronia*); Artigas & Papavero, 1988b: 210 (n. comb.), 251, fig. 159. Type, MZUSP.

Genus **Lastaurus** Loew, 1851

Dasypogon, subg. **Lastaurus** Loew, 1851: 11. Type-species, *Lastaurus anthracinus* Loew, 1851 (orig. des.) = *lugubris* (Macquart, 1846).

Morimna Walker, 1851: 104. Type-species, *Morimna mallophorides* Walker, 1851 (mon.) = *Lastaurus fallax* (Macquart, 1838). *Lastauropsis* Carrera, 1949: 107. Type-species, *Lastauropsis villosus* Carrera, 1949 (orig. des.). Ref. – Artigas & Papavero, 1988: 210 (syn.).

alticola Carrera & Machado-Allison, 1968: 500. Type-locality: Bolivia, Cochabamba, Tiraque, 3200 m. Distr. – Bolivia (Cochabamba), Argentina (Jujuy). Refs. – Martin & Papavero, 1970: 29 (cat.); Artigas & Papavero, 1988b: 210. Type, BASEL.

crassitarsis (Macquart), 1838: 36 (1839: 152) (*Dasypogon*). Type-locality "Brazil". Distr. – Brazil (Rio de Janeiro, São Paulo, Santa Catarina). Refs. – Walker, 1854: 435 (*Dasypogon*); Williston, 1891: 76 (*Dasypogon*); Kertész, 1909: 127 (cat.; *Dasypogon*); Carrera, 1949: 100, fig. 167 (*Lastauroides*), 1958a: 146 (*Lastauroides*); Carrera & Papavero, 1962: 54 (*Lastauroides*); Hull, 1962 (1): 241 (*Lastauroides*); Martin & Papavero, 1970: 29 (cat.; *Lastauroides*); Artigas & Papavero, 1988b: 210 (n. comb.). Type, MNHNP.

atratus Bigot, 1878: 415 (*Diogmites*). Type-locality: "Brazil". Refs. – Williston, 1891: 75 (cat.; *Deromyia*); Kertész, 1909: 114 (cat.; *Deromyia*); Bromley, 1946: 107 (cat.; as *atrata*, error).

crassitarsus Bromley, 1946: 107 (cat.; *Dasypogon*), error. Type, OXF.

fallax (Macquart), 1846: 191 (1846: 63), pl. 7, fig. 5 (*Dasypogon*). Type-locality: "Colombia" (as "Nouvelle Grenade"). Distr. – Colombia, Venezuela (Aragua, Distrito Federal, Táchira), Ecuador, Peru (Cuzco), ?Bolivia (Chapare, Cochabamba), ?:Argentina (Tucumán). Refs. – Loew, 1851: 12 (n. comb.); Walker, 1854: 435 (*Dasypogon*); Schiner, 1867: 373; Williston, 1891: 73 (cat.); Kertész, 1909: 144 (cat.); Carrera, 1949: 91; Hull, 1962 (1): 239; Carrera & Machado-Allison, 1963: 235, 1966: 490; Martin & Papavero, 1970: 30 (cat.); Lamas M., 1972: 307 (cat.); Artigas & Papavero, 1988: 210, 255, figs. 190-194. Type, OXF.

bombimorpha Rondani, 1850: 368 (*Dasipogon*). Type-locality: "America equatoriale". Refs. – Osten Sacken, 1891: 418 (*Dasypogon*); Kertész, 1909: 126 (cat.; *Dasypogon*); Carrera, 1949: 92 (n. comb.); Hull, 1962 (1): 239; Martin & Papavero, 1970: 29 (cat.). Type, MZUN.

mutabilis Loew, 1851: 12. Type-locality: "Colombia [Moritz; cf. Papavero, 1973d: 297-298]". Refs. – Osten Sacken, 1887: 180; Aldrich, 1905: 266 (cat.); Kertész, 1909: 145 (cat.); Hull, 1962 (1): 239; Martin & Papavero, 1970: 30 (cat.). Type, BZM [no. 578; cf. Schumann, 1973: 136].

mallophorides Walker, 1851: 104, pl. 6, figs. 2, 22 (*Morimna*). Type-locality: "Colombia". Refs. – Walker, 1854: 429 (*Morimna*); Williston, 1891: 73 (cat.); Martin & Papavero, 1970: 30 (cat.). Type, BMNH.

flavipellitus Enderlein, 1914b: 173 (as *fallax* var.). Type-locality: Ecuador: Bolívar, Balzapamba; Tungurahua: Hacienda Santa Inés, 1244 m (10°25'S, 78°12'W) [cf. Brown, 1941: 846] and Baños. Syntypes, IZPAN.

mallophoroides Kertész, 1909: 144 (cat.); Carrera, 1949: 91; Hull, 1962 (1): 239; Carrera & Machado-Allison, 1968: 493, error.

fenestratus Bigot, 1878: 424. Type-locality: "Colombia" (as "Nouvelle Grenade"). Distr. – Colombia. Refs. – Williston, 1891: 73 (cat.); Kertész, 1909: 144 (cat.); Carrera, 1949: 92; Hull, 1962 (1): 239; Carrera & Machado-Allison, 1966: 495; Martin & Papavero, 1970: 30 (cat.). Type, OXF.

lugubris (Macquart), 1846: 192 (1846: 64) (*Dasypogon*). Type-locality: "Colombia" (as Nouvelle Grenade). Distr. – Mexico, Colombia, Venezuela (Aragua, Carabobo, Distrito Federal, Táchira), ?Argentina (Jujuy, Tucumán). Refs. – Loew, 1851: 12 (n. comb.); Walker, 1854: 435 (*Dasypogon*); Schiner, 1867: 373; Williston, 1891: 73 (cat.); Kertész, 1909: 144 (cat.); Curran, 1941: 272 (in key; *Mallophora*); Carrera, 1953a: 274; Hull, 1962 (1): 239; Carrera & Machado-Allison, 1966: 188; Martin & Papavero, 1970: 30 (cat.); Artigas & Papavero, 1988b: 210, 251, fig. 160, 259, fig. 207. Type, lost.

anthracinus Loew, 1851: 12 (*Dasypogon*). Type-locality: "Mexico". Refs. – Osten Sacken, 1874: 185, 1887: 179, pl. 3, fig. 10; Aldrich, 1905: 265 (cat.); Kertész, 1909: 144 (cat.); Carrera, 1949: 90; Hull, 1962 (1): 239, figs. 159, 548, 1069, 1078, 1950, 2008; Martin & Papavero, 1970: 29 (cat.). Type, BZM [no. 180; cf. Schumann, 1973: 86].

robustus Carrera, 1949: 93, figs. 24, 57, 67, 103. Type-locality: Brazil, Minas Gerais, Araguari. Distr. – Brazil (Minas Gerais). Refs. – Hull, 1962 (1): 239; Carrera & Machado-Allison, 1966: 496; Martin & Papavero, 1970: 30 (cat.); Artigas & Papavero, 1988b: 210, 251, fig. 161. Type, MZUSP.

transiens (Walker), 1849: 364 (*Dasypogon*). Type-locality: "Venezuela". Refs. – Walker, 1854: 110 (*Dasypogon*, *Morimna*); Williston, 1891: 73 (cat.; n. comb.); Kertész, 1909: 145 (cat.); Carrera, 1949: 91; Hull, 1962 (1): 239; Martin & Papavero, 1970: 30 (cat.).

tricolor Carrera & Machado-Allison, 1968: 497. Type-locality: Argentina, Tucumán, Araoz. Distr. – Argentina (Jujuy, Tucumán). Refs. – Martin & Papavero, 1970: 30 (cat.); Artigas & Papavero, 1988: 210. Type, IML.

villosus (Carrera), 1949: 108, figs. 29, 54, 77, 107 (*Lastauropsis*). Type-locality: Brazil, Paraná, Rio Negro. Distr. – Brazil (Paraná). Refs. – Carrera & Papavero, 1962: 49; Hull, 1962 (1): 239, figs. 1072, 1081; Martin & Papavero, 1970: 30 (cat.); *Lastauropsis*; Artigas & Papavero, 1988b: 210. Type, MZUSP.

Genus *Neodiogmites* Carrera, 1949

Neodiogmites Carrera, 1949: 85. Type-species, *Dasypogon melanogaster* Wiedemann, 1821 (orig. des.).
Lastauroides Carrera, 1949: 94. Type-species, *Lastauroides alexanderi* Carrera, 1949 (orig. des.). Ref. – Artigas & Papavero, 1988b: 212 (syn.).
Lastaurax Carrera, 1949: 109. Type-species, *Lastaurax lanei* Carrera, 1949 (orig. des.). Ref. – Artigas & Papavero, 1988b: 212 (syn.).
Ref. – Artigas & Papavero, 1988b: 212 (key to spp.).

alexanderi (Carrera), 1949: 95, figs. 25-26, 50, 68, 105 (*Lastauroides*). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Juçaral. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Carrera, 1958a: 146 (*Lastauroides*); Carrera & Papavero, 1962: 53; Hull, 1962 (1): 241 (*Lastauroides*); Martin & Papavero, 1970: 29 (cat.; *Lastauroides*); Artigas & Papavero, 1988b: 212 (in key; n. comb.), 259, fig. 206. Type, IOC.
atriapex (Carrera & Papavero), 1962: 52 (*Lastauroides*). Type-locality: Brazil, São Paulo, Araçatuba (rio Jacaretinga). Distr. – Brazil (São Paulo). Refs. – Martin & Papavero, 1970: 29 (cat.; *Lastauroides*); Artigas & Papavero, 1988b: 213 (n. comb.). Type, MZUSP.
carrerai Artigas & Papavero, 1988b: 213, 254, figs. 183-189. Type-locality: Brazil, Espírito Santo, Santa Teresa. Distr. – Brazil (Espírito Santo). Type, MZUSP.
hirtuosus (Wiedemann), 1821: 227 (*Dasypogon*). Type-locality: “Brazil” [Olfers; cf. Papavero, 1971e: 69]. Distr. – Brazil (Rio de Janeiro, São Paulo). Refs. – Wiedemann, 1828: 402 (*Dasypogon*); Walker, 1854: 443 (*Dasypogon*); Schiner, 1866: 679 (*Dasypogon*); Williston, 1891: 67 (cat.; *Dasypogon*); Kertész, 1909: 128 (cat.; *Dasypogon*); Carrera, 1949: 97, fig. 27 (*Lastauroides*; n. comb.), 1958a: 146 (*Lastauroides*); Carrera & d’Andretta, 1961: 69 (*Lastauroides*; prey); Carrera & Papavero, 1962: 53; Hull, 1962 (1): 241, figs. 545, 1074, 1083 (*Lastauroides*); Martin & Papavero, 1970: 29 (cat.; *Lastauroides*); Artigas & Papavero, 1988b: 213 (n. comb.), 151, fig. 157. Syntypes, BZM [no. 541; cf. Schumann, 1973: 116].
hirtosus Bromley, 1946: 107 (cat.; *Dasypogon*), error.
lanei (Carrera), 1949: 110, figs. 28, 51, 83, 108 (*Lastaurax*). Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. – Brazil (Rio de Janeiro). Refs. – Hull, 1962 (1): 240, figs. 1071, 1080 (*Lastaurax*); Martin & Papavero, 1970: 28 (cat.; *Lastaurax*); Artigas & Papavero, 1988b: 214 (n. comb.). Type, MZUSP.
melaleucus (Schiner), 1868: 165 (*Dasypogon*). Type-locality: “South America”. Refs. – Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 230 (cat.; *Dasypogon*); Carrera, 1949: 98 (*Lastauroides*); Hull, 1962 (1): 241 (*Lastauroides*); Martin & Papavero, 1970: 29 (cat.; *Lastauroides*). Type, NMW.
melanogaster (Wiedemann), 1821: 215 (*Dasypogon*). Type-locality: “Brazil”. Distr. – Brazil (Rio de Janeiro, Paraná, Santa Catarina), Argentina (Misiones). Refs. – Wiedemann, 1828: 368 (*Dasypogon*); Macuart, 1834: 294 (*Dasypogon*), 1838: 35 (*Dasypogon*); Walker, 1849: 366 (*Dasypogon*), 1854: 430 (*Dasypogon*); Schiner, 1866: 675 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 130 (cat.; *Dasypogon*); Bromley, 1946: 86 (*Diogmites*); Carrera, 1947d: 267 (*Diogmites*) (preys), 1949: 86, figs. 22, 44, 84, 113, 144, 151 (n. comb.), 1960a: 158; Carrera & Papavero, 1962: 46; Hull, 1962 (1): 235, figs. 156B, 541, 1068, 1077; Martin & Papavero, 1970: 28 (cat.); Artigas & Papavero, 1988b: 214. Syntypes, NMW.
grandis Macquart, 1846: 63, pl. 6, fig. 8. Type-locality: “Brazil”. Refs. – Walker, 1854: 436 (*Dasypogon*); Williston, 1891: 128 (cat.; *Dasypogon*); Kertész, 1909: 128 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Diogmites*); Hull, 1962 (1): 228 (*Dasypogon*). Type, lost.
rapax Walker, 1851: 88 (*Dasypogon*). Type-locality: “South America”. Refs. – Walker, 1854: 430 *Dasypogon*; Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 131 (cat.; *Dasypogon*). Type, BMNH.
mixtus (Carrera), 1949: 101, fig. 55 (*Lastauroides*). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. – Brazil (São Paulo). Refs. – Hull, 1962 (1): 241 (*Lastauroides*); Martin & Papavero, 1970: 29 (cat.; *Lastauroides*); Artigas & Papavero, 1988b: 214 (n. comb.), 251, fig. 158. Type, MZUSP.
modestus Carrera, 1949: 103, fig. 56 (*Lastauroides*). Type-locality: Brazil, São Paulo, Campos do Jordão. Refs. – Carrera & d’Andretta, 1961: 69 (*Lastauroides*; preys); Hull, 1962 (1): 241 (*Lastauroides*); Martin & Papavero, 1970: 29 (cat.; *Lastauroides*); Artigas & Papavero, 1988b: 214 (syn.). Type, MZUSP.
niger (Carrera), 1949: 99 (*Lastauroides*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Distr. – Brazil (Rio de Janeiro, São Paulo). Refs. – Hull, 1962 (1): 241 (*Lastauroides*); Artigas & Papavero, 1988bg: 214 (n. comb.). Type, MZUSP.
albomarginatus Carrera, 1949: 98 (*Lastauroides*). Type-locality: Brazil, São Paulo, Paranaípacaba (as Alto da Serra). Refs. – Hull, 1962 (1): 241 (*Lastauroides*); Martin & Papavero, 1970: 29 (cat.; *Lastauroides*); Artigas & Papavero, 1988b: 214 (syn.). Type, MZUSP.
tauauana Artigas & Papavero, 1988: 214. Type-locality: Brazil, Espírito Santo, Itirapina. Distr. – Brazil (Espírito Santo). Type, MZUSP.
tenebrosus Carrera, 1949: 87, figs. 21, 73. Type-locality: Brazil, Bahia, Jequié. Distr. – Brazil (Bahia). Refs. – Hull, 1962 (1): 235; Martin & Papavero, 1970: 28 (cat.). Type, MZUSP.

Genus **Phonicocleptes** Lynch Arribálzaga, 1881

Phonicocleptes Lynch Arribálzaga, 1881: 18. Type-species, *Phonicocleptes busiris* Lynch Arribálzaga, 1881 (mon.).

busiris Lynch Arribálzaga, 1881: 21. Type-locality: Argentina, Buenos Aires, Chacabuco. Distr. – Uruguay, Argentina. Refs. – Carrera, 1949: 59, 1953a: 273; Hull, 1962 (1): 235, figs. 65, 574, 1057, 1066, 1794, 1799; Martin & Papavero, 1970: 28 (cat.); Artigas & Papavero, 1988b: 259, fig. 208, 260, fig. 210–214.

longipes (Macquart), 1838: 30 (1839: 146), pl. 2, fig. 4 (*Dasypogon*). Type-locality: Brazil, northern state of São Paulo. Distr. – Peru (Loreto), Brazil (São Paulo, Paraná). Refs. – Walker, 1854: 440 (*Dasypogon*); Williston, 1891: 70 (cat.); *Microstylum*; Kertész, 1909: 152 (cat.; *Microstylum*); Bromley, 1946: 108 (cat.; *Archilestris*); Hull, 1962: 141 (*Archilestris*); Martin & Papavero, 1970: 18 (*Archilestris*); Lamas M., 1972: 307 (cat.); Artigas & Papavero, 1988b: 260, fig. 209 e 215. Type, MNHNP.

spectans Walker, 1851: 88 (*Dasypogon*). Type-locality: unknown. Refs. – Walker, 1854: 484 (*Dasypogon*); Kertész, 1909: 132 (cat.; *Dasypogon*); Carrera, 1949: 60 (n. comb.); Hull, 1962 (1): 235; Martin & Papavero, 1970: 28 (cat.); Papavero, 1971d: 20 (n. syn.). Type, BMNH.

langei Carrera, 1947d: 268 (descr. and prey). Type-species, Brazil, Paraná, Vila Velha. Refs. – Carrera, 1949: 61, figs. 66, 72, 122; Carrera & d'Andretta, 1953: 66; Carrera & Papavero, 1962: 47; Hull, 1962 (1): 235; Martin & Papavero, 1970: 28 (cat.); Papavero, 1971d: 20 (n. syn.). Type, Mus. Paranaense.

Tribe **Megapodini** Carrera, 1949

Acanthocneminae Hermann, 1912: 9 (in Laphriinae).

Megapodini Carrera, 1949: 5. Type-genus, *Megapoda* Macquart, 1834.

Megapodininae Hull, 1962 (1): 6.

Megapodinae Hull, 1962 (1): 420.

Megapodina Papavero, 1975: 228.

Cyrtophriina Papavero, 1975: 229. Type-species, *Cyrtophrys* Loew, 1851.

Senobasiina Papavero, 1975: 229. Type-genus, *Senobasis* Macquart, 1838.

Senobasina Sabrosky, 1999: 279, corrected spelling.

Ref. – Papavero, 1975 (rev.).

Genus **Cyrtophrys** Loew, 1851

Dasypogon, subg. **Cyrtophrys** Loew, 1851: 3. Type-species, *Dasypogon (Cyrtophrys) attenuatus* Loew, 1851 (orig. des.).

Myolestes Brèthes, 1904: 338. Type-species, *Myolestes lynchii* Brèthes, 1904 (mon.). Refs. – Arnaud, 1974 (erroneously as a valid genus); Papavero, 1975: 229 (syn.).

Myiolestes Kertész, 1909: 123 (preocc. Bonaparte, 1850), emend.

Mirolestes Curran, 1935: 1 (unjustified *nom. nov.* for *Myolestes* Brèthes, 1904). Type-species, *Myolestes lynchii* Brèthes, 1904 (aut.).

Ref. – Papavero, 1975: 231 (key to spp.).

albimanus (Carrera), 1949: 114, fig. 94 (*Mirolestes*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Distr. – Brazil (Rio de Janeiro). Ref. – Hull, 1962 (1): 252 (*Mirolestes*); Martin & Papavero, 1970: 31 (cat.; *Mirolestes*); Papavero, 1975: 231, fig. 17 (n. comb.); Artigas & Papavero, 1988b: 232, figs. 58, 65–69. Type, MZUSP.

attenuatus (Loew), 1851: 3 (*Dasypogon (Cyrtophrys)*). Type-locality: “Brazil” (probably Rio de Janeiro, Nova Friburgo; cf. Papavero, 1971e: 87). Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Kertész, 1909: 119 (cat.); Bromley, 1946: 106 (cat.); Carrera, 1949: 121; Hull, 1962 (1): 250, figs. 180, 552, 566, 1091, 1100, 1893 (*Lastauroides*); Martin & Papavero, 1970: 30 (cat.); Papavero, 1975: 232, figs. 1, 10, 18, 44, 50–52, 77, 93–95 (n. comb.); Artigas & Papavero, 1988b: 238, fig. 94. Syntypes, BZM [no. 10133; cf. Schumann, 1973: 89].

tenuis Walker, 1851: 86 (*Dioctria*). Type-locality: “South America”. Refs. – Walker, 1854: 387 (*Dioctria*); Williston, 1891: 69 (cat.; *Dioctria*); Kertész, 1909: 114 (cat.; *Dioctria*); Hull, 1962 (1): 29 (*Dioctria*); Martin & Papavero, 1970: 31 (cat.; *Mirolestes*); Papavero, 1975: 232 (syn.). Martin & Papavero, 1970: 18 (cat.; *Mirolestes*).

atennatus Williston, 1891: 74 (cat.; as from Chile, in error), error.

barbiellinii Curran, 1935: 1 (*Mirolestes*). Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 107 (cat.; *Mirolestes*);

Carrera, 1949: 113, figs. 10, 62, 96, 120, 132, 134, 181; Carrera & Vulcano, 1961: 69 (prey); Carrera & Papavero, 1962: 55; Hull, 1962 (1): 252 (*Mirolestes*); Martin & Papavero, 1970: 31 (cat.); Papavero, 1975: 232 (syn.). Type, AMNH.

facialis (Curran), 1935: 1 (in key), 2 (*Mirolestes*). Type-locality: Brazil, São Paulo. Distr. – Brazil (Rio de Janeiro, São Paulo, Santa Catarina). Refs. – Bromley, 1946: 107 (cat.; *Mirolestes*); Hull, 1962 (1): 252, figs. 71, 1821, 1829 (*Mirolestes*); Martin & Papavero, 1970: 31 (cat.; *Mirolestes*); Papavero, 1975: 232, figs. 19, 96-98 (n. comb.). Type, AMNH.

pleuralis Curran, 1935: 1 (in key), 3 (*Mirolestes*). Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 107 (cat.; *Mirolestes*); Carrera, 1949: 233 (as syn. of *lynchii* Brèthes, in error); Hull, 1962 (1): 252 (*Mirolestes*; as syn. of *lynchii* Brèthes, 1901); Martin & Papavero, 1970: 31 (cat.; *Mirolestes*, syn.). Type, AMNH.

fascialis Carrera, 1949: 116, figs. 59, 95 (*Mirolestes*), 1953a: 274 (*Mirolestes*); Carrera & Papavero, 1962: 54, error.

lynchii (Brèthes), 1901: 338 (*Myolestes*). Type-locality: Argentina, Tucumán. Distr. – Venezuela (Carabobo), Brazil (Mato Grosso, São Paulo, Santa Catarina), Argentina (Tucumán). Refs. – Kertész, 1909: 123 (cat.; *Myolestes*); Curran, 1935: 1 (in key) (*Mirolestes*); Carrera, 1949: 116, figs. 59, 95 (*Mirolestes*); Carrera & Papavero, 1962: 54; Hull, 1962 (1): 252, fig. 530, 1092, 1101 (*Mirolestes*); Carrera & Machado-Allison, 1963: 235; Martin & Papavero, 1970: 31 (cat.; *Mirolestes*); Papavero, 1975: 235, figs. 20, 99-101 (n. comb.).

sp. Papavero, 1975: 236 (Venezuela, Carabobo, Valle Seco).

Genus **Deromyia** Philippi, 1865

Deromyia Philippi, 1865: 705. Type-species, *Deromyia gracilis* Philippi, 1865 (sub. des., Williston, 1901: 311, as of Rondani, in error) = *fuscipennis* (Blanchard, 1852).

Ref. – Papavero, 1975: 236 (key to spp.).

fuscipennis (Blanchard), 1852: 371, pl. 1, figs. 6a-b (*Gonypus*). Type-locality: Chile, Santiago. Neotype-locality: Chile, Curicó, Estero La Jaula. Distr. – Chile (Arauco, Cautín, Concepción, Colchagua, Curicó, Linares, Malleco, Ñuble, O'Higgins, Santiago, Talca, Valparaíso), Argentina (Río Negro). Refs. – Walker, 1855: 771 (*Leptogaster*); Kertész, 1909: 116 (cat.); Bromley, 1932: 267 (*Deromyia*); Stuardo Ortiz, 1946: 82 (cat.); Carrera & Papavero, 1962: 61; Hull, 1962 (1): 250; Martin, 1968e: 3 (cat., *Leptogaster*); Artigas, 1970a: 113, figs. 173-178, 441, 1971: 11, figs. 11-12; Martin & Papavero, 1970: 31 (cat.); Papavero, 1975: 237, figs. 2, 11, 21, 53-55, 78, 102-104; Artigas & Papavero, 1988b: 232, fig. 59. Type, lost. NT DZUC [des. by Artigas, 1970a: 114].

nigrescens Blanchard, 1852: 371 (*Leptogaster*), error.

fascipennis Philippi, 1865: 705 (*Leptogaster*), 706; Williston, 1891: 75 (cat.), error.

gracilis Philippi, 1865: 706, pl. 26, figs. 29, 29a. Type-locality: “Chile”. Refs. – Williston, 1891: 75 (cat.); Kertész, 1909: 116 (cat.); Stuardo Ortiz, 1946: 82 (cat.); Carrera, 1949: 122; Artigas, 1961: 5, figs. 2, 4, 6; Hull, 1962 (1): 250, figs. 532, 1089, 1098. Type, lost.

nigriventris (Bigot), 1878: 427 (*Lochites*). Type-locality: “Chile”. Distr. – Chile (Arauco, Malleco, Ñuble). Refs. – Williston, 1891: 75 (cat.; *Senobasis*); Kertész, 1909: 125 (cat.; *Lochites*); Brèthes, 1925: 105 (*Lochitomyia*); Stuardo Ortiz, 1946: 82 (cat.; *Senobasis*); Hull, 1962 (1): 233 (*Blepharepium*), 422 (*Senobasis*); Martin & Papavero, 1970: 31 (as syn. of *fuscipennis* Blanchard, in error); Artigas, 1970a: 116, figs. 179-180, 446, 1971: 11, figs. 13-14; Papavero, 1975: 238. Type, OXF.

negriventris Artigas, 1970a: 113 (in key), error.

Genus **Megapoda** Macquart, 1834

Megapoda Macquart, 1834: 288. Type-species, *Megapoda cyanea* Macquart, 1834 (mon.) = *Megapoda labiata* (Fabricius, 1805).

labiata (Fabricius), 1805: 160 (*Laphria*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Guyana, Brazil (Amazonas, Pará, Mato Grosso, Paraíba, Bahia, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo). Refs. – Wiedemann, 1821: 232 (*Laphria*), 1828: 499 (*Laphria*); Macquart, 1838: 59 (1839: 175) (n. comb.); Walker, 1849: 371 (*Laphria*), 1855: 509; Schiner, 1868: 690; Wulp, 1882b: 103; Roeder, 1887: 76; Williston, 1889: 256, 1891: 77 (cat.); Kertész, 1909: 161 (cat.); Hermann, 1912: 18, fig. 9; Bromley, 1946: 108 (cat.); Carrera, 1949: 7, fig. 1, 1952c: 62, figs. 13-20, 1960a: 157; Hull, 1962 (1): 424, figs. 247, 635, 1275, 1284, 1564, 1568, 1816, 2111, 2139; Martin & Papavero, 1970: 55 (cat.); Papavero, 1975: 246, figs. 4, 14, 26, 46, 59-61, 80, 108-110; Artigas & Papavero, 1988b: 232, fig. 60, 234, figs. 71-75, 238, fig. 95; Castillo, Jerez & Artigas, 1994 (egg chorion); Geller-Grimm, 1999: 210 [specimen in SENCK]. Type, COPN [cf. Zimšen, 1964: 468].

rufimanus Perty, 1833: 181, pl. 36, fig. 6 (*Dasyopogon*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (as “Sebastianopolis”). Ref. – Kertész, 1909: 161 (*Megapoda*). Type, lost.

cyanea Macquart, 1834: 288. Type-locality: “Brazil”. Ref. – Macquart, 1838: pl. 5, fig. 3. Type, lost.

Genus **Pronomopsis** Hermann, 1912

Pronomopsis Hermann, 1912: 18. Type-species, *Pronomopsis chalybea* Hermann, 1912 (orig. des.).

Ref. – Papavero, 1975: 249 (key to spp.).

chalybea Hermann, 1912: 19, fig. 10, pl. 1, fig. 4. Type-locality: Argentina, Mendoza. Distr. – Argentina (Mendoza). Refs. – Carrera, 1949: 8, 1952c: 60; Hull, 1962 (1): 423; Artigas, 1964: 8, 1970a: 253, fig. 411; Martin & Papavero, 1970: 56 (cat.); Papavero, 1975: 249. Type, ZSBS.

pseudorubripes Lamas M., 1972: 312, fig. 4. Type-locality: Peru, Cuzco. Distr. – Peru (Cuzco). Type, Departamento de Zoología y Entomología, Universidad Nacional “San Antonio Abad”, Cuzco, Peru.

rubripes Hermann, 1912: 20. Type-locality: Peru, Puno, Juliaca, 4000 m. Distr. – Peru (Puno), Chile (Antofagasta). Refs. – Hull, 1962 (1): 423, figs. 240, 660, 1238, 1247, 2105, 2144; Artigas, 1964: 4, figs. 2, 4-6, 9-13; Artigas, 1970a: 254, figs. 213, 215-219, 412, 1971: 31, figs. 73-74; Martin & Papavero, 1970: 56 (cat.); Lamas M., 1972: 312 (cat.); Papavero, 1975: 250, figs. 7, 27, 62-64, 81, 111-113; Artigas & Papavero, 1988b: 235, figs. 76-81TP ZSBS.

talabrensis Artigas, 1964: 6, figs. 1, 3, 8, 12, 14. Type-locality: Chile, Antofagasta, Talabre. Distr. – Chile (Antofagasta). Refs. – Artigas, 1970a: 256, fig. 486, 1971: 32, figs. 71-72; Martin & Papavero, 1970: 56 (cat.); Papavero, 1975: 251. Type, DZUC.

Genus **Pseudorus** Walker, 1851

Pseudorus Walker, 1851: 103. Type-species, *Pseudorus piceus* Walker, 1851 (mon.).

Ampyx Walker, 1855: 564 (preocc. Dalman, 1827). Type-species, *Ampyx varipennis* Walker, 1855 (mon.) = *Pseudorus distendens* (Wiedemann, 1838). Ref. – Papavero, 1975: 251 (syn.).

Doryclus Jaennicke, 1867: 366. Type-species, *Asilus distendens* Wiedemann, 1828 (orig. des.). Ref. – Papavero, 1975: 251 ((syn.)). *Pseudoryclus* Carrera, 1952c: 61, 75. Type-species, *Megapoda rufiventris* Roeder, 1887 (orig. des.). Ref. – Papavero, 1975: 251 (syn.).

Ref. – Papavero, 1975: 251 (key to spp.).

bicolor Bellardi, 1861: 111 (11), pl. 1, fig. 20. Type-locality: Mexico, Veracruz, Playa Vicente. Distr. – Mexico (Nayarit, Michoacán, Oaxaca, Guerrero, Jalisco, Veracruz, Yucatán). Refs. – Osten Sacken, 1887: 183; Williston, 1889: 256, 1901: 315; Aldrich, 1905: 267 (cat.); Kertész, 1909: 161 (cat.); Hermann, 1912: 14; Carrera, 1949: 15; Carrera & d’Andretta, 1950: 162, fig. 13; Carrera, 1952c: 76 (*Pseudoryclus*); Hull, 1962 (1): 428 (*Pseudoryclus*); Martin & Papavero, 1970: 57 (cat.; *Pseudoryclus*); Papavero, 1975: 254, figs. 35, 86, 114-116; Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipteral.). Type, TORO.

conspectus Tomasovic in Tomasovic & Braet, 2001: 380, figs. 1-2, 9-11 [with notes on a mimetic complex involving Braconidae and Ichneumonidae (Hymenoptera), Reduviidae (Hemiptera) and Arctiidae Ctenuchinae (Lepidoptera)]. Type-locality: French Guiana, Montagne de Kaw, Relais Patawa (40°32'42.20"N, 52°09'09.19"W). Distr. – French Guiana. Type, FUSA Gembloux.

dandrettae Carrera, 1949: 15 (as *d’andrettae*). Type-locality: Brazil, São Paulo, Itaporanga (Nucleo Barão de Antonina). Distr. – Brazil (Rio de Janeiro, São Paulo, Santa Catarina). Refs. – Carrera, 1952c: 82; Hull, 1962 (1): 430; Martin & Papavero, 1970: 56 (cat.); Papavero, 1975: 256, figs. 8, 16, 37, 68-70, 88; Geller-Grimm, 1997: 459, figs. 2a-g (descr. of male). Type, MZUSP.

dimidiatus Oldroyd, 1964: 5 (as *dimiata*, error), 9, figs. 1-2, 6.3. Type-locality: Brazil, Amazonas. Distr. – Venezuela, Peru, Brazil (Amazonas, Pará). Refs. – Martin & Papavero, 1970: 56 (cat.); Papavero, 1975: 256, figs. 42, 71-731; Fisher, 1984: 28. Type, BMNH.

distendens (Wiedemann), 1828: 571 (*Asilus*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Mexico to Paraguay, southern Brazil and Argentina (Misiones, Tucumán, Buenos Aires). Refs. – Schiner, 1866: 690 (*Asilus*); Jaennicke, 1867: 366, pl. 44, fig. 3 (*Doryclus*); Roeder, 1887: 76 (*Doryclus*); Williston, 1889: 256 (*Doryclus*), 1891: 77 (cat.; *Doryclus*); Aldrich, 1905: 268 (cat.; *Doryclus*); Kertész, 1909: 161 (cat.; *Doryclus*); Hermann, 1912: 12 (*Doryclus*); Curran, 1942: 56 (*Doryclus*); Bromley, 1946:a 108 (cat.; *Doryclus*); Carrera, 1949: 9 (*Doryclus*); Bromley, 1951: 17 (in key; *Doryclus*); Carrera, 1952c: 80, figs. 23-24, 27 (*Doryclus*; *cyaniventris*, *varipennis*, *latipes*, *guentherii* and *panamensis* as syn.); Hull, 1962 (1): 427, figs. 236, 647, 1278, 1287, 2104, 2118, 2169 (*Doryclus*); Martin & Papavero, 1970: 56 (cat.; *Doryclus*); Lamas M., 1972: 312 (cat.); Papavero, 1975: 257, figs. 9, 15, 36, 74-76, 87, 117-119; Knutson,

- 1976: 509 (*Doryclus*; pupal case); Fisher, 1984: 28; Artigas & Papavero, 1988b: 236, figs. 82-86, 238, fig. 97; Dennis & Knutson, 1988: 860. Type, SENCK [cf. Geller-Grimm, 1999: 208].
- crassitarsis* Macquart, 1846: 198 (1846: 70), pl. 7, figs. 11, 11a. Type-locality: "Brazil". Refs. – Walker, 1855: 509 (*Megapoda*); Kertész, 1909: 162 (cat.; *Doryclus*); Curran, 1942: 56 (*Doryclus*); Bromley, 1946: 108 (cat.; *Doryclus*); Carrera, 1949: 10 (*Doryclus*); Bromley, 1951: 17 (in key; *Doryclus*); Carrera, 1952c: 81, figs. 21-22, 25-26, 28-29 (*Doryclus*); Hull, 1962 (1): 427 (*Doryclus*); Martin & Papavero, 1970: 56 (cat.; *Doryclus*). Type, lost.
- cyaniventris* Macquart, 1846: 199 (1846: 71), pl. 7, fig. 12 (*Megapoda*). Type-locality: "Mexico". Refs. – Walker, 1855: 508 (*Megapoda*); Roeder, 1887: 76 (*Doryclus*); Bromley, 1951: 17 (in key; *Doryclus*); Carrera, 1952c: 78; Martin, 1965a: 27; Papavero & Ibáñez Bernal, 2001: 84 (hist. Mexican dipterol.). Type, MNHNP.
- varipennis* Walker, 1855: 564 (*Ampyx*). Type-locality: Brazil, Pará, Santarém. Ref. – Osten Sacken, 1887: 182 (*Doryclus*); Curran, 1942: 56 (*Doryclus*); Carrera, 1949: 11 (*Doryclus*); Bromley, 1951: 17 (in key; *Doryclus*). Type, BMNH.
- latipes* Wulp, 1870: 216, pl. 9, figs. 7-12 (*Doryclus*). Type-locality: "Surinam". Refs. – Curran, 1942: 56 (*Doryclus*); Bromley, 1951: 17 (in key; *Doryclus*); Martin & Papavero, 1970: 56 (cat.; *Doryclus*). Type, ZMA.
- guentherii* Lynch Arribálzaga, 1882: 186 (*Doryclus*). Type-locality: Argentina, Buenos Aires. Refs. – Williston, 1891: 77 (*Megapoda*); Kertész, 1909: 161 (cat.; *Megapoda*); Hermann, 1912: 12 (*Doryclus*); Bromley, 1946: 108 (cat.; *Doryclus*); Carrera, 1949: 13 (*Doryclus*); Bromley, 1951: 17 (in key; *Doryclus*); Martin & Papavero, 1970: 56 (cat.; *Doryclus*). Type, MACN.
- panamensis* Curran, 1942: 56 (*Doryclus*). Type-locality: Panama, Canal Zone, Corozal. Ref. – Bromley, 1951: 17 (in key; *Doryclus*); Hull, 1962 (1): 427 (*Doryclus*); Martin & Papavero, 1970: 56 (cat.; *Doryclus*). Type, AMNH.
- carrerai* Bromley, 1951: 17 (in key), 18 (*Doryclus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Refs. – Hull, 1962 (1): 427 (*Doryclus*); Martin & Papavero, 1970: 56 (cat.; *Doryclus*); Papavero, 1975: 258 (syn.). Type, AMNH.
- plaumanni* Bromley, 1951: 18 (*Doryclus*). Type-locality: Brazil, Santa Catatrina, Nova Teutônia. Refs. – Hull, 1962 (1): 427 (*Doryclus*); Martin & Papavero, 1970: 56 (cat.; *Doryclus*); Papavero, 1975: 258 (syn.). Type, AMNH.
- distends* Notario, Michela, Fiorentino & Castresana, 2000: 17, error ("emergieron de trozas en las que sólo se encontraron *B[rasilianus] lacordairei* [(Gahan, 1892)], *B[rasilianus] murinus* [(Gahan, 1892)] [Coleoptera, Cerambycidae] y *C[hrysobothris] holochalcea?* [Burmeister, 1872] [Coleoptera, Buprestidae]").
- hermanni** Carrera, 1949: 14. Type-locality: unknown [Brazil]. Distr. – Brazil (Santa Catarina). Refs. – Carrera, 1852c: 61; Oldroyd, 1964: 8, figs. 4, 6.1; Hull, 1962 (1): 430; Martin & Papavero, 1970: 56 (cat.); Papavero, 1975: 260, figs. 38, 49, 89, 120-122. Lectotype NMW [des. by Oldroyd, 1954a: 9].
- piceus* Walker of Schiner, 1866: 703 and Hermann, 1912: 13, fig. 5, misident.
- holcephalinus** Papavero, 1975: 261, figs. 39, 47, 90. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (Santa Catarina). Type, MZUSP.
- dandrettae* Carrera of Oldroyd, 1964: 5, 10, figs. 3, 6.3, misident.
- martini** Papavero, 1975: 262, figs. 40, 91, 123-125. Type-locality: Mexico, Puebla, south of Matamoros. Distr. – Mexico (Morelos, Puebla). Ref. – Artigas & Papavero, 1988b: 232, figs. 61-62. Type, MZUSP.
- mexicanus** (Bromley), 1951: 17 (*Doryclus*). Type-locality: Mexico, Sonora, Agua Prieta. Distr. – Mexico (Sonora). Refs. – Hull, 1962 (1): 427 (*Doryclus*); Papavero, 1975: 263 (n. comb.). Type, USNM.
- piceus** Walker, 1851: 103, pl. 4, figs. 5, 5a. Type-locality: "South America". Distr. – Brazil (Espírito Santo). Refs. – Walker, 1854: 493; Williston, 1889: 256, 1891: 77 (cat.); Kertész, 1909: 161 (cat.); Hermann, 1912: 13; Bromley, 1946: 108 (cat.); Carrera, 1949: 15; Hull, 1962 (1): 430, text-fig. 29, figs. 235, 668, 1279, 1288, 2168; Oldroyd, 1964: 7, figs. 5, 6.1; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 254, figs. 48, 92. Type, BMNH.
- rufiventris** (Roeder), 1887: 78 (*Megapoda*). Type-locality: Brazil, Amazonas, Maués. Distr. – Peru (Huancayo), Brazil (Amazonas). Refs. – Williston, 1891: 77 (cat.; *Megapoda*); Kertész, 1909: 161 (cat.; *Megapoda*); Hermann, 1912: 18 (*Megapoda*); Bromley, 1946: 108 (cat.; *Megapoda*); Carrera, 1952c: 75 (*Pseudoryclus*); Hull, 1962 (1): 428 (*Pseudoryclus*); Martin & Papavero, 1970: 57 (cat.; *Pseudoryclus*); Lamas M., 1972: 312 (cat.; *Pseudoryclus*); Papavero, 1975: 264, fig. 43 (n. comb.).

Genus *Senobasis* Macquart, 1838

- Senobasis** Macquart, 1838: 52 (1839: 168). Type-species, *Senobasis analis* Macquart, 1838 (sub. des., Bromley, 1934: 332). *Stenobasis* Agassiz, 1846: 338, emend.
- Astylium* Rondani, 1850: 185. Type-species, *Astylium dorsiger* Rondani, 1850 (orig. des.).
- Lochites* Schiner, 1866: 655 (key, 671 (12868: 163; second erection of genus) (preocc. Cabanis & Heine, 1860). Type-species, *Laphria ornata* Wiedemann, 1819 (orig. des.).
- Astylum* Kertész, 1909: 83, emend.
- Stenobasis* Kertész, 1909: 123, emend.
- Lochitomyia* Brèthes, 1925: 105 (nom. nov. for *Lochites* Schiner, 1866). Type-species, *Laphria ornata* Wiedemann, 1819 (aut.).

Ref. – Papavero, 1975: 265 (key to spp.).

- aedon** (Walker), 1849: 343 (*Dasypogon*). Type-locality: “Venezuela”. Distr. – Colombia, Venezuela. Refs. – Walker, 1854: 432 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 83 (*Stichopogon*); Bezzi, 1910c: 155 (*Neopogon*); Hull, 1962 (1): 106 (*Stichopogon*); Martin & Papavero, 1970: 5 (cat.; *Stichopogon*); Papavero, 1975: 269, figs. 126, 128-131 (n. comb.). Type, BMNH.
- staurophora** Schiner, 1868: 164 (*Lochites*). Type-locality: “Colombia”. Lectotype-locality: “Venezuela” (Papavero, 1975: 269). Refs. – Kertész, 1909: 125 (cat.; *Lochites*); Williston, 1891: 75 (cat.; n. comb.); Kertész, 1909: 125 (cat.; *Lochites*); Carrera, 1949: 21; Hull, 1962 (1): 422; Carrera & Machado-Allison, 1963: 239; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 269 (syn.). Lectotype NMW [des. by Papavero, 1975: 269].
- almeidai** Carrera, 1946c: 121. Type-locality: Brazil, São Paulo, Araraquara. Distr. – Brazil (São Paulo). Refs. – Carrera, 1949: 18, 1952c: 66; Hull, 1962 (1): 422; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 288. Type, MZUSP.
- analis** Macquart, 1838: 53 (1839: 169), pl. 5, figs. 1, 1a, 1b. Type-locality: French Guiana, Mana river (as “Brésil, Lamana”, in error). Distr. – Colombia, Guyana, French Guiana, Peru, Brazil (Amazonas, Amapá). Refs. – Walker, 1854: 500; Williston, 1891: 74 (cat.); Kertész, 1909: 123 (cat.; *Stenobasis*); Curran, 1934c: 332; Bromley, 1946: 106 (cat.); Carrera, 1949: 19; Hull, 1962 (1): 422, figs. 675, 1276, 1286, 2107, 2131, 2165; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 271, figs. 28, 131-133; Fisher, 1984: 28. Type, MNHNP.
- tibialis** Curran, 1934c: 332. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Refs. – Carrera, 1949: 27; Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 271 (syn.). Type, AMNH.
- apicalis** (Schiner), 1867: 370 (*Lochites*). Type-locality: “Brazil” [Helmreich [sic]] (probably Minas Gerais, where Helmreich stayed most of his time); cf. Papavero, 1973d: 295]. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Williston, 1891: 75 (n. comb.); Kertész, 1909: 124 (cat.; *Lochites*); Williston, 1891: 75; Bromley, 1946: 106 (cat.); Carrera, 1949: 18, 1952c: 68; Hull, 1962 (1): 422; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 271, fig. 85; Artigas & Papavero, 1988b: 237, figs. 87-93. Type, NMW.
- lopesi** Carrera, 1949: 19, fig. 165. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Refs. – Carrera, 1952c: 65; Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 271 (syn.). Type, MZUSP.
- bromleyana** Carrera, 1949: 22, fig. 30. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. – Brazil (Goiás, Minas Gerais, São Paulo). Refs. – Carrera, 1952c: 67; Carrera & Vulcano, 1961: 70 (preys); Hull, 1962 (1): 422; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 374, figs. 134-136; Artigas & Papavero, 1988b: 238, fig. 96. Type, MZUSP.
- claripennis** (Schiner), 1867: 369 (*Lochites*). Type-locality: “Brazil (Bescke)” (most probably Nova Friburgo, state of Rio de Janeiro; cf. Papavero, 1971b: 87-89). Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Santa Catarina). Refs. – Williston, 1891: 75 (cat.); Kertész, 1909: 125 (cat.; *Lochites*); Bromley, 1946: 106 (cat.; n. comb.); Carrera, 1949: 25, fig. 164 (part), 1952c: 67 (part), 1960a: 157; Carrera & Vulcano, 1961: 70 (preys); Hull, 1962 (1): 422; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 276, figs. 65-67, 82, 143-145; Artigas & Papavero, 1988b: 232, figs. 63-64. Type, NMW.
- clavigera** (Rondani), 1850: 185, pl. 4, fig. 8 (*Astylium*). Type-locality: “Venezuela”. Distr. – Venezuela, Peru (Huánuco). Refs. – Osten Sacken, 1891: 417 (*Astylium*); Kertész, 1909: 83 (cat.; *Astylium*); Carrera & d’Andretta, 1953: 65; Hull, 1962 (1): 422; Martin & Papavero, 1970: 57 (cat.); Lamas M., 1972: 312 (cat.); Papavero, 1975: 277. Type, TORO.
- clavigeroides** Papavero, 1975: 277. Type-locality: Ecuador, Santo Domingo, 500 m. Distr. – Ecuador, Peru. Type, MZUSP.
- clavigera* Rondani of Carrera, 1952c: 70, misident.
- corsair** Bromley, 1951b: 12. Type-locality: Panama, Río Indio. Distr. – Costa Rica, Panama, Colombia. Refs. – Hull, 1962 (1): 422; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 277, figs. 29, 45, 83, 149-151; Shelly, 1985a: 253 (reproductive behaviour). Type, USNM.
- flukei** Carrera, 1952c: 68, fig. 7. Type-locality: Ecuador, Manabí, Palmar, 200 m. Distr. – Ecuador, Peru. Refs. – Hull, 1962 (1): 422; Martín & Papavero, 1970: 58 (cat.); Papavero, 1975: 281, figs. 30, 152-154. Type, MZUSP.
- frosti** Bromley, 1951b: 12. Type-locality: Panama, Pecora. Distr. – Panama. Refs. – Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 282. Type, USNM.
- gyrophora** (Schiner), 1868: 164, pl. 2, fig. 10 (*Lochites*). Type-locality: “Brazil (Coll. Winthem)” [probably from Rio de Janeiro]. Distr. – Brazil (Rio de Janeiro, Santa Catarina). Refs. – Williston, 1891: 75 (cat.); Kertész, 1909: 125 (*Lochites*); Bromley, 1946: 106 (cat.; n. comb.); Carrera, 1949: 29, figs. 2, 32, 166; Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 282, figs. 84, 155-157. Type, NMW.
- lanei** Carrera, 1949: 23, fig. 3. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Carrera, 1952c: 73, 1960a: 157; Carrera & Vulcano, 1961: 69 (preys); Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 282, figs. 40, 158-160. Type, MZUSP.
- boraceana** Carrera, 1952c: 74. Type-locality: Brazil, São Paulo, Salesópolis (Estação Biológica de Boracéia). Refs. – Carrera, 1958a: 146; Hull, 1962 (1): 422; Martin & Papavero, 1970: 57 (cat.); Papavero, 1975: 282 (syn.). Type, MZUSP.
- lenkoi** Papavero, 1975: 283, figs. 137-139. Type-locality: Brazil, Mato Grosso, Utariiti. Distr. – Brazil (Mato Grosso). Type, MZUSP.
- mendax** Curran, 1934c: 333. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Distr. – Colombia, Venezuela, Ecuador, Peru, Guyana, French Guiana, Brazil (Amazonas). Refs. – Carrera, 1949: 21, 1952c: 67; Hull, 1962 (1): 422; Carrera & Machado-Allison, 1963: 239; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 284, figs. 161-163;

Fisher, 1984: 28. Type, AMNH.

mundata (Wiedemann), 1838: 569 (*Dasypogon*). Type-locality: "Brazil". Distr. – Brazil (Rio de Janeiro, São Paulo, Santa Catarina). Refs. – Loew, 1851: 11 (*Stenobasis*); Walker, 1854: 444 (*Dasypogon*); Schiner, 1866: 681 (*Dasypogon*); Williston, 1891: 67 (cat.; *Dasypogon*); Kertész, 1909: 130 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Carrera, 1949: 28, 1952c: 72, 1960a: 157; Hull, 1962 (1): 228 (*Dasypogon*), 422 [*Senobasis*]; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 285, figs. 164–168. Type, SENCK [cf. Geller-Grimm, 1999: 211].

gyrophora Schiner of Carrera, 1949: 29, misident.

notata (Bigot), 1878: 416 (?*Diogmites*). Type-locality: "Amazonia". Distr. – Ecuador, Peru (Loreto), Brazil (Amazonas, Pará, Amapá). Refs. – Williston, 1891: 75 (cat.; *Deromyia*); Kertész, 1909: 117 (cat.; *Deromyia*); Carrera & d'Andretta, 1953: 65 (?*Senobasis*); Hull, 1962 (1): 232 (*Diogmites*); Martin & Papavero, 1970: 26 (cat.; *Diogmites*); Papavero, 1975: 287, figs. 34, 170–172; Fisher, 1984: 28. Type, lost.

weyrauchi Carrera, 1952c: 72, fig. 9. Type-locality: Peru, Loreto, Pucallpa, 200 m. Refs. – Carrera & d'Andretta, 1953: 64; Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Lamas M., 1972: 312 (cat.); Papavero, 1975: 287 (syn.).

ornata (Wiedemann), 1819: 49 (*Laphria*). Type-locality: "Brazil" [Freireiss (somewhere in southeastern Brazil, probably Rio de Janeiro or Espírito Santo; cf. Papavero, 1971b: 57–58)]. Distr. – Brazil (Espírito Santo, Rio de Janeiro). Refs. – Wiedemann, 1821: 222 (*Dasypogon*), 1828: 386, 585 (*Dasypogon*); Walker, 1854: 444 (*Dasypogon*); Schiner, 1866: 677 (*Lochites*), 1867: 369 (*Lochites*); Williston, 1891: 75 (cat.); Kertész, 1909: 125 (cat.; *Lochites*); Bromley, 1946: 106 (cat.; n. comb.); Carrera, 1949: 30; Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 287, figs. 33, 167–169. Type, SENCK [cf. Geller-Grimm, 1999: 213].

rhombungulata Carrera, 1949: 20, pl. 16, fig. 150. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Refs. – Hull, 1962 (1): 422; Martin & Papavero, 1970: 58 (cat.); Papavero, 1975: 287 (syn.). Type, IOC.

Subfamily Laphriinae Macquart, 1838

Laphritae Macquart, 1838: 56 (1839: 172). Type-genus, *Laphria* Meigen, 1803.

Laphritidae Bigot, 1852: 485.

Laphridae Bigot, 1857b: 536.

Laphriinae Schiner, 1860: 209.

Laphriida Gistel, 1848: 152.

Laphrites Walker, 1855: 507.

Laphriina Rondani, 1856: 32.

Laphrina Rondani, 1856: 160, 212.

Laphira Valdés-Raguès, 1908: 309.

Tribe Andrenosomatini Hull, 1962

Erecmoneminae Hermann, 1912: 21, part.

Andrenosomini Hull, 1962 (1): 348. Type-genus, *Andrenosoma* Rondani, 1856.

Andrenosomatini Sabrosky, 1999: 45, corrected spelling.

Genus *Andrenosoma* Rondani, 1856

Andrenosoma Rondani, 1856: 160. Type-species, *Asilus ater* Linnaeus, 1758 (orig. des.).

albolineatum (Macquart), 1850: 378 (1850: 74) (*Laphria*). Type-locality: Bolivia, Mojos [= Depto. Beni]. Distr. – Bolivia. Refs. – Williston, 1891: 79 (cat.; *Laphria*); Kertész, 1909: 176 (cat.; *Laphria*); Martin & Papavero, 1970: 42 (cat.). Type, MNHNP.

appendiculatum (Macquart), 1846: 203 (1846: 75), pl. 7, fig. 18 (*Laphria*). Type-locality: French Guiana, Cayenne. Distr. – French Guiana. Refs. – Walker, 1855: 536 (*Laphria*); Williston, 1891: 81 (cat.; *Andrenosoma*); Kertész, 1909: 197 (cat.; *Nusa*); Hull, 1962 (1): 350; Martin & Papavero, 1970: 42 (cat.).

avicenniae Farr, 1965: 12, figs. 5–6. Type-locality: Jamaica, Clarendon, Portland Ridge. Distr. – Jamaica. Ref. – Martin & Papavero, 1970: 42 (cat.). Type, BMNH.

batesi Bromley, 1931a: 132. Type-locality: Brazil, "Amazon river". Distr. – Brazil (Amazonia). Refs. – Bromley, 1946: 110 (cat.); Martin & Papavero, 1970: 42 (cat.). Type, BMNH.

- camposi** Curran, 1931b: 20. Type-locality: Ecuador, Guayas, Guayaquil. Distr. – Ecuador. Refs. – Hull, 1962 (1): 350; Martin & Papavero, 1970: 42 (cat.). Type, AMNH.
- chalybeum** Williston, 1885: 56. Type-locality: “San Domingo”. Distr. – Cuba, Dominican Republic (Azua, Barahona, Hato Mayor, La Altagracia, Pedernales), Puerto Rico. Refs. – Aldrich, 1905: 1269 (cat.; *Nusa*); Kertész, 1909: 198 (cat.; *Nusa*); Bromley, 1929a: 284, pl. 1, fig. 17; Hull, 1962 (1): 350; Martin & Papavero, 1970: 42 (cat.); Alayo Dalmau & García Ávila, 1983: 46 (cat.); Fisher, 1986: 90; Scarbrough & Perez-Gelabert, 2006: 17.
- cinctum** (Bellardi), 1861: 118 (18), pl. 1, fig. 19 (*Laphria*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Osten Sacken, 1887: 188; Aldrich, 1905: 269 (cat.; *Nusa*); Kertész, 1909: 198 (cat.; *Nusa*); Hermann, 1912: 254 (*Nusa*, as syn. of *clausicella* Macquart, 1850); Hull, 1962 (1): 350; Martin & Papavero, 1970: 42 (cat.); Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, TORO.
- cinereum** (Bellardi), 1861: 116 (16), pl. 1, fig. 16 (*Laphria*). Type-locality: “Mexico”. Distr. – Mexico. Refs. – Osten Sacken, 1887: 188; Aldrich, 1905: 269 (cat.; *Nusa*); Kertész, 1909: 198 (cat.; *Nusa*); Hull, 1962 (1): 350; Martin & Papavero, 1970: 42 (cat.); Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Syntypes, TORO.
- corallium** Martin, 1966c: 327. Type-locality: Mexico, Oaxaca, Valerio Trujano. Distr. – Mexico (Oaxaca). Ref. – Martin & Papavero, 1970: 42 (cat.). Type, CAS.
- cruentum** (McAtee), 1919: 244 (*Nusa*). Type-locality: U. S. A., Florida. Distr. – U. S. A. (Florida), Mexico. Refs. – Bromley, 1950: 233; Hull, 1962 (1): 350; Martin & Wilcox, 1965: 386 (cat.).
- crassum** Bromley, 1929a: 285, fig. 9, pl. 1, figs. 11, 13. Type-locality: Cuba, Camaguey. Refs. – Hull, 1962 (1): 350; Martin & Papavero, 1970: 42 (cat.); Alayo Dalmau & García Ávila, 1983: 46 (cat.). Type, USNM.
- currani** Bromley, 1931a: 133. Type-locality: Brazil, Rio Grande do Sul, Santa Cruz. Distr. – Brazil (Rio Grande do Sul). Refs. – Bromley, 1946: 110 (cat.); Martin & Papavero, 1970: 42 (cat.). Type, HAMBURG.
- erythropyga** (Wiedemann), 1828: 509 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Peru (Lima), Brazil. Refs. – Walker, 1849: 375 (*Laphria*), 1855: 535 (*Laphria*); Schiner, 1866: 691; Wulp, 1882b: 104; Williston, 1891: 83 (cat.; *Laphria*); Kertész, 1909: 199 (cat.; *Nusa*); Hermann, 1912: 265, fig. 85 (*Nusa*); Curran, 1930b: 20 (in key); Bromley, 1946: 110 (cat.); Carrera & d’Andretta, 1953: 64; Carrera, 1960a: 156; Hull, 1962 (1): 350; Martin & Papavero, 1970: 42 (cat.); Lamas M., 1972: 308 (cat.) TP SENCK [cf. Geller-Grimm, 1999: 208].
- heros** Bromley, 1931a: 131. Type-locality: Surinam, Demerara. Distr. – Surinam, Brazil (Pará). Refs. – Bromley, 1946: 110 (cat.); Carrera, 1960a: 155; Martin & Papavero, 1970: 43 (cat.). Type, BMNH.
- igneum** Bromley, 1929b: 46. Type-locality: Mexico, Veracruz, Coatepec. Distr. – Mexico (Veracruz). Refs. – Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.). Type, USNM.
- lewisi** Farr, 1965: 13, figs. 7-8. Type-locality: Jamaica, St. Thomas, Morant Point. Distr. – Jamaica. Ref. – Martin & Papavero, 1970: 43 (cat.). Type, USNM.
- mesoxanthum** (Wiedemann), 1828: 513 (*Laphria*). Type-locality: “Brazil”. Refs. – Macquart, 1838: 67 (*Laphria*); Walker, 1855: 538 (*Laphria*); Schiner, 1866: 691; Williston, 1891: 81 (cat.); Kertész, 1909: 200 (cat.; *Nusa*); Bromley, 1946: 110 (cat.); Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.).
- nigrum** Bromley, 1931a: 132. Type-locality: Ecuador, Guayas, Guayaquil. Distr. – Ecuador, Peru (La Libertad, Lima). Refs. – Martin & Papavero, 1970: 43 (cat.); Lamas M., 1972: 308 (cat.). Type, BMNH.
- purpurascens** (Walker), 1856: 457 (*Laphria*). Type-locality: Brazil, Amazonas. Distr. – Brazil (Amazonas). Refs. – Williston, 1891: 81 (cat.); Kertész, 1909: 200 (cat.; *Nusa*); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.).
- pygophorum** Schiner, 1868: 175. Type-locality: “South America”. Refs. – Williston, 1891: 81 (cat.); Kertész, 1909: 200 (cat.; *Nusa*); Hermann, 1912: 258, pl. 5, fig. 63 (*Nusa*); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 350; Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.). Type, NMW.
- pyrrhacra** (Wiedemann), 1828: 517 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1855: 526 (*Laphria*); Schiner, 1866: 692, 1868: 175; Kertész, 1909: 199 (cat.; *Nusa*, as syn. of *fulvicauda* Say, 1823); Hermann, 1912: 265 (*Nusa*); Bromley, 1946: 110 (cat.); Carrera & Vulcano, 1961: 70 (preys). Type, SENCK [cf. Geller-Grimm, 1999: 213].
- quadrimaculatum** Bromley, 1929a: 285, pl. 1, fig. 19. Type-locality: “Cuba”. Distr. – Cuba. Refs. – Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.); Alayo Dalmau & García Ávila, 1983: 456 (cat.). Type, MCZ.
- rubidum** (Williston), 1901: 318 (*Nusa*). Type-locality: Mexico, Veracruz, Atoyac. Distr. – U. S. A. (Texas), Mexico (Veracruz). Refs. – Aldrich, 1905: 270 (cat.; *Nusa*); Kertész, 1909: 200 (cat.; *Nusa*); Hull, 1962 (1): 350; Martin & Wilcox, 1965: 386 (cat.); Martin & Papavero, 1970: 43 (cat.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol.). Type, BMNH.
- ruficauda** (Williston), 1885: 55 (*Laphria*). Type-locality: “Santo Domingo”. Distr. – Dominican Republic (Azua, El Seibo, La Altagracia, Pedernales). Refs. – Aldrich, 1905: 273 (cat.; *Laphria*); Kertész, 1909: 193 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; n. comb.); Scarbrough & Perez-Gelabert, 2006: 18 (as *ruficaudum*, error).
- rufiventris** (Blanchard), 1852: 364, pl. 1, fig. 5 (*Laphria*). Type-locality: Chile, Concepción. Distr. – Chile (Aconcagua, Biobío, Coquimbo, Linares, Maule, O’Higgins, Santiago, Valparaíso). Refs. – Philippi, 1865: 684 (*Laphria*); Schiner, 1868: 176 (n. comb.); Williston, 1891: 81 (cat.); Kertész, 1909: 200 (cat.; *Nusa*); Hermann, 1912: 261 (*Nusa*); Curran, 1930b: 20 (in key); Bromley, 1932: 168; Stuardo Ortiz, 1946: 84 (cat.); Hull, 1962 (1): 350; Artigas, 1970a: 239, figs. 197-202, 466, 1971: 30, fig. 67-68; Martin & Papavero, 1970: 43 (cat.). Type, MNHNP.
- calogastra** Philippi, 1865: 684 (*Laphria*). Type-locality: Chile, Valdivia. Refs. – Williston, 1891: 79 (cat.; *Laphria*); Kertész, 1909: 179 (cat.; *Laphria*); Hermann, 1912: 262 (*Nusa*); Curran, 1930b: 20 (in key); Bromley, 1932: 268;

Stuardo Ortiz, 1946: 83 (cat.); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 43 (cat.; syn.). Type, lost.
sarcophaga (Hermann), 1912: 263 (*Nusa*). Type-locality: Peru, Huánuco, mouth of Pachitea river, 150 m, and Cuzco, Vilcanota (as Villanota, error). Distr. – Peru (Huánuco). Refs. – Curran, 1930b: 20 (in key); Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.); Lamas M., 1972: 308 (cat.); Fisher, 1984: 30. Type, ?
sexpunctatum (Williston), 1901: 319, pl. 6, fig. 4 (*Nusa*). Type-locality: Mexico, Guerrero, Amula. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 270 (cat.; *Nusa*); Kertész, 1909: 200 (cat.; *Nusa*); Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.
subheros Bromley, 1931a: 131. Type-locality: Brazil, Rio Grande do Sul, Santa Cruz. Distr. – Brazil (Amazonas to Espírito Santo, south to Santa Catarina and Rio Grande do Sul). Refs. – Bromley, 1946: 111 (cat.); Carrera, 1960a: 155; Martin & Papavero, 1970: 43 (cat.).
varipes (Banks), 1920: 67 (*Laphria*). Type-locality: “Cuba”. Distr. – Cuba. Refs. – Bromley, 1929a: 284; Hull, 1962 (1): 350; Martin & Papavero, 1970: 43 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 46 (cat.).
xanthocnemum (Wiedemann), 1828: 509 (*Laphria*). Type-locality: “Brazil”. Distr. – ?Mexico, Venezuela, Guyana, Peru (Junín), Brazil, Paraguay. Refs. – Macquart, 1838: 67 (*Laphria*); Walker, 1855: 691 (*Laphria*); Schiner, 1866: 691 (n. comb.); Wulp, 1882b: 104; Williston, 1891: 81 (cat.); Aldrich, 1905: 170 (cat.; *Nusa*); Hermann, 1905: 31, fig. 19; Kertész, 1909: 201 (cat.; *Nusa*); Hermann, 1912: 258; Curran, 1930b: 20 (in key); Engel, 1930: 468 (*Nusa*); Bromley, 1934: 342, 1946a: 111 (cat.); Carrera, 1947a: 45; Carrera & d’Andretta, 1953: 64; Carrera, 1960a: 156; Hull, 1962 (1): 350; Carrera & Machado-Allison, 1963: 241; Martin & Papavero, 1970: 43 (cat.); Lamas M., 1972: 308 (cat.); Notario, Michela, Fiorentino & Castresana, 2000: 17 (“emergieron de trozas en las que sólo se encontraron *B[rasilianus lacordairei* [(Gahan, 1892)], *B[rasilianus murinus* [(Gahan, 1892)] [Coleoptera, Cerambycidae] y *C[hrysobothris holoclalcea?* [Burmeister, 1872] [Coleoptera, Buprestidae]”). Type, SENCK.

Genus **Cerotainiops** Curran, 1930

Cerotainiops Curran, 1930a: 11. Type-species, *Cerotainiops rufiventris* Curran, 1930 (orig. des.).

omus Pritchard, 1942: 22. Type-locality: U. S. A., California. Distr. – U. S. A. (California, Arizona), Mexico (Baja California Norte, Baja Califotnia Sur, Sonora). Refs. – Hull, 1962 (1): 355; Martin & Wilcox, 1965: 388 (cat.).
wilcoxi Pritchard, 1942: 23. Type-locality: U. S. A., Arizona, Tempe. Distr. – U. S. A. (Arizona), Mexico (Baja California). Refs. – Hull, 1962 (1): 355; Martin & Wilcox, 1965: 388 (cat.); Martin & Papavero, 1970: 45 (cat.).
abdominalis Brown of authors, misident.
pritchardi Hull, 1960: 70. Type-locality: Mexico, Baja California. TP, CAS.

Genus **Dasyllis** Loew, 1851

Laphria, subg. **Dasyllis** Loew, 1851: 20. Type-species, *Laphria haemorrhoa* Wiedemann, 1830 (orig. des.).

Ref. – Artigas, Papavero & Pimentel, 1988: 245, figs. 1-2 (key to spp.).

albicollis Bigot, 1878: 229. Type-locality: “South America”. Distr. – Paraguay. Refs. – Williston, 1891: 79 (cat.); Kertész, 1909: 172 (cat.); Hermann, 1912: 238; Bromley, 1951: 21 (list); Carrera, 1958b: 219; Hull, 1962 (1): 359; Martin & Papavero, 1970: 45 (cat.); Artigas, Papavero & Pimentel, 1988: 245. Type, OXF.
croceiventris (Wiedemann), 1821: 234 (*Laphria*). Type-locality: “Brazil”. Distr. – Brazil (Amazonas, Pará, Amapá, Rio de Janeiro, São Paulo). Refs. – Wiedemann, 1828: 503 (*Laphria*); Loew, 1851: 20; Walker, 1855: 537 (*Laphria*); Schiner, 1866: 691; Williston, 1891: 79 (cat.); Kertész, 1909: 172 (cat.); Hermann, 1912: 237; Bromley, 1946: 110 (cat.), 1951: 21 (list); Carrera, 1958b: 217, figs. 2-3, 1960a: 157; Carrera & Vulcano, 1961: 70 (preys); Hull, 1962 (1): 359; Martin & Papavero, 1970: 45 (cat.); Artigas, Papavero & Pimentel, 1988: 245. Syntypes, BZM [no. 418; cf. Schumann, 1973: 100].
erythrura Hermann, 1912: 238. Type-locality: “Paraguay”. Distr. – Paraguay. Refs. – Bromley, 1951: 21 (list); Carrera, 1958b: 218; Hull, 1962 (1): 359; Martin & Papavero, 1970: 45 (cat.); Artigas, Papavero & Pimentel, 1988: 245. Syntypes, ZSBS.
fascipennis (Macquart), 1834: 284 (*Laphria*). Type-locality: “Cayenne”. Distr. – French Guiana, Brazil (Pará, Mato Grosso). Refs. – Walker, 1855: 536 (*Laphria*); Williston, 1891: 79 (cat.); Aldrich, 1905: 271 (cat.); Kertész, 1909: 271 (cat.); Hermann, 1912: 237; Bromley, 1934b: 337, 1951: 21, fig. 1 (list); Carrera, 1958b: 216, fig. 4, 1960a: 157; Hull, 1962 (1): 359; Martin & Papavero, 1970: 45 (cat.); Lamas M., 1972: 309 (cat.). Type, lost.
praepotens Macquart, 1846: 202 (1846: 74), pl. 7, fig. 17 (*Laphria*). Type-locality: French Guiana, Cayenne. Refs. – Walker, 1855: 535 (*Laphria*); Schiner, 1868: 172; Hermann, 1912: 237. Type, lost.
haemorrhoa (Wiedemann), 1830: 645 (*Laphria*). Type-locality: Brazil, Bahia [Gomez; cf. Papavero, 1971e: 48]. Distr. –

Panama, French Guiana, Peru, Brazil (Amazonas, Bahia, Espírito Santo, Rio de Janeiro), Bolivia. Refs. – Walker, 1855: 536 (*Laphria*); Schiner, 1866: 694; Williston, 1891: 79 (cat.); Kertész, 1909: 79 (cat.); Hermann, 1912: 237; Bromley, 1934b: 337, 1946a: 110 (cat.), 1951: 21 (list); Carrera, 1958b: 214, fig. 1; Hull, 1962 (1): 359, figs. 768, 1261, 1270, 2080, 2082, 2129, 2175; Carrera & Machado-Allison, 1963: 241; Martin & Papavero, 1970: 45 (cat.); Lamas M., 1972: 309 (cat.); Fisher, 1984: 30 (“A spectacular mimic of orchid bees of the *Eulaema meriana* grouo”). Type, BZM [no. 417; cf. Schumann, 1973: 116].

rufa (Bromley), 1931a: 132 (*Andrenosoma*). Type-locality: “Santo Domingo”. Distr. – Dominican Republic (Azua, La Altagracia, Pedernales, Santo Domingo), Haiti. Refs. – Martin & Papavero, 1970: 43 (cat.; *Andrenosoma*); Scarbrough & Perez-Gelabert, 2006: 20. Type, BMNH.

Genus **Pilica** Curran, 1931

Pilica Curran, 1931b: 20. Type-species, *Laphria formidolosa* Walker, 1860 (orig. des.).

Ref. – Artigas, Papavero & Pimentel, 1988: 247 (key to South American species).

acunai (Bromley), 1929a: 286, pl. 1, fig. 7 (*Andrenosoma*). Type-locality: Cuba, Camaguey. Distr. – Cuba. Ref. – Hull, 1962 (1): 350 (*Andrenosoma*); Martin & Papavero, 1970: 44 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 456 (cat.). Type, (remains) USNM.

cyrtophora (Hermann), 1912: 255, fig. 84 (*Nusa*). Type-locality: Brazil, Minas Gerais. Distr. – Brazil (Minas Gerais). Refs. – Bromley, 1946: 110 (cat.; *Andrenosoma*); Hull, 1962 (1): 353; Martin & Papavero, 1970: 44 (cat.).

erax (Bromley), 1934b: 345 (*Andrenosoma*). Type-locality: Guyana, Cuyuni-Mazaruni, Bartica. Distr. – Guyana. Refs. – Hull, 1962 (1): 353; Martin & Papavero, 1970: 44 (cat.). Type, AMNH (only allotype, a teneral specimen).

erythrogaster (Wiedemann), 1828: 523 (*Laphria*). Type-locality: “Brazil” [Freyreiss, cf. Papavero, 1971e: 57-58]. Distr. – Peru (Cuzco), Brazil (Rondônia, Amazonas, Pará, Amapá, Maranhão, Pernambuco, Espírito Santo, Rio de Janeiro). Refs. – Walker, 1855: 538 (*Laphria*); Schiner, 1866: 692 (*Andrenosoma*), 1867: 382 (*Andrenosoma*); Williston, 1891: 81 (cat.; *Andrenosoma*); Brauer & Bischof, 1903: 244 (*Andrenosoma*); Kertész, 1909: 199 (cat.; *Nusa*); Bromley, 1946: 110 (cat.; *Andrenosoma*); Carrera, 1960a: 156 (*Andrenosoma (Pilica)*); Hull, 1962 (1): 353; Carrera & Machado-Allison, 1963: 240 (*Andrenosoma (Pilica)*); Martin & Papavero, 1970: 44 (cat.); Lamas M., 1972: 309 (cat.); Fisher, 1984: 30; Artigas, Papavero & Pimentel, 1988: 248. Syntypes, BZM [cf. Schumann, 1973: 107], SENCK [cf. Geller-Grimm, 1999: 208].

contusa Wiedemann, 1828: 587 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1855: 537 (*Laphria*); Schiner, 1866: 693 (*Laphria*); Williston, 1891: 79 (*Laphria*); Kertész, 1909: 180 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*); Artigas, Papavero & Pimentel, 1988: 248 (syn.). Type, SENCK [cf. Geller-Grimm, 1999: 207].

clausicella Macquart, 1850: 378 (1850: 74), pl. 7, fig. 6 (*Laphria*). Type-locality: “Guyana”. Refs. – Williston, 1891: 81 (cat.; *Andrenosoma*); Hermann, 1912: 254 (*Nusa*); Bromley, 1934b: 342 (*Andrenosoma*); Hull, 1962 (1): 353. Type, OXF.

laticornis Walker, 1855: 535 (*Laphria*). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Williston, 1891: 79 (*Laphria*), 81 (*Andrenosoma*); Kertész, 1909: 200 (cat.; *Nusa*); Hermann, 1912: 254 (*Nusa*; as syn. of *clausicella* Macquart, 1850); Bromley, 1946: 110 (cat.; *Andrenosoma*); Hull, 1962 (1): 350 (*Andrenosoma*). Type, BMNH.

formidolosa (Walker), 1860: 280 (*Laphria*). Type-locality: “Mexico”. Distr. – Mexico (Veracruz), Guatemala, Nicaragua, Panama. Refs. – Bellardi, 1861: 16, pl. 1, fig. 18 (*Laphria*); Osten Sacken, 1887: 188 (*Andrenosoma*); Aldrich, 1905: 270 (cat.; *Nusa*); Kertész, 1909: 199 (cat.; *Nusa*); Hermann, 1912: 250, fig. 82 (*Nusa*); Hull, 1962 (1): 353, figs. 239, 638, 1240, 1249, 2076, 2085; Martin & Papavero, 1970: 44 (cat.); Fisher, 1983: 755, 1991: 769; Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.). Type, BMNH.

funebris Artigas, Papavero & Pimentel, 1988: 249. Type-locality: Brazil, São Paulo, Onda Verde (Fazenda São João). Distr. – Brazil (Amazonas, Pará, Mato Grosso, Mato Grosso do Sul, Pernambuco, Minas Gerais, São Paulo, Santa Catarina). Type, MZUSP.

lupus (Bromley), 1931a: 130 (*Andrenosoma*). Type-locality: Brazil, Amazonas, Tefé (as Ega). Distr. – Brazil (Amazonas, Pará). Refs. – Bromley, 1946: 110 (cat.); Carrera, 1960a: 156 (*Andrenosoma*); Martin & Papavero, 1970: 43 (cat.; *Andrenosoma*); Artigas, Papavero & Pimentel, 1988: 249. Type, BMNH.

olbus (Walker), 1849: 375 (*Laphria*). Type-locality: “Honduras”. Distr. – Honduras, Guatemala, Costa Rica, Venezuela, Peru. Refs. – Hermann, 1912: 258 (*Nusa*); Enderlein, 1914a: 252 (*Laphria*); Bromley, 1946: 110 (cat.; *Andrenosoma*); Hull, 1962 (1): 353; Carrera & Machado-Allison, 1963: 240 (*Andrenosoma (Pilica)*); Martin & Papavero, 1970: 44 (cat.); Lamas M., 1972: 309 (cat.). Type, BMNH.

minos Bromley, 1931a: 130 (*Andrenosoma*). Type-locality: Brazil, “Amazon river”. Distr. – Venezuela, Guyana, Ecuador, Brazil (Amazonas). Ref. – Bromley, 1946: 110 (cat.; *Andrenosoma*). Type, BMNH.

cyaniventris Bromley, 1934b: 344, fig. 23 (*Andrenosoma*). Type-locality: Guyana, Tukeit. Refs. – Hull, 1962 (1): 353; Artigas, Papavero & Pimentel, 1988: 249 (syn.). Type, AMNH.

- phoenicogaster** (Hermann), 1912: 248 (*Nusa*, as *phönicogaster*). Type-locality: Bolivia, Mapiri, Sarampiuni, 700 m; San Ernesto; San Carlos, 800 m; and “Bolivia”. Distr. – Peru, Bolivia, Brazil (Amazonas, Pará). Refs. – Hull, 1962 (1): 353; Martin & Papavero, 1970: 44 (cat.); Lamas M., 1972: 309 (cat.); Fisher, 1984: 30 (*Andrenosoma*); Artigas, Papavero & Pimentel, 1988: 249. Syntypes, MUN, BMNH, AMNH, NMW, TP SMTD [cf. Kallweit, 1992: 197].
- pyrrhopya** (Wiedemann), 1828: 515 (*Laphria*). Type-locality: “Brazil: [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Venezuela, Guyana, Peru (Huánuco, Lima), Brazil (Amazonas, Pará, Amapá, Mato Grosso, Rio de Janeiro, São Paulo, Paraná), Paraguay. Refs. – Walker, 1855: 535 (*Laphria*); Schiner, 1866: 692 (*Andrenosoma*); Williston, 1891: 81 (cat.; *Andrenosoma*); Kertész, 1909: 200 (cat.; *Nusa*); Hermann, 1912: 254, fig. 83 (*Nusa*); Bromley, 1946: 111 (cat.; *Andrenosoma*); Carrera, 1947d: 270 (prey); Carrera, 1960a: 156 (*Andrenosoma (Piica)*); Hull, 1962 (1): 353; Martin & Papavero, 1970: 44 (cat.); Lamas M., 1972: 309 (cat.); Artigas, Papavero & Pimentel, 1988: 250. Type, SENCK [cf. Geller-Grimm, 1999: 213].
- punctata* Bromley, 1934b: 394 (*Andrenosoma*). Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Refs. – Carrera, 1960a: 156 (*Andrenosoma (Pilica)*); Martin & Papavero, 1970: 44 (cat.); Lamas M., 1972: 309 (cat.); Artigas, Papavero & Pimentel, 1988: 250 (syn.). Type, AMNH.
- rubidapex** (Hermann), 1912: 247 (*Nusa*). Type-locality: “Venezuela”. Distr. – Venezuela, Guyana, Brazil (Amapá). Refs. – Hull, 1962 (1): 353; Martin & Papavero, 1970: 44 (cat.); Artigas, Papavero & Pimentel, 1988: 251. Type, MUN.
- elegans* Bromley, 1934b: 343 (*Andrenosoma*). Type-locality: Guyana, Kamakusa. Ref. – Hull, 1962 (1): 353.
- rufipennis** (Wiedemann), 1828: 522 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Venezuela, Guyana, Surinam, Brazil (Amazonas, Minas Gerais, Rio de Janeiro, São Paulo). Refs. – Macquart, 1846: 73, 1848: 24 (*Laphria*); Walker, 1855: 536 (*Laphria*); Schiner, 1866: 692 (*Laphria*), 1867: 380 (*Laphria*); Williston, 1891: 80 (cat.; *Laphria*); Hermann, 1912: 248 (*Nusa*); Bromley, 1934b: 342 (*Andrenosoma*); Carrera, 1960a: 156 (*Andrenosoma (Pilica)*); Hull, 1962 (1): 353; Carrera & Machado-Allison, 1963: 240 (*Andrenosoma (Pilica)*); Martin & Papavero, 1970: 44 (cat.); Artigas, Papavero & Pimentel, 1988: 251. Lectotype SENCK [des. by Artigas, Papavero & Pimentel, 1988: 247]; Paralectotype BZM [cf. Schumann, 1973: 140].
- flamipennis* Bromley, 1931a: 133 (*Andrenosoma*). Type-locality: Guyana, Cuyuni-Mazaruni, Bartica. Ref. – Bromley, 1946: 110 (cat.; *Andrenosoma*). Type, ANSP.
- †sp. Poinar Jr., 1992: 178. Mexico (Oligocene/Miocene).
- sp. Scarbrough & Perez-Gelabert, 2006: 21 (Dominican Republic: Azua, La Altagracia).
- zanutoi** Artigas, Papavero & Pimentel, 1988: 251. Type-locality: Brazil, Mato Grosso, Barra dos Bugres (Reserva Ecológica de Serra das Araras). Distr. – Brazil (Mato Grosso, São Paulo). Type, MPEG.

Genus **Pogonosoma** Rondani, 1865

Pogonosoma Rondani, 1865: 160. *Type-species*, *Asilus maroccanus* Fabricius, 1794 (orig. des.).

arachnoides Bigot, 1878: 227. Type-locality: “Mexico”. Distr. – Mexico. Refs. – Williston, 1883: 33; Osten Sacken, 1887: 186 (*Laphria*); Aldrich, 1905: 269 (cat.); Kertész, 1909: 175 (cat.); Hull, 1962 (1): 351; Martin & Papavero, 1970: 44 (cat.); Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipterol). Type, OXF.

Tribe **Atomosiini** Lynch Arribálzaga, 1882

Atomosiaria Lynch Arribálzaga, 1882: 144. Type-genus, *Atomosia* Macquart, 1838. Ref. – Dikow, 2003: 465.
Eremoneminae Hermann, 1912: 21, part.
Atomosinae Hermann, 1912: 21.
Atomosiinae G. H. Hardy, 1928: 472.
Atomasinae Engel, 1929: 161.

Genus **Aphestia** Schiner, 1866

Aphestia Schiner, 1866: 663 (key), 673. *Type-species*, *Aphestia brasiliensis* Schiner, 1866 (orig. des.) = *Aphestia annulipes* (Macquart, 1838).

annulipes (Macquart), 1838: 74 (1839: 190) (*Atomosia*). Type-locality: “Brazil”. Distr. – Brazil (Rondônia, Mato Grosso, Goiás, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Santa Catarina). Refs. – Walker, 1855: 569 (*Atomosia*); Williston, 1891: 78 (cat.; *Atomosia*); Kertész, 1909: 163 (cat.; *Atomosia*); Hermann, 1912: 122; Bromley, 1946: 108

- (cat.; *Atomosia*); Carrera & d'Andretta, 1952: 256 (preys); Hull, 1962 (1): 383, figs. 292, 648, 1195, 1204; Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 57, figs. 18-24.
- affinis* Macquart, 1850: 379 (1850: 75) (*Atomosia*). Type-locality: "Brazil". Refs. – Williston, 1891: 77 (cat.); Kertész, 1909: 163 (cat.); Bromley, 1946: 108 (cat.; *Atomosia*); Hull, 1962 (1): 373 (*Atomosia*); Martin & Papavero, 1970: 46 (cat.; *Atomosia*); Artigas, Papavero, & Serra, 1991: 57 (syn.).
- brasiliensis* Schiner, 1866: 378. Type-locality: "Brazil". Refs. – Williston, 1891: 78 (cat.); Kertész, 1909: 167 (cat.); Hermann, 1912: 122 (syn.); Bromley, 1946: 110 (cat.). Type, NMW.
- calceata* Schiner, 1867: 379. Type-locality: "Brazil". Refs. – Williston, 1891: 78 (cat.); Kertész, 1909: 167 (cat.); Hermann, 1912: 123; Bromley, 1946: 110 (cat.); Hull, 1962 (1): 383; Martin & Papavero, 1970: 50 (cat.). Type, NMW.
- calcarata* Williston, 1901: 317, error.
- nigra* Bigot, 1878: 235. Type-locality: "Mexico". Distr. – Mexico, Guyana. Refs. – Williston, 1883: 33; Osten Sacken, 1887: 185 (*Atomosia*); Aldrich, 1905: 268 (cat.); Kertész, 1909: 164 (cat.; *Atomosia*); Hermann, 1912: 123; Hull, 1962 (1): 383; Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 57; Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipterol).
- mexicana* Williston, 1901: 317, pl. 6, fig. 2. Type-locality: Mexico, Guerrero, Amula and Veracruz, Atoyac. Refs. – Aldrich, 1905: 268 (cat.); Kertész, 1909: 167 (cat.); Hull, 1962 (1): 383; Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 58 (syn.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.
- andrenoides* Bromley, 1934b: 339 (*Atomosia*). Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Refs. – Curran, 1935: 5 (in key); Hull, 1962 (1): 373 (*Atomosia*); Martin & Papavero, 1970: 46 (cat.; *Atomosia*); Artigas, Papavero & Serra, 1991: 58 (syn.). Type, AMNH.

Genus *Aphractia* Artigas, Papavero & Serra, 1991

Aphractia Artigas, Papavero & Serra, 1991: 58. Type-species, *Atractia longicornis* Hermann, 1912 (orig. des.).

- longicornis*** (Hermann), 1912: 190, fig. 74 (*Atractia*). Type-locality: Brazil, Bahia. Distr. – Brazil (Bahia). Refs. – Bromley, 1946: 108 (cat.; *Atractia*); Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 58 (n. comb.). Type, NMW.
- rubida*** (Hermann), 1912: 179 (*Atractia*). Type-locality: Brazil, Rio Grande do Sul. Distr. – Brazil (Rio Grande do Sul). Refs. – Bromley, 1946: 110 (cat.; *Atractia*); Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 58 (n. comb.). Type, NMW.
- vivax*** (Hermann), 1912: 177, fig. 67 (*Atractia*). Type-locality: Peru, Cuzco, Umuhaniali, Ururubamba river, 500 m and "Peru". Distr. – Peru (Cuzco). Refs. – Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Lamas M., 1972: 310 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 58 (n. comb.). Syntypes, ZSBS, SMTD [cf. Kallweit, 1992: 198].

Genus *Atomosia* Macquart, 1838

Atomosia Macquart, 1838: 73 (1839: 189). Type-species, *Atomosia incisuralis* Macquart, 1838 (sub. des., Coquillett, 1910: 512) = *puella* (Wiedemann, 1828).

Paratractia Hull, 1962 (1): 384. Type-species, *Laphria dasypus* Wiedemann, 1828 (orig. des.).

- anacaona*** Scarbrough & Perez-Gelabert, 2006: 22 (in key and text), figs. 3-7. Type-locality: Dominican Republic, Santiago, Road-243 Babosico, on way to Jánico. Distr. – Dominican Republic (Santiago). Type, USNM.
- anonyma*** Williston, 1901: 316. Type-locality: Mexico, Guerrero, Chilpancingo. Refs. – Aldrich, 1905: 268 (cat.); Curran, 1930b: 16 (in key), 1935: 6 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Syntypes, AMNH, BMNH.
- argyrophora*** Schiner, 1868: 169. Type-locality: "Brazil". Refs. – Williston, 1891: 78 (cat.); Kertész, 1909: 163 (cat.); Hermann, 1912: 163, fig. 61; Curran, 1930b: 15 (in key), 1935: 5 (in key); Bromley, 1946: 109 (cat.); Carrera & d'Andretta, 1952: 254 (preys), 1961: 70 (preys); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, NMW.
- armata*** Hermann, 1912: 152, fig. 58. Type-locality: "Brazil". Distr. – Brazil, Paraguay. Refs. – Curran, 1930b: 15 (in key), 1935: 5 (in key); Bromley, 1946: 109 (cat.); Carrera, 1947a: 44; Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58, figs. 87-88 (cat.). Syntypes, MUN, MIL.
- barbiellinii*** Curran, 1935: 5 (in key), 7. Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 109 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, AMNH.

- beckeri** Jaennicke, 1867: 359. Type-locality: "Mexico". Refs. – ?Lynch Arribálzaga, 1880: 52, 1883a: 144 (cat.); Williston, 1891: 78 (cat.); Aldrich, 1905: 268 (cat.); Kertész, 1909: 163 (cat.); Hermann, 1912: 156; Curran, 1930b: 16 (in key), 1935: 6 (in key); Hull, 1962 (1): 373; Martin, 1965a: 25; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.); Papavero & Ibáñez Bernal, 2001: 114 (hist. Mexican dipterol.). Type, MUN.
- bequaerti** Bromley, 1934b: 339. Type-locality: Brazil, Pará, Santarém. Refs. – Curran, 1935: 6 (in key); Bromley, 1946: a 109 (cat.); Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.).
- bigoti** Bellardi, 1861: 120 (20). Type-locality: "Mexico". Refs. – Aldrich, 1905: 268 (cat.); Kertész, 1909: 163 (cat.); Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.); Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, OXF. Type, TORO.
- ?**brevicornis** Macquart, 1838: 76 (1839: 192) (*Laphria*). Type-locality: "Brazil". Refs. – Walker, 1855: 568 (n. comb.); Williston, 1891: 78 (cat.); Kertész, 1909: 163 (cat.); Bromley, 1946: 109 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.). Type, lost.
- cerverai** Bromley, 1929a: 282, fig. 6. Type-locality: Cuba, Soledad. Refs. – Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Alayo Dalmau & García Ávila, 1983: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, USNM.
- ciguaya** Scarbrough & Perez-Gelabert, 2006: 24 (in key), 25, figs. 8-12, 119. Type-locality: Dominican Republic, San Juan, road-232, road San Juan/Vallejuelo, near river. Distr. – Dominican Republic (Azua, Barahona, Elías Piña, Independencia, Montecristi, Pedernales, Peravia, Sánchez Ramírez, San Juan). Type, USNM.
- coxalis** Curran, 1930b: 16 (in key), 18. Type-locality: Brazil, Mato Grosso do Sul, Corumbá. Refs. – Curran, 1935: 7 (in key); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, AMNH.
- cyanescens** Rondani, 1848: 92. Type-locality: "Brazil" [i. e., Pará, Belém; cf. Papavero, 1973d: 342-343]. Refs. – Williston, 1891: 78 (cat.); Kertész, 1909: 163 (cat.); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). TYPE MZUN.
- danforthi** Curran, 1935: 6 (in key), 8. Type-locality: Puerto Rico, Coamo Springs. Refs. – Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, AMNH.
- dasypus** (Wiedemann), 1828: 527 (misprinted as 257) (*Laphria*). Type-locality: "Brazil". Refs. – Walker, 1855: 568; Schiner, 1866: 693 (?*Atomosia*, ?*Atractia*); Williston, 1891: 79 (cat.); ?*Atomosia*, ?*Atractia*; Kertész, 1909: 166 (?*Atractia*); Bromley, 1946: 108 (cat.); Hull, 1962: 386, text-fig. 27, figs. 228, 281, 639, 1280, 1289 (*Paratractia*); Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.; n. comb.); Castillo, Jerez & Artigas, 1994: (*Paratractia*; egg chorion). Type, BZM [no. 2278; cf. Schumann, 1973: 101].
- fredericoi** Carrera, 1952a: 209. Type-locality: Brazil, São Paulo, São Paulo (Ipiranga). Refs. – Carrera & d'Andretta, 1952: 255 (preys); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, MZUSP.
- frontalis** Curran, 1930b: 16 (in key), 17. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Curran, 1935: 6 (in key); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, AMNH.
- geniculata** (Wiedemann), 1821: 241 (*Laphria*). Type-locality: "Brazil". Refs. – Wiedemann, 1828: 528 (*Laphria*); Walker, 1855: 568; Schiner, 1866: 693, 1868: 169; Wulp, 1882b: 105; Williston, 1891: 78 (cat.); Kertész, 1909: 164 (cat.); Hermann, 1912: 142, fig. 52; Curran, 1930b: 16 (in key); Engel, 1930: 467; Curran, 1935: 6 (in key); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, ?NMW.
- hondurana** James, 1953: 53. Type-locality: Honduras, Zamorano (Escuela Agrícola Panamericana). Refs. – Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, MICH.
- jagua** Scarbrough & Perez-Gelabert, 2006: 22 (in key), 28, figs. 13-17. Type-locality: Dominican Republic, Independencia, Road-272, caseta 1, Parque Nacional Sierra de Bahoruco. Distr. – Dominican Republic (Independencia, Pedernales, Peravia). Type, USNM.
- jimagua** Scarbrough & Perez-Gelabert, 2006: 22 (in key), 31, figs. 18-22. Type-locality: Dominican Republic, Pedernales, Parque Nacional Jaragua, trail to Carlitos, ca. 6 km south of highway 44, 106 m. Distr. – Dominican Republic (Pedernales). Type, USNM.
- limbiventris** Thomson, 1869: 466. Type-locality: Uruguay, Montevideo. Refs. – Lynch Arribálzaga, 1883a: 144 (cat.); Williston, 1891: 78 (cat.); Kertész, 1909: 164 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, STOCKHOLM.
- limbativentris* Lynch Arribálzaga, 1880: 50, emend.
- lineata** Curran, 1930b: 16 (in key), 19. Type-locality: Brazil, Mato Grosso do Sul, Corumbá. Refs. – Curran, 1935: 7 (in key); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, AMNH.
- macquarti** Bellardi, 1861: 120 (20). Type-locality: "Mex. ico". Distr. – Mexico (Gerrero, Tabasco, Tamaulipas, Veracruz, Yucatán), Costa Rica Refs. – Williston, 1901: 316; Aldrich, 1905: 269 (cat.); Kertész, 1909: 164 (cat.); Hermann, 1912: 163; Curran, 1930b: 16 (in key), 1935: 6 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 46 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.); Coronado Blanco & Ruiz Cancino, 1999: 81 (as predator of *Unaspis citri* (Comstock, 1883) (Hemiptera, Homoptera, Diaspididae)); Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, TORO.
- maestrae** Bromley, 1929a: 282, fig. 7. Type-locality: Cuba, Sierra Maestra. Refs. – Curran, 1935: 6 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Alayo Dalmau & García Ávila, 1983: 47 (cat.); Artigas, Papavero & Serra, 1991:

- 59 (cat.). Type, USNM.
- melanopogon** Hermann, 1912: 144, fig. 53. Type-locality: U. S. A., Texas. Distr. – U. S. A. (Arizona, Colorado, Iowa, Texas), Mexico, Honduras. Refs. – Curran, 1930b: 15, 16 (in key), 17, 1935: 5, 6 (in key); Hull, 1962 (1): 373; Martin & Wilcox, 1965: 387 (cat.); Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, ?NMW.
- metallescens** Hermann, 1912: 150, fig. 57. Type-locality: Argentina, Mendoza. Refs. – Curran, 1930b: 15 (in key), 1935: 5 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Syntypes, MUN, NMW.
- metallica** Bromley, 1929a: 283. Type-locality: Cuba, Sierra Maestra, Palma Mocha. Refs. – Curran, 1935: 6 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, USNM.
- mucida** Osten Sacken, 1887: 184. Type-locality: Mexico, Sinaloa, Presidio. Refs. – Williston, 1901: 315; Aldrich, 1905: 269 (cat.); Kertész, 1909: 164 (cat.); Curran, 1930b: 16 (in key), 1935: 6 (in key); Hull, 1962 (1): 373; Martin & Wilcox, 1965: 387 (cat.); Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, BMNH.
- nigroaenea** Walker, 1851: 154. Type-locality: Colombia, Bogotá (in label of type). Refs. – Walker, 1855: 569; Williston, 1891: 78 (cat.); Kertész, 1909: 164 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, BMNH.
- nuda** Hermann, 1912: 147, fig. 55. Type-locality: Trinidad, Belmont (in label of type). Refs. – Curran, 1930b: 15 (in key), 1935: 5 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, MUN.
- panamensis** Curran, 1930: 16 (in key), 17. Type-locality: Panama, Canal Zone, Barro Colorado I. Refs. – Curran, 1935: 6 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, AMNH.
- parva** (Bigot), 1857a: 330, pl. 20, figs. 2, 2a (*Dasytopogon*; preocc. Rondani, 1850). Type-locality: “Cuba”. Refs. – Aldrich, 1905: 267 (cat.; *Dasytopogon*); Kertész, 1909: 130 (cat.; *Dasytopogon*); Bromley, 1929a: 281, pl. 1, fig. 8 (n. comb.); Curran, 1935: 5 (in key); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 47 (cat.). Type, MNHNP (almost entirely destroyed, only right pleura and one wing and tergites remain).
- pilipes** Thomson, 1869: 465. Type-locality: Argentina, Buenos Aires. Refs. – Wulp, 1882b: 105; Lynch Arribálzaga, 1883a: 144 (cat.); Kertész, 1909: 165 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, STOCKHOLM.
- pilosipes* Lynch Arribálzaga, 1880: 52, error.
- pubescens** Bromley, 1929a: 281, pl. 1, fig. 3. Type-locality: Cuba, Soledad. Type, Refs. – Curran, 1935: 5 (in key); Hull, 1962 (1): 373, fig. 229; Martin & Papavero, 1970: 47 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). MCZ.
- puella** (Wiedemann), 1828: 531 (*Laphria*). Type-locality: unknown. Refs. – Rondani, 1848: 91; Walker, 1855: 570; Schiner, 1866: 693; Williston, 1883: pl. 2, fig. 8; Osten Sacken, 1891: 418; Johnson, 1903: 114; Aldrich, 1905: 269 (cat.); Kertész, 1909: 165 (cat.); Hermann, 1912: 141, figs. 51a, 51b; Curran, 1930b: 16 (in key), 1935: 6 (in key); Fatrtigg, 1945: 18; Bromley, 1950: 233; Hull, 1962 (1): 373, figs. 223, 637, 1309, 1318, 2110, 2121-2123, 2153, 2156; Martin & Wilcox, 1965: 387 (cat.); Martin & Papavero, 1970: 47 (cat.); Scarbrough & Sraver, 1979: 630 (predatory behavior and prey); Alayo Dalmau & Garcia Ávila, 1983: 47 (cat.); Artigas, Papavero & Serra, 1991: 59, figs. 75-81 (cat.). Type, NMW.
- incisuralis* Macquart, 1838: 76 (1839: 192), pl. 7, fig. 1. Type-locality: “Cuba”. Refs. – Walker, 1855: 567; Bigot, 1857a: 330; Aldrich, 1905: 268 (cat.); Kertész, 1909: 164 (cat.); Bromley, 1929a: 280; Curran, 1935: 6 (in key); Hull, 1962 (1): 373. Type, MNHNP.
- rica** Curran, 1935: 6 (in key), 9. Type-locality: Puerto Rico, Mayaguez. Refs. – Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, AMNH.
- rosalesi** Carrera & Machado-Allison, 1963: 242, fig. 1. Type-locality: Venezuela, Carabobo, Maruiara. Ditr. – Venezuela (Aragua, Bolívar, Carabobo, Lara, Mérida, Portuguesa). Refs. – Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, FAUCV.
- rufipes** Macquart, 1847: 55 (1847: 39). Type-locality: U. S. A., Pennsylvania. Distr. – U. S. A (Pennsylvania, Georgia, Florida, Texas), Mexico. Refs. – Hermann, 1912: 157, fig. 60; Curran, 1930b: 16 (in key); Bromley, 1934c: 163; Curran, 1935: 7 (in key); Fattig, 1945: 18; Martin & Wilcox, 1965: 387 (cat.); Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, MNHNP.
- eupoda* Bigot, 1878: 234 (*Cormansis*). Type-locality: “Mexico”. Refs. – Williston, 1883: 33 (*Cormansis*); Osten Sacken, 1887: 185; Aldrich, 1905: 268 (cat.); Kertész, 1909: 163 (cat.); Hermann, 1912: 161 (as *rufipes* var.); Curran, 1930b: 16 (in key), 1935: 7 (in key); Hull, 1962 (1): 373; Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipterol). Type, OXF.
- soror* Bigot, 1878: 236. Type-locality: “Mexico”. Refs. – Williston, 1883: 33; Johnson, 1903: 114; Aldrich, 1905: 269 (cat.); Kertész, 1909: 166 (cat.); Hermann, 1912: 161 (as *rufipes* var.); Curran, 1930b: 16 (in key), 1935: 6 (in key); Hull, 1962 (1): 373; Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipterol). Type, OXF.
- punctifera* Hermann, 1912: 161 (as *rufipes* var.). Type-locality: “Mexico”. Refs. – Curran, 1930b: 16 (in key), 1935: 6 (in key). Type, MUN.
- ?**scoriacea** (Wiedemann), 1828: 529 (*Laphria*). Type-locality: “Mexico”. Refs. – Walker, 1855: 569; Schiner, 1866: 693; Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 165 (cat.); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, SENCK [cf. Geller-Grimm, 1999: 214].

- selene** Curran, 1935: 5 (in key), 7. Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 47 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, AMNH.
- setosa** Hermann, 1912: 145, fig. 54. Type-locality: “Brazil”. Refs. – Curran, 1930b: 16 (in key), 1935: 6 (in key); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 48 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Syntypes, MIL, MUN.
- similis** Bigot, 1857a: 330, pl. 20, figs. 4, 4a. Type-locality: “Cuba”. Refs. – Aldrich, 1905: 165 (cat.); Kertész, 1909: 165 (cat.); Hermann, 1912: 149, fig. 56; Bromley, 1929a: 282, pl. 1, fig. 15 (n. comb.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 48 (cat.); Alayo Dalmau & García Ávila, 1983: 47 (cat.).
- sp., near dasypus** (Wiedemann, 1828) Artigas, Papavero & Serra, 1991: 81, figs. 82-86.
- tenuis** Curran, 1930b: 16 (in key), 19. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Curran, 1935: 7 (in key); Bromley, 1946: 108 (cat.); Carrera, 1947d: 269 (preys); Carrera & d’Andretta, 1952: 254 (preys); Hull, 1962 (1): 373; Martin & Papavero, 1970: 48 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, AMNH.
- tibialis** Macquart, 1846: 204 (1846: 76). Type-locality: México, Yucatán, Mérida. Refs. – Walker, 1855: 567; Wulp, 1882b: 105; Williston, 1891: 78 (cat.); Aldrich, 1905: 269 (cat.); Kertész, 1909: 166 (cat.); Curran, 1930b: 15 (in key), 1935: 5 (in key); Bromley, 1934b: 339, 1946a: 108 (cat.); Carrera, 1947d: 269 (prey); Carrera & d’Andretta, 1952: 255 (preys), 1961: 71 (preys); Hull, 1962 (1): 373; Artigas, Papavero & Serra, 1991: 59 (cat.); Papavero & Ibáñez Bernal, 2001: 85 (hist. Mexican dipterol.). Type, lost.
- unicolor** Macquart, 1838: 74 (1839: 190), pl. 7, figs. 2-3. Type-locality: Brazil, Rio Grande do Sul. Refs. – Walker, 1855: 569; Williston, 1891: 78 (cat.); Kertész, 1909: 166 (cat.); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin & Papavero, 1970: 48 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, MNHNP.
- venustula** Lynch Arribálzaga, 1880: 50. Type-locality: Argentina, Buenos Aires, Las Conchas and Gran Chaco. Distr. – Paraguay, Argentina. Refs. – Wulp, 1882b: 106; Lynch Arribálzaga, 1883a: 144 (cat.); Williston, 1891: 78 (cat.); Kertész, 1909: 166 (cat.); Hermann, 1912: 162; Curran, 1930b: 15 (in key); Engel, 1930: 467; Curran, 1935: 5 (in key); Carrera, 1947a: 44; Carrera & d’Andretta, 1952: 255 (prey), 1961: 71 (preys); Hull, 1962 (1): 373; Martin & Papavero, 1970: 48 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.). Type, lost.
- xanthopus** (Wiedemann), 1828: 529 (*Laphria*). Type-locality: “Brazil”. Refs. – Walker, 1855: 539 (*Laphria*); Schiner, 1866: 693; Wulp, 1882b: 105; Williston, 1891: 78 (cat.); Kertész, 1909: 166 (cat.); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 373; Martin, 1965a: 25; Martin & Papavero, 1970: 48 (cat.); Artigas, Papavero & Serra, 1991: 59 (cat.).
- yurabia** Scarbrough & Perez-Gelabert, 2006: 22 (in key), 33, figs. 23-27. Type-locality: Dominican Republic, San Juan, road-233, few km east of Vallejuelo, 704 m. Distr. – Dominican Republic (Pedernales, San Juan). Type, USNM.

Genus *Atomosiella* Wilcox, 1937

Atomosiella Wilcox, 1937: 40. Type-species, *Atomosia antennata* Banks, 1920 (orig. des.)

antennata (Banks), 1920: 66 (*Atomosia*). Type-locality: U. S. A., Arizona. Distr. – U. S. A. (Arizona, California, New Mexico), Mexico (Baja California). Refs. – Wilcox, 1937: 41, fig. 3 (n. comb.); Hull, 1962 (1): 399, figs. 220, 671, 1334, 1343; Martin & Wilcox, 1965: 387 (cat.); Artigas, Papavero & Serra, 1991: 58 (cat.). Type, ?MCZ.

Genus *Atoniomyia* Hermann, 1912

Atonia Williston, 1889: 257 (preocc. Gistel, 1848). Type-species, *Atomosia mikii* Williston, 1886 (sub. des., Williston, 1901: 316).

Atoniomyia Hermann, 1912: 81 (*nom. nov.* for *Atonia* Williston, 1889). Type-species, *Atomosia mikii* Williston, 1886 (aut.). Ref. – Artigas, Papavero & Serra, 1991: 78, figs. 66-72.

Neatonia Bromley, 1935a: 130, unavailable name (no type-species designated).

albiceps (Hermann), 1912: 85 (key; as *albifacies*, error), 97, fig. 38 (*Atonia*). Type-locality: Brazil, Mato Grosso. Refs. – Bromley, 1946: 108 (cat.); Carrera, 1946b: 120; Hull, 1962 (1): 415; Martin & Papavero, 1970: 54 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, BUD.

ancylocera (Schiner), 1868: 170 (*Atomosia*). Type-locality: “Venezuela”. Distr. – Venezuela, Brazil (São Paulo). Refs – Williston, 1889: 257 (*Atomosia*), 1891: 78 (cat.; ?*Atonia*); Kertész, 1909: 167 (cat.; *Atonia*); Hermann, 1912: 88, figs. 31-32 (*Atonia*); Carrera, 1946b: 119, figs. 2, 13, 17; Hull, 1962 (1): 415; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, NMW.

brevistylata (Williston), 1901: 316, pl. 6, fig. 1 (*Atonia*). Type-locality: Mexico, Tabasco, Teapa. Distr. – U. S. A. (Arizona), Mexico (Tabasco). Refs. – Aldrich, 1905: 268 (cat.; *Atonia*); Kertész, 1909: 167 (cat.; *Atonia*); Hermann, 1912: 84;

- Hull, 1962 (1): 415; Martin & Wilcox, 1965: 387 (cat.); Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Type, BMNH.
- fulvipes** Carrera, 1946b: 122, figs. 7, 10, 20. Type-locality: Brazil, Mato Grosso do Sul, Salobra. Refs. – Hull, 1962 (1): 415; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, MZUSP.
- grossa** Carrera, 1946b: 125, figs. 9, 12, 19. Type-locality: Brazil, São Paulo, Severínia. Refs. – Hull, 1962 (1): 415; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, MZUSP.
- hispidella** (Hermann), 1912: 95, fig. 37 (*Atonia*). Type-locality: Brazil, São Paulo. Refs. – Bromley, 1946: 108 (cat.); Carrera, 1946b: 120, figs. 8, 15, 18; Hull, 1962 (1): 415, figs. 276, 276a, 672, 1333, 1342, 2047; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Syntypes, MIL, MUN.
- laterepunctata** (Hermann), 1912: 93, fig. 36 (*Atonia*). Type-locality: Peru, Huánuco, mouth of Pachitea river; Cuzco, Umuhanckiali at Urubamba river and Pasco, Puerto Yessup; Bolivia, Mapiri, Sarampiuni; Peru, Cuzco, Vilcanota. Distr. – Peru (Cuzco, Huánuco, Pasco), Bolivia. Refs. – Carrera & d'Andretta, 1950: 161 (misident.); Hull, 1962 (1): 415; Martin, 1965a: 25; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Syntypes, NMW, MIL, BUD, BMNH, TP SMTD [cf. Kallweit, 1992: 196].
- lateropunctata* Carrera & Machado-Allison, 1963: 244, error.
- lateripunctata* Hull, 1962 (2): 617 (caption to fig. 276c), error.
- mikii** (Williston), 1886: 290 (*Atomosia*). Type-locality: Dominican Republic, Santo Domingo. Distr. – Dominican Republic (Azua, Barahona, Santo Domingo). Refs. – Williston, 1889: 257 (*Atonia*); Aldrich, 1905: 268 (cat. *Atonia*); Hull, 1962 (1): 415; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.); Scarbrough & Perez-Gelabert, 2006: 36, figs. 28-32. Type, ?lost.
- mickii* Hermann, 1912: 83, error.
- mollis** (Hermann), 1912: 88, fig. 33 (*Atonia*). Type-locality: Peru, Pasco, Unini at Ucayali river, and Cuzco, Vilcanota. Distr. – Peru (Cuzco, Pasco). Refs. – Carrera, 1946b: 120, figs. 6, 14; Hull, 1962 (1): 415; Martin & Papavero, 1970: 53 (cat.); Lamas M., 1972: 311 (cat.); Fisher, 1984: 31; Artigas, Papavero & Serra, 1991: 60 (cat.). Syntypes, BUD, MUN, SMTD [cf. Kallweit, 1992: 196].
- pinguis** (Hermann), 1912: 90, fig. 34 (*Atonia*). Type-locality: Peru, Huánuco, mouth of Pachitea river. Distr. – Peru (Huánuco). Refs. – Carrera & d'Andretta, 1950: 161, figs. 5, 22 (misident.); Hull, 1962 (1): 415; Martin, 1965a: 26; Martin & Papavero, 1970: 53 (cat.); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Syntypes, MUN, SMTD [cf. Kallweit, 1992: 197].
- scalarata** (Hermann), 1912: 92, fig. 34 (*Atonia*). Type-locality: Peru, Cuzco, Rosalina at Urubamba river; Bolivia, Mapiri, Chimate. Distr. – Peru (Cuzco), Bolivia. Refs. – Hull, 1962 (1): 415, fig. 276c; Martin & Papavero, 1970: 53 (cat.); Lamas M., 1972: 311 (cat.); Fisher, 1984: 31; Artigas, Papavero & Serra, 1991: 60 (cat.). Syntypes, NMW, MUN, BMNH, SMTD [cf. Kallweit, 1992: 197].
- setigera** (Hermann), 1912: 86, pl. 3, figs. 24-25 (*Atonia*). Type-locality: “Brazil”. Refs. – Hull, 1962 (1): 415; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, NMW.
- sp.** Scarbrough & Perez-Gelabert, 2006: 38 (Dominican Republic: Barahona).
- viduata** (Wiedemann), 1819: 50 (*Laphria*). Type-locality: “Brazil”. Distr. – Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Type, NMW. Refs. – Wiedemann, 1821: 241 (*Laphria*), 1828: 528 (*Laphria*); Walker, 1855: 538 (*Laphria*); Schiner, 1866: 693 (*Atomosia*), 1868: 378 (*Atomosia*); Williston, 1891: 78 (cat. *Atomosia*); Kertész, 1909: 166 (cat. *Atomosia*); Hermann, 1912: 85, fig. 30 (*Atonia (Laphria)*); Bromley, 1946: 108 (cat.); Carrera, 1946b: 117, figs. 1, 11, 16; Hull, 1962 (1): 373 (*Atomosia*), 415 [*Atoniomyia*]; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.); Geller-Grimm, 1999: 215 (material in SENCK). Type, NMW.

Genus *Atractia* Macquart, 1838

Atractia Wiedemann, 1838: 151 (1839: 267). Type-species, *Asilus psilogaster* Wiedemann, 1828 (mon.).

- psilogaster** (Wiedemann), 1828: 456 (*Laphria*). Type-locality: “Brazil”. Refs. – Macquart, 1838: 152; Walker, 1855: 754; Schiner, 1866: 685, 1868: 412; Williston, 1891: 89 (cat.); Kertész, 1909: 167 (cat.); Hermann, 1912: 174, fig. 65; Bromley, 1946: 108 (cat.); Hull, 1962 (1): 379; Martin & Papavero, 1970: 49 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Syntypes, NMW, BZM [no. 2272; cf. Schumann, 1973: 147].
- coronata** Schiner, 1867: 412. Type-locality: “Brazil”. Refs. – Williston, 1891: 89 (cat.); Kertész, 1909: 166 (cat.); Hermann, 1912: 174 (syn.); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 379. Type, NMW.
- psylogaster** Carrera & Vulcano, 1961: 72 (preys), error.
- pulverulenta** Schiner, 1867: 412. Type-locality: “Brazil”. Refs. – Williston, 1891: 89 (cat.); Kertész, 1909: 167 (cat.); Hermann, 1912: 176, fig. 66; Bromley, 1946: 110 (cat.); Hull, 1962 (1): 379; Martin & Papavero, 1970: 49 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, NMW.

Genus **Bathropsis** Hermann, 1912

Bathropsis Hermann, 1912: 68. Type-species, *Bathropsis peruviana* Hermann, 1912 (orig. des.).

basalis Curran, 1930b: 6, fig. 1. Type-locality: Panama, Canal Zone, Barro Colorado I. Refs. – Hull, 1962 (1): 407; Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, AMNH.

delgadoi Kaletta, 1978: 56. Type-locality: Venezuela, Aragua, Hacienda El Periquito, Ocumare de la Costa. Ref. – Artigas, Papavero & Serra, 1991: 60 (cat.). Type, FAUCV.

peruviana Hermann, 1912: 69, fig. 23, pl. 2, figs. 13–14. Type-locality: Peru, Cuzco, Vilcanota. Distr. – Peru (Cuzco). Refs. – Hull, 1962 (1): 407, figs. 251, 665, 1294, 1303, 1349, 1358; Martin & Papavero, 1970: 53 (cat.); Lamas M., 1972: 311; Artigas, Papavero & Serra, 1991: 60 (cat.). Type, MUN.

plazai Kaletta, 1986: 264. Type-locality: Venezuela, Aragua, Rancho rande, 1100 m. Distr. – Venezuela (Aragua). Type, IZAUCV.

Genus **Cerotainia** Schiner, 1866

Cerotainia Schiner, 1866: 662 (key), 673 (1868: 170, second erection of genus). Type-species, *Laphria xanthoptera* Wiedemann, 1828 (orig. des.).

Ceratotaenia Lynch Arribálzaga, 1880: 52, emend.

Cyphotomyia Williston, 1889: 257. Type-species, *Cyphotomyia lynchii* Williston, 1889 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 60 (syn.).

Protichisma Hermann, 1912: 35. Type-species, *Protichisma longimanus* Hermann, 1912 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 60 (syn.).

albibarbis (Curran), 1930b: 7 (*Protichisma*). Type-locality: Panama, Canal Zone, France Field. Ref. – Hull, 1962 (1): 392 (*Protichisma*); Martin & Papavero, 1970: 51 (cat.; *Protichisma*); Artigas, Papavero & Serra, 1991: 60 (cat.; n. comb.). Type, AMNH.

argyropasta Hermann, 1912: 61. Type-locality: Peru, Ucayali river. Peru (Loreto). Refs. – Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 51 (cat.); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, TP SMTD [cf. Kallweit, 1992: 194].

argyropus Schiner, 1868: 170. Type-locality: “Colombia” (in error; actually Venezuela in type label). Distr. – Venezuela, Peru. Refs. – Williston, 1891: 78 (cat.); Kertész, 1909: 168 (cat.); Hermann, 1912: 51, fig. 15; Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 51 (cat.); Lamas M., 1972: 311 (cat.); Kaletta, 1986: 258; Artigas, Papavero & Serra, 1991: 60 (cat.). Type, NMW.

argyropyga Hermann, 1912: 52, fig. 16. Type-locality: “Peru”. Distr. – Peru. Refs. – Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 51 (cat.); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 60 (cat.). Type, MUN.

aurata Schiner, 1868: 171. Type-locality: “Colombia”. Ref. – Williston, 1891: 78 (cat.); Kertész, 1909: 168 (cat.); Hermann, 1912: 54, fig. 18; Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 51 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, NMW.

bella Schiner, 1867: 380. Type-locality: “Brazil” [probably from the state of Rio de Janeiro, as it was collected by Bescke; cf. Papavero, 1971e: 87]. Refs. – Williston, 1891: 78 (cat.); Kertész, 1909: 168 (cat.); Hermann, 1912: 46; Curran, 1930b: 12 (in key); Bromley, 1946: 110 (cat.); Hull, 1962 (1): 395; Martin & Papavero, 1970: 51 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, NMW.

bolivari Kaletta, 1986: 258. Type-locality: Venezuela, Carabobo, Borburata. Distr. – Venezuela (Carabobo). Type, IBTUCV.

brasiliensis Schiner, 1867: 379. Type-locality: “Brazil” (probably from the state of Rio de Janeiro, as it was collected by Bescke; cf. Papavero, 1971e: 87]. Distr. – Brazil (Rio de Janeiro, São Paulo, Paraná, Santa Catarina). Refs. – Williston, 1891: 78 (cat.); Kertész, 1909: 168 (cat.); Hermann, 1912: 51; Curran, 1934a: 3 (in key); Bromley, 1946: 110 (cat.); Carrera, 1958a: 145; Hull, 1962 (1): 395; Martin & Papavero, 1970: 51 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, NMW.

camposi Curran, 1934a: 3. Type-locality: Ecuador, Guayas, Posorja. Refs. – Hull, 1962 (1): 395; Martin & Papavero, 1970: 51 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, AMNH.

clavijoi Kaletta, 1986: 263. Type-locality: Venezuela, Aragua, Camatagua. Distr. – Venezuela (Aragua). Type, IZAUCV.

dasythrix Hermann, 1912: 48, fig. 13. Type-locality: Peru, Huánuco, mouth of Pachitea river; Bolivia, Mapiri, Sarapiuni, 700 m. Distr. – Venezuela (Aragua), Peru (Huánuco), Bolivia. Refs. – Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Carrera & Machado-Allison, 1963: 244; Martín & Papavero, 1970: 51 (cat.); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Syntypes, MUN, SMTD [cf. Kallweit, 1992: 194].

debilis Hermann, 1912: 48, fig. 21. Type-locality: Peru, Huánuco, mouth of Pachitea river, 150 m; Junín, Chanchamayo, 800 m; Bolivia, Sarapiuni; Peru, Cuzco, Vilcanota. Distr. – Venezuela (Cojedes), Peru (Cuzco, Huánuco, Junín), Bolivia. Refs. – Curran, 1930b: 12 (in key), 1934a: 3 (in key); Hull, 1962 (1): 395; Carrera & Machado-Allison, 1963: 245;

- Martin & Papavero, 1970: 52 (cat.); Lamas M., 1972: 311 (cat.); Fisher, 1984: 32; Artigas, Papavero & Serra, 1991: 61 (cat.). Syntypes, MUN, BUD, NMW, BMNH, SMTD [cf. Kallweit, 1992: 195].
- dubia** Bigot, 1878: 238. Type-locality: "Mexico". Refs. – Williston, 1883: 33; Osten Sacken, 1887: 185 (*Atomosia*); Aldrich, 1905: 268 (cat.); Kertész, 1909: 168 (cat.); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.); Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipterol). Type, OXF.
- feminea** Curran, 1930b: 12 (in key), 14. Type-locality: Panama, Canal Zone, Corzal. Refs. – Curran, 1934a: 3 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, AMNH.
- flavipes** Hermann, 1912: 57, fig. 20. Type-locality: Peru, Junín, Chanchamayo, 800 m; Paraguay, Asunción. Distr. – Peru (Junín), Paraguay. Refs. – Curran, 1930b: 12 (in key), 1934a: 3 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Syntypes, MUN, BUD, SMTD [cf. Kallweit, 1992: 195].
- gerulewiczii** Kaletta, 1986: 161. Type-locality: Venezuela, Aragua, Rancho Grande, 1100 m. Distr. – Venezuela (Aragua). Type, IZAUCV.
- jamaicensis** Johnson, 1919: 430 (*as macrocera* var.). Type-locality: "Jamaica". Refs. – Gowdey, 1926: 77 (*as macrocera* var.); Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Farr, 1965: 9, figs. 3, 4a; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, MCZ.
- laticeps** Bromley, 1929a: 279, pl. 1, fig. 2. Type-locality: Cuba, Zapata, Santo Tomás. Refs. – Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Alayo Dalmau & García Ávila, 1983: 47 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.; syn.). Type, USNM.
- jaimacensis** Hull, 1962 (1): 395, error.
- jolyi** Kaletta, 1986: 260. Type-locality: Venezuela, Miranda, La Tumusa, Valle del Tuy. Distr. – Venezuela (Miranda). Ref. – Rueda, 2004. Type, IZAUCV.
- leonina** Hermann, 1912: 49, fig. 14. Type-locality: Argentina, Mendoza. Distr. – Argentina (Córdoba, Mendoza, Tucumán). Refs. – Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61, figs. 89-97 (cat.); Rueda, 1998: 170, figs. 1a-c. Type, MUN.
- longimana** (Hermann), 1912: 37, pl. 1, figs. 7-8 (*Protichisma*). Type-locality: Peru, Madre de Dios, Callanga. Distr. – Peru (Huánuco, Madre de Dios). Refs. – Hull, 1962 (1): 392, figs. 278, 649, 1293, 1302, 1346, 1355 (*Protichisma*); Martin & Papavero, 1970: 51 (cat.; *Protichisma*); Lamas M., 1972: 311 (cat.; *Protichisma*); Artigas, Papavero & Serra, 1991: 61 (cat.; n. comb.). Type, BUD.
- lynchii** (Williston), 1889: 158 (*Cyphotomyia*). Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Distr. – Peru (Cuzco), Bolivia, Brazil (Mato Grosso, São Paulo). Refs. – Williston, 1891: 79 (cat.; *Cyphotomyia*); Kertész, 1909: 169 (cat.; *Cyphotomyia*); Hermann, 1912: 68; Bromley, 1946: 110 (cat.; *Cyphotomyia*); Hull, 1962 (1): 397, figs. 267, 641, 1345, 1354, 2157 (*Cyphotomyia*); Martin & Papavero, 1970: 52 (cat.; *Cyphotomyia*); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 61, figs. 98-104 (cat.; n. comb.). Type, ?AMNH.
- marginata** Hermann, 1912: 62. Type-locality: Peru, Pasco, Unini, Ucayali river (as Pachitea river, error). Refs. – Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Lamas M., 1972: 311 (cat.); Fisher, 1984: 32; Artigas, Papavero & Serra, 1991: 61 (cat.). Type, SMTD [cf. Kallweit, 1992: 196].
- melanosoma** Scarbrough & Knutson, 1989: 520, figs. 1-2. Type-locality: Dominica, Clarke Hall, near Layou River. Distr. – Dominica. Type, USNM.
- minima** Curran, 1930b: 11 (in key), 12. Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. – Panama, Venezuela. Refs. – Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Kaletta, 1986: 157; Artigas, Papavero & Serra, 1991: 61 (cat.). Type, AMNH.
- nigra** Bigot, 1878: 238. Type-locality: "Mexico". Refs. – Williston, 1883: 33; Osten Sacken, 1887: 185 (*Atomosia*); Aldrich, 1905: 268 (cat.); Kertész, 1909: 168 (cat.); Curran, 1930b: 16 (in key; *Atomosia*), 1935: 6 (in key; *Atomosia*); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.); Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipterol). Type, OXF.
- nigripennis** (Bellardi), 1861: 119 (19) (*Atomosia*). Type-locality: "Mexico". Refs. – Aldrich, 1905: 268 (cat.); Kertész, 1909: 168 (cat.); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.); Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol). Type, TORO.
- ornatipes** James, 1953: 54. Type-locality: Honduras, Zamorano. Distr. – Honduras, Venezuela. Refs. – Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Kaletta, 1986: 257; Artigas, Papavero & Serra, 1991: 61 (cat.). Type, WASH.
- propinqua** Schiner, 1868: 171. Type-locality: "Colombia" (in error; actually Venezuela in type's label). Refs. – Williston, 1891: 78 (cat.); Hermann, 1912: 56, fig. 19; Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.). Type, NMW.
- porpinqua** Kertész, 1909: 168 (cat.), error.
- rhopalocera** (Lynch Arribálzaga), 1882: 189 (*Ceratotaenia*). Type-locality: Argentina, Buenos Aires, Chacabuco. Refs. – Williston, 1891: 79 (cat.); Kertész, 1909: 168 (cat.); Hermann, 1912: 54, fig. 17; Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, lost.
- sarae** Rueda, 1998: 172, figs. 2a-c. Type-locality: Argentina, Catamarca, El Rodeo. Distr. – Argentina (Catamarca). Type, IML.
- sola** Scarbrough & Perez-Gelabert, 2006: 39, figs. 33-36, 120. Type-locality: Dominican Republic, Pedernales, 25 km north of Cabo Rojo, 679 m. Distr. – Dominican Republic (Hato Mayor, Independencia, La Altagracia, Pedernales). Type, USNM.

- unicolor** Hermann, 1912: 64, fig. 22. Type-locality: “Peru”. Distr. – Peru. Refs. – Curran, 1930b: 12 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, MUN.
- violaceithorax** (Lynch Arribálzaga), 1880: 52 (*Ceratotaenia*). Type-locality: Argentina, Buenos Aires. Refs. – Lynch Arribálzaga, 1882: 145 (*Ceratotaenia*), 1883a: 145 (cat.; *Ceratotaenia*); Williston, 1891: 79 (cat.); Kertész, 1909: 168 (cat.); Hermann, 1912: 54; Curran, 1930b: 12 (in key); Bromley, 1946: 110 (cat.); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, lost.
- willistoni** Curran, 1930b: 12 (in key), 14. Type-locality: Brazil, Mato Grosso, Chapada dos Guimarães. Refs. – Curran, 1934a: 3 (in key); Bromley, 1946: 110 (cat.); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, AMNH.
- xanthoptera** (Wiedemann), 1828: 530 (*Laphria*). Type-locality: unknown (“America” on type label). Refs. – Walker, 1855: 562 (*Laphria*); Schiner, 1866: 693; Kertész, 1909: 168 (cat.); Hermann, 1912: 46; Curran, 1930b: 11 (in key); Hull, 1962 (1): 395; Martin & Papavero, 1970: 52 (cat.); Artigas, Papavero & Serra, 1991: 61 (cat.). Type, NMW.

Genus **Cyphomyiactia** Artigas, Papavero & Serra, 1991

- Cyphomyiactia** Artigas, Papavero & Serra, 1991: 61. Type-species, *Cyphomyiactia costai* Artigas, Papavero & Serra, 1991 (orig. des.).
- costai** Artigas, Papavero & Serra, 1991: 62. Type-locality: Brazil, Goiás, Goiânia (Campinas). Distr. – Brazil (Goiás). Type, MZUSP.

Genus **Dissmeryngodes** Hermann, 1912

- Dissmeryngodes** Hermann, 1912: 75. Type-species, *Laphria antica* Wiedemann, 1828 (orig. des.).
- Centrolaphria* Enderlein, 1914a: 241. Type-species, *Centrolaphria columbiana* Enderlein, 1914 (orig. des.) = *Dissmeryngodes nigripes* (Macquart, 1838). Ref. – Artigas, Papavero & Serra, 1991: 62 (syn.).
- Catonomyia* Hull, 1962 (1): 380. Type-species, *Catonomyia spiculata* Hull, 1962 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 62 (syn.).
- Ref. – Artigas, Papavero & Serra, 1991: 62 (key to spp.).

- amapa** Artigas, Papavero & Serra, 1991: 63. Type-locality: Brazil, Amapá, Serra do Navio. Distr. – Brazil (Amapá). Type, MZUSP.
- anticus** (Wiedemann), 1828: 530 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Brazil (Rio de Janeiro, São Paulo). Refs. – Walker, 1855: 568 (*Atomosia*); Schiner, 1866: 693 (*Atomosia*); Williston, 1891: 79 (cat.; *Laphria*); Kertész, 1909: 163 (cat.; *Atomosia*); Hermann, 1912: 76 (*Dissmeryngodes (Laphria)*); Bromley, 1946: 108 (cat.; *Atomosia*); Carrera, 1958a: 145, 1960a: 150; Hull, 1962 (1): 416; Martin & Papavero, 1970: 54 (cat.); Artigas, Papavero & Serra, 1991: 63, figs. 11-17 (n. comb.); Castillo, Jerez & Artigas, 1994 (egg chorion). Type, SENCK [cf. Geller-Grimm, 1999: 206].
- iracema** Artigas, Papavero & Serra, 1991: 63. Type-locality: Brazil, São Paulo, Ribeirão Preto (Fazenda Iracema). Distr. – Brazil (São Paulo). Type, MZUSP.
- nigripes** (Macquart), 1838: 74 (1839: 190) (*Atomosia*). Type-locality: “Brazil”. Distr. – Colombia, Peru (Huánuco), Brazil (Pará). Refs. – Walker, 1855: 569 (*Atomosia*); Williston, 1891: 78 (cat.; *Atomosia*); Kertész, 1909: 164 (cat.; *Atomosia*); Bromley, 1934b: 338 (*Atractia*), 1946a: 108 (cat.; *Atomosia*); Hull, 1962 (1): 379, fig. 651 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*). Type, MNHNP.
- dispar** Walker, 1855: 570 (*Atomosia*). Type-locality: Brazil, Pará, Santarém. Refs. – Williston, 1891: 78 (cat.; *Atomosia*); Kertész, 1909: 163 (cat.; *Atomosia*); Hermann, 1912: 76 (*Dissmeryngodes (Atomosia)*); Bromley, 1946: 108 (cat.; *Atomosia*); Carrera, 1960a: 150; Hull, 1962 (1): 416, figs. 260, 659, 1330, 1339, 1571, 2125; Martin & Papavero, 1970: 54 (cat.); Lamas M., 1972: 312 (cat.); Artigas, Papavero & Serra, 1991: 63, figs. 1-10 (syn.). Type, BMNH.
- columbiana** Enderlein, 1914a: 242 (*Centrolaphria*). Type-locality: Colombia, Caldas, Victoria (Hacienda Pehlke), 900 m [cf. Papavero, 1973d: 299]. Refs. – Hull, 1962 (1): 416; Martin & Papavero, 1970: 54 (cat.; *Centrolaphria*); Artigas, Papavero & Serra, 1991: 63 (syn.). Type, IZPAN.
- spiculatus** (Hull), 1962 (1): 381, fig. 660 (*Catonomyia*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Refs. – Martin & Papavero, 1970: 48 (cat.; *Lycosimyia*); Artigas, Papavero & Serra, 1991: 63 (n. comb.). Type, ?lost [never sent to MZUSP].

Genus **Eumecosoma** Schiner, 1866

- Eumecosoma** Schiner, 1866: 662 (key), 673 (1868: 171, second erection of genus). Type-species, *Laphria pleuritica* Wiedemann, 1828 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 72, figs. 27-33.
- Josmayala** Kaletta, 1978: 58. Type-species, *Josmalaya tiarensis* Kaletta, 1978 (orig. des.) = *Eumecosoma tarsale* Curran, 1939. Refs. – Artigas, Papavero & Serra, 1991: 64 (syn.); Fisher & Hespenheide, 1992: 629 (note 2; syn.).
- ayala** Kaletta, 1974: 101. Type-locality: Venezuela, Aragua, Tiara (Campamento Rangel). Distr. – Venezuela (Aragua, Miranda). Ref. – Artigas, Papavero & Serra, 1991: 64 (cat.). Type, FAUCV.
- caeruleum** Scarbrough & Perez-Gelabert, 2006: 42, figs. 37-39 (as *caerulum*, error, as in p. 43 they say “Latin *caerulum* [sic], for blue”). Type-locality: Dominican Republic, La Vega, Parque Nacional Armando Bermúdez, km 1.3 along trail west of La Ciénaga, 900-1100 m. Distr. – Dominican Republic (Independencia, La Vega). Type, USNM.
- calverti** Hine, 1917: 206. Type-locality: Costa Rica, Juan Vilas. Distr. – Costa Rica. Refs. – Hull, 1962 (1): 380; Martin & Papavero, 1970: 49 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Type, OSU.
- carmina** Kaletta, 1974: 103. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. – Venezuela (Aragua). Ref. – Artigas, Papavero & Serra, 1991: 64 (cat.). Type, FAUCV.
- dicromum** Bigot, 1878: 236. Type-locality: “Brazil”. Refs. – Williston, 1891: 77 (cat.); Hull, 1962 (1): 380; Martin & Papavero, 1970: 49 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Type, OXF.
- dichroma** Kertész, 1909: 169 (cat.); Hermann, 1912: 72; Bromley, 1946: 110 (cat.), emend.
- hirsutum** Hermann, 1912: 73, fig. 27. Type-locality: Bolivia, Mapiri, Sarampiuni, 700 m and “Peru”. Distr. – Venezuela (Aragua, Guárico, Lara, Miranda), Peru, Bolivia. Refs. – Curran, 1930b: 8 (in key); Hull, 1962 (1): 380; Martin & Papavero, 1970: 49 (cat.); Kaletta, 1974: 100; Artigas, Papavero & Serra, 1991: 64 (cat.). Syntypes, SMTD [cf. Kallweit, 1992: 196].
- mellipes** (Wiedemann), 1828: 526 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1855: 538 (*Laphria*); Schiner, 1866: 693 (*Laphria*); Williston, 1891: 79 (cat.; *Laphria*); Kertész, 1909: 191 (cat.; *Laphria*); Bromley, 1946: 111 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Lamas M., 1972: 310 (cat.; n. comb.). Type, SENCK [cf. Geller-Grimm, 1999: 211].
- metallescens** Schiner, 1868: 171. Type-locality: “Venezuela”. Distr. – Venezuela (Táchira), Peru. Refs. – Williston, 1891: 77 (cat.); Kertész, 1909: 169 (cat.); Hermann, 1912: 73, fig. 26; Curran, 1930b: 8 (in key); Hull, 1962 (1): 380; Martin & Papavero, 1970: 50 (cat.); Lamas M., 1972: 310 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Syntypes, NMW.
- metalecens** Kaletta, 1974: 100, error.
- molle** Bromley, 1934b: 340. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica. Refs. – Hull, 1962 (1): 380; Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Type, AMNH.
- pleuriticum** (Wiedemann), 1828: 527 (misprinted as 257) (*Laphria*). Type-locality: “Brazil”. Refs. – Walker, 1855: 568 (*Atomosia*); Schiner, 1866: 693; Williston, 1891: 77 (cat.); Kertész, 1909: 169 (cat.); Heremann, 1912: 72, fig. 24; Curran, 1930b: 8 (in key; as *pleuritica*, error); Bromley, 1946: 110 (cat.); Hull, 1962 (1): 380, figs. 290, 681, 1290, 1299; Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Geller-Grimm, 1999: 213 (material in SENCK). Syntypes, NMW.
- shropshirei** Curran, 1930b: 8, fig. 2. Type-locality: Panama, Canal Zone, Barro Colorado I. Refs. – Hull, 1962 (1): 380; Shelly, 1987: 180 (natural history); Artigas, Papavero & Serra, 1991: 64 (cat.). Type, AMNH.
- staurophorum** Schiner, 1868: 172. Type-locality: “Venezuela”. Distr. – Venezuela, Peru (Huánuco). Refs. – Williston, 1891: 77 (cat.); Kertész, 1909: 169 (cat.); Hermann, 1912: 73, fig. 25; Curran, 1930b: 8 (in key); Hull, 1962 (1): 380; Martin & Papavero, 1970: 50 (cat.); Lamas M., 1972: 310 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Type, NMW.
- tarsale** Curran, 1930b: 8 (in key), 11 (as *tarsalis*, error). Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. – Panama, Venezuela (Aragua). Refs. – Hull, 1962 (1): 380; Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Type, AMNH.
- tiarensis** Kaletta, 1978: 60, pl. 2, figs. 1-4 (*Josmayala*). Type-locality: Venezuela, Aragua, Tiara (Campamento Rangel). Distr. – Refs. – Artigas, Papavero & Serra, 1991: 64 (cat.; n. comb.); Fisher & Hespenheide, 1992: 629 (note 2: syn.). Type, FAUCV.

Genus **Hodites** Hull, 1962

Hodites Hull, 1962 (1): 375. Type-species, *Hodites punctissima* Hull, 1962 (orig. des.).

punctissima Hull, 1962 (1): 376, figs. 286, 642, 1273, 1282. Type-locality: Brazil, Distrito Federal, Corumbá de Goiás (Fazenda Monjolinho). Distr. – Brasil (Distrito Federal). Refs. – Martin & Papavero, 1970: 48 (cat.); Artigas, Papavero & Serra, 1991: 64, figs. 73-74 (cat.). Type, MZUSP.

Genus **Hybozelodes** Hermann, 1912

Atractia Macquart of Hermann, 1912: 168, part, misident.

Hybozelodes Hermann, 1912: 197. Type-species, *Hybozelodes nigellus* Hermann, 1912 (orig. des.).

Lophoceraea Hermann, 1912: 193. Type-species, *Lophoceraea pennata* Hermann, 1912 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 64 (syn.).

acuticornis Carrera, 1945b: 171. Type-locality: Brazil, Rio de Janeiro, Mangaratiba. Distr. – Brazil (Rio de Janeiro, São Paulo). Refs. – Carrera, 1958a: 146, fig. 9; Hull, 1962 (1): 389; Martin & Papavero, 1970: 50 (cat.); Artigas, Papavero & Serra, 1991: 64, figs. 53–54 (cat.). Type, MZUSP.

albipes Hermann, 1912: 201. Type-locality: Peru, Cuzco, Mishagua, Urubamba river. Distr. – Peru (Cuzco). Refs. – Martin & Papavero, 1970: 50 (cat.); Lamas M., 1972: 310 (cat.); Artigas, Papavero & Serra, 1991: 64 (cat.). Type, SMTD [cf. Kallweit, 1992: 194].

clausicella (Carrera), 1960a: 150, figs. 3–4 (*Atractia*). Type-locality: Brazil, São Paulo, Cajuru. Refs. – Martin & Papavero, 1970: 48 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 64 (cat.; n. comb.). Type, MZUSP.

comatus (Hermann), 1912: 183, fig. 69 (*Atractia*). Type-locality: “Costa Rica”. Refs. – Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.). Syntypes, MUN.

conjugens (Hermann), 1912: 189, fig. 72 (*Atractia*). Type-locality: “Brazil”. Refs. – Bromley, 1946ag: 108 (cat.; *Atractia*); Hull, 1962 (1): 379 (*Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.). Type, NMW.

dispar (Hermann), 1912: 186, fig. 71 (*Atractia*). Type-locality: Peru (several localities not originally mentioned). Refs. – Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; gg *Atractia*); Lamas M., 1972: 310 (cat.); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.). Syntypes, ZSBS, SMTD [cf. Kallweit, 1992: 195].

fulvipes (Hermann), 1912: 185, fig. 70 (*Atractia*). Type-locality: Brazil, Santa Catarina, Blumenau. Refs. – Bromley, 1946: 108 (cat.; *Atractia*); Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.). Type, NMW.

lucidus (Hermann), 1912: 192, fig. 74 (*Atractia*). Type-locality: Brazil, Minas Gerais, Mar d’Hespanha (as Espírito Santo, in error). Brazil (Minas Gerais, Rio de Janeiro). Refs. – Bromley, 1946: 108 (cat.; *Atractia*); Carrera, 1960a: 150 (*Atractia*); Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.). Syntypes, MUN.

marginatus (Osten Sacken), 1887: 212 (*Atractia*). Type-locality: Nicaragua, Chontales. Refs. – Aldrich, 1905: 274 (cat.; *Atractia*); Kertész, 1909: 166 (cat.; *Atractia*); Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.); Shelly, 1984a: 120 (*Atractia*; prey selection). Type, BMNH.

?**minutus** (Wiedemann), 1828: 530 (*Laphria*). Type-locality: “Brazil”. Refs. – Walker, 1855: 569 (*Atomosia*); Schiner, 1866: 693 (?*Atractia*); Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 166 (cat.; *Atractia*); Bromley, 1946: 108 (cat.; *Atractia*); Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; ?*Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.). Type, BZM [no. 2268; cf. Schumann, 1973: 234].

nigellus Hermann, 1912: 199, fig. 76. Type-locality: Peru, Cuzco, Mishagua, Urubamba river. Distr. – Peru (Cuzco). Refs. – Hull, 1962 (1): 389, figs. 217, 653; Martin & Papavero, 1970: 50 (cat.); Lamas M., 1972: 310 (cat.); Artigas, Papavero & Serra, 1991: 65 (cat.). Syntypes, MUN, SMTD [cf. Kallweit, 1992: 196].

pennatus (Hermann), 1912: 196, figs. 75a–b, pl. 5, figs. 55–56 (*Lophoceraea*). Type-locality: Peru, Huánuco, mouth of Pachitea river, 150m; Bolivia, Mapiri, Sarapiqui. Distr. – Peru (Huánuco), Bolivia. Refs. – Hull, 1962 (1): 390, figs. 263, 678, 1326, 1336 (*Lophoceraea*); Martin & Papavero, 1970: 51 (cat.; *Lophoceraea*); Lamas M., 1972: 310 (cat.; *Lophoceraea*); Fisher, 1984: 33 (*Lophoceraea*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.). Syntypes, MUN, SMTD [cf. Kallweit, 1992: 197].

pictus (Hermann), 1912: 181, fig. 68 (*Atractia*). Type-locality: “Venezuela”. Refs. – Hull, 1962 (1): 379 (*Atractia*); Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.). Type, NMW.

platycerus Hermann, 1912: 202, fig. 77. Type-locality: Peru, Cuzco, Mishagua, Urubamba river. Distr. – Peru (Cuzco). Refs. – Hull, 1962 (1): 389; Martin & Papavero, 1970: 50 (cat.); Lamas M., 1972: 310 (cat.); Artigas, Papavero & Serra, 1991: 65 (cat.). Syntypes, SMTD [cf. Kallweit, 1992: 197].

Genus **Lamprozona** Loew, 1851

Laphria, subg. **Lamprozona** Loew, 1851: 18. Type-species, *Laphria (Lamprozona) auricincta* Loew, 1851 (mon.).

Automolina Hermann, 1912: 101. Type-species, *Automolina chilensis* Hermann, 1912 (orig. des.).

auricincta (Loew), 1851: 18 (*Laphria (Automolina)*). Type-locality: “Chile”. Distr. – Chile (Aconcagua, Arauco, Cautín, Linares, Maule, Santiago, Valdivia, Valparaíso). Refs. – Schiner, 1866: 706, 1868: 171; Williston, 1891: 79 (cat.); Kertész, 1909: 169 (cat.); Hermann, 1912: 99; Stuardo Ortiz, 1946: 83 (cat.); Hull, 1962 (1): 402; Artigas, 1970a: 243, figs. 203–205, 207–208, 440; Martin & Papavero, 1970: 53 (cat.); Artigas, 1971: 30, fig. 69–70; Artigas, Papavero & Serra, 1991: 65, figs. 25–26.

Papavero

atratus Philippi, 1865: 687 (*Dasypogon*). Type-locality: Chile, Valdivia. Refs. – Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 126 (cat.; *Dasypogon*); Bromley, 1932: 268; Stuardo Ortiz, 1946: 83 (cat.); Hull, 1962 (1): 402, figs. 224, 670, 1308, 1317, 2172. Type, MNHNS.

sericeus Philippi, 1865: 691 (*Dasypogon*). Type-locality: “Chile”. Refs. – Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 131 (cat.; *Dasypogon*); Stuardo Ortiz, 1946: 82 (cat.; *Dasypogon*); Hull, 1962 (1): 402 (*Dasypogon*). Type, MHNHS.

castaneipes Bigot, 1878: 234. Type-locality: “Chile”. Refs. – Williston, 1891: 79 (cat.); Kertész, 1909: 169 (cat.); Hull, 1962 (1): 402. Type, OXF.

cyanescens Bigot, 1878: 235 (*Cormansis*). Type-locality: “Chile”. Refs. – Williston, 1891: 78 (cat.; *Atomosia*); Kertész, 1909: 163 (cat.; *Atomosia*); Hermann, 1912: 100 (n. comb.); Brèthes, 1925: 105 (*Atomosia*); Hull, 1962 (1): 402 (as syn. of *chilensis* Hermann, 1912). Syntypes, OXF.

castanipes Bigot, 1879: ix, emend.

chilensis Brèthes, 1925: 105 (*Atomosia*; preocc. Hermann, 1912). Type-locality: “Chile”. Ref. – Stuardo Ortiz, 1946: 83 (cat.; *Atomosia*); Hull, 1962: 402.

chilensis (Hermann), 1912: 103 (*Automolina*). Type-locality: Chile, Concepción. Refs. – Bromley, 1932: 268 (*Automolina*); Stuardo Ortiz, 1946: 83 (cat.; *Automolina*); Hull, 1962 (1): 271, 409, figs. 262, 667, 1327, 1335 (*Automolina*); Artigas, 1970a: 246, figs. 206, 209-212, 437 (n. comb.); Martin & Papavero, 1970: 53 (cat.); Artigas, Papavero & Serra, 1991: 65 (cat.). Type, ZSBS.

Genus **Lycosimyia** Hull, 1958

Lycosimyia Hull, 1958a: 102. Type-species, *Lycosimyia carrerae* Hull, 1958 (orig. des.).

Ref. – Artigas, Papavero & Serra, 1991: 77, figs. 57-65.

carrerae Hull, 1958a: 103 (as *carrerae*, in error). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Distr. – Brazil (Rio de Janeiro). Refs. – Hull, 1962 (1): 375, figs. 285, 1291, 1300; Martin & Papavero, 1970: 48 (cat.); Artigas, Papavero & Serra, 1991: 65 (cat.). Type, MZUSP.

fluviatilis (Carrera), 1960a: 152, fig. 5 (*Atractia*). Type-locality: Brazil, Pará, Óbidos. Distr. – Brazil (Pará). Refs. – Martin & Papavero, 1970: 49 (cat.; *Atractia*); Artigas, Papavero & Serra, 1991: 65 (cat.). Type, MNRJ.

Genus **Oidardis** Hermann, 1912

Oidardis Hermann, 1912: 76. Type-species, *Oidardis gibbosa* Hermann, 1912 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 72, figs. 34-38.

Othoniomyia Hermann, 1912: 119. Type-species, *Othoniomyia triangularis* Hermann, 1912 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 65 (syn.).

Othoniomyia, subg. *Sphagolestes* Hull, 1962 (1): 391. Type-species, *Othoniomyia (Sphagolestes) nigrum* Hull, 1962 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 65 (syn.).

Voluptarius Kaletta, 1978: 48. Type-species, *Voluptarius curupaoensis* Kaletta, 1978 (orig. des.). Ref. – Artigas, Papavero & Serra, 1991: 65 (syn.).

Iucundus Kaletta, 1978: 52. Type-species, *Iucundus aveledoi* Kaletta, 1978 (orig. des.).

aenescens Hermann, 1912: 80. Type-locality: Peru, Pasco, Unini, Ucayali river. Distr. – Peru (Pasco). Refs. – Hull, 1962 (1): 409; Martin & Papavero, 1970: 53 (cat.); Lamas M., 1972: 311 (cat.); Artigas, Papavero & Serra, 1991: 65 (cat.). Syntypes, SMTD [cf. Kallweit, 1992: 194].

aveledoi (Kaletta), 1978: 54, pl. 1, figs. 4, 6-9 (*Iucundus*). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. – Venezuela (Aragua). Ref. – Artigas, Papavero & Serra, 1991: 65, figs. 39-40 (cat.; n. comb.). Type, FAUCV.

curupaoensis (Kaletta), 1978: 50, pl. 1, figs. 1-3, 5 (*Voluptarius*). Type-locality: Venezuela, Miranda, Curupao. Distr. – Venezuela (Miranda). Ref. – Artigas, Papavero & Serra, 1991: 65, figs. 41-43 (cat.; n. comb.). Type, FAUCV.

gibba (Curran), 1930b: 8 (in key), 10 (*Eumecosoma*). Type-locality: Panama, Canal Zone, Barro Colorado I. Refs. – Hull, 1962 (1): 380 (*Eumecosoma*); Martin & Papavero, 1970: 49 (cat.; *Eumecosoma*); Shelly, 1986b: 124 (foraging success; variation with attack distance and angle), 1987: 180 (natural history); Artigas, Papavero & Serra, 1991: 65, figs. 44-45 (cat.; n. comb.). Type, AMNH.

gibbosa Hermann, 1912: 78. Type-locality: Peru, Cuzco, Umuhuankiali, 500 m, Puerto Bermúdez, 700 m; Bolivia, Chimate, 650 m. Distr. – Peru (Pasco), Bolivia. Refs. – Hull, 1962 (1): 409, figs. 218, 677, 1260, 1269; Martin & Papavero, 1970: 53 (cat.); Lamas M., 1972: 311 (cat.); Fisher, 1984: 31; Artigas, Papavero & Serra, 1991: 65 (cat.). Syntypes, MUN,

SMTD [cf. Kallweit, 1992: 195].
nigra (Hull), 1962 (1): 391 (*Othoniomyia (Sphagolestes)*). Type-locality: Brazil, São Paulo. Refs. – Martin & Papavero, 1970: 51 (cat.; *Othoniomyia*); Artigas, Papavero & Serra, 1991: 65 (cat.; n. comb.).
triangularis (Hermann), 1912: 120, figs. 47–48 (*Othoniomyia*). Type-locality: Brazil, Santa Catarina, Blumenau. Distr. – Brazil (Rio de Janeiro, São Paulo, Paraná, Santa Catarina). Refs. – Bromley, 1946: 108 (cat.); Carrera, 1946a: 98 (*Othoniomyia*), 1958a: 145 (*Othoniomyia*); Hull, 1962 (1): 391, figs. 249, 654, 1352, 1361, 1669, 1814; Martin & Papavero, 1970: 51 (cat.; *Othoniomyia*); Artigas, Papavero & Serra, 1991: 65, figs. 46–47 (cat.; n. comb.). Syntypes, MIL, NMW, MUN.

Genus **Strombocodia** Hermann, 1912

Strombocodia Hermann, 1912: 165. Type-species, *Strombocodia elegans* Hermann, 1912 (orig. des.).

elegans Hermann, 1912: 167, fig. 62, pl. 4, fig. 51, pl. 5, fig. 52. Type-locality: Peru, Cuzco, Umuhanckiali, Urubamba river, 500 m. Distr. – Peru (Cuzco). Refs. – Hull, 1962 (1): 375, fig. 2511; Martin & Papavero, 1970: 48 (cat.); Lamas M., 1972: 310 (cat.); Artigas, Papavero & Serra, 1991: 65 (cat.). Type, MUN.

Unrecognized genus and species of Atomosiini

Genus **Rhatimomyia** Lynch Arribálzaga, 1882

Rhatimomyia Lynch Arribálzaga, 1882: 135. Type-species, *Rhatimomyia nitidula* Lynch Arribálzaga, 1882 (mon.).
Rhatitomyia Williston, 1891: 78, emend.

nitidula Lynch Arribálzaga, 1882: 137. Type-locality: Argentina, Buenos Aires, Chacabuco. Distr. – Argentina (Buenos Aires). Refs. – Williston, 1891: 78 (cat.; *Rhatitomyia*); Kertész, 1909: 167 (cat.); Hull, 1962 (1): 400, figs. 230, 652, 1316, 1325; Martin & Papavero, 1970: 53 (cat.). Type, lost.

Tribe **Dasytrichini** Lynch Arribálzaga, 1882

Dasythricaria Lynch Arribálzaga, 1882: 245. Type-genus, *Dasythrix* Loew, 1851. Ref. – Dikow, 2003: 465.
Erecmoneminae Hermann, 1912: 21, part.
Dasytrichini Artigas, Papavero & Pimentel, 1988: 212, 237, as new, error.

Genus **Cryptomerynx** Enderlein, 1914

Cryptomerynx Enderlein, 1914a: 254. Type-species, *Cryptomerynx laphriicornis* Enderlein, 1914 (orig. des.).
Alipiolaphria Carrera, 1951: 116. Type-species, *Alipiolaphria mirandai* Carrera, 1951 (orig. des.). Ref. – Artigas, Papavero & Pimentel, 1988: 237 (syn.).

laphriicornis Enderlein, 1914a: 254, figs. 1–2. Type-locality: “Colombia [Pehlke]”. Refs. – Hull, 1962 (1): 344, figs. 2515, 2522; Artigas, Papavero & Pimentel, 1988: 215, figs. 3–4. Type, IZPAN.
mirandai (Carrera), 1951: 116, figs. 4–6, 11–14 (*Alipiolaphria*). Type-locality: Brazil, Paraná, Curitiba. Distr. – Brazil (Paraná). Refs. – Hull, 1962 (1): 333; Artigas, Papavero & Pimentel, 1988: 237 (n. comb.). Type, MZUSP.

Genus **Dasythrix** Loew, 1851

Laphria, subg. **Dasythrix** Loew, 1851: 21. Type-species, *Laphria (Dasythrix) inornata* Loew, 1851 (orig. des.).

inornata (Loew), 1851: 21 (*Laphria (Dasythrix)*). Type-locality: “South America” (Uruguay, Montevideo (Sello) [cf. Papavero, 1971e: 56-60, 69-80, 99-102] in TP label). Refs. – Williston, 1891: 81 (cat.); Kertész, 1909: 207 (cat.); Hull, 1962 (1): 343 (*Nusa*); Martin & Papavero, 1970: 41 (cat.; ?*Nusa*); Artigas, Papavero & Pimentel, 1988: 237. Type, BZM [no. 432; cf. Schumann, 1973: 121].

leucophaea Lynch Arribálzaga, 1880: 55. Type-locality: Argentina, Buenos Aires. Refs. – Wulp, 1882b: 195; Lynch Arribálzaga, 1883a: 145 (cat.); Williston, 1891: 81 (cat.); Engel, 1930: 468; Hull, 1962 (1): 343 (*Nusa*); Martin & Papavero, 1970: 41 (cat.; ?*Nusa*); Artigas, Papavero & Pimentel, 1988: 237. Type, MACN.

Genus **Smeryngolaphria** Hermann, 1912

Smeryngolaphria Hermann, 1912: 226. Type-species, *Smeryngolaphria pictipennis* Hermann, 1912 (mon.) = *maculipennis* (Macquart, 1846).

Panamasilus Curran, 1930: 20. Type-species, *Panamasilus xylota* Curran, 1930 (orig. Des.) = *numitor* (Osten Sacken, 1887).

Ref. – Artigas, Papavero & Pimentel, 1988: 239 (key to spp.).

gorayebi Artigas, Papavero & Pimentel, 1988: 240, figs. 7, 14, 17-20, 21-22. Type-locality: Brazil, Pará, Monte Dourado. Distr. – Brazil (Pará, Amapá). Type, MPEG.

gurupi Artigas, Papavero & Pimentel, 1988: 241, figs. 9, 16. Type-locality: Brazil, Pará, rio Gurupi, Aldeia Gurupi. Distr. – Brazil (Pará). Type, MZUSP.

maculipennis (Macquart), 1846: 201 (1846: 73), pl. 7, fig. 14 (*Laphria*). Type-locality: “Guyana”. Distr. – Guyana, Brazil (Amazonas, Pará, Amapá), Bolivia. Refs. – Walker, 1855: 539 (*Laphria*); Williston, 1891: 81 (cat.; *Andrenosoma*); Kertész, 1909: 200 (cat.; *Nusa*); Hermann, 1912: 227; Curran, 1942: 57; Carrera, 1951: 119, 1960a: 152; Hull, 1962 (1): 333; Martin & Papavero, 1970: 39 (cat.); Artigas, Papavero & Pimentel, 1988: 242, figs. 12, 23-26. Syntypes, OXF. *pictipennis* Hermann, 1912: 227. Type-locality: Bolivia, Mapiri, San Ernesto, 300 m. Ref. – Curran, 1942: 57 (syn.). Syntypes, TP SMTD [cf. Kallweit, 1992: 197].

aurata Enderlein, 1914a: 252 (*Laphria*). Type-locality: Brazil, Pará, Óbidos (as Amazonas, in error). Ref. – Bromley, 1946: 111 (cat.). Type, IZPAN.

melanura (Wiedemann), 1828: 508 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Brazil (Espírito Santo, Rio de Janeiro). Refs. – Macquart, 1838: 539 (*Laphria*); Walker, 1855: 539 (*Laphria*); Schiner, 1866: 691 (*Laphria*); Williston, 1891: 79 (cat.; *Laphria*); Kertész, 1909: 191 (cat.; *Laphria*); Hermann, 1912: 226; Bromley, 1946: 111 (cat.); Carrera, 1951: 113, figs. 7-9; Hull, 1962 (1): 333; Martin & Papavero, 1970: 39 (cat.); Artigas, Papavero & Pimentel, 1988: 243, figs. 11, 27-30. Type, SENCK [cf. Geller-Grimm, 1999: 211; face eaten away by dermestids].

numitor (Osten Sacken), 1887: 185 (*Laphria*). Type-locality: Nicaragua, Chontales. Distr. – Nicaragua, Panama. Refs. – Aldrich, 1905: 272 (cat.; *Laphria*); Kertész, 1909: 192 (cat.; *Laphria*); Martin & Papavero, 1970: 39 (cat.); Artigas, Papavero & Pimentel, 1988: 243, figs. 5, 10, 31-34. Type, BMNH.

xylota Curran, 1930b: 20, fig. 3 (*Panamasilus*). Type-locality: Panama, Canal Zone, Barro Colorado I. Refs. – Carrera, 1951: 120; Hull, 1962 (1): 333. Type, AMNH.

panamensis Curran, 1942: 57 (*Panamasilus*; ?*lapsus calami* for *xylota*).

seabrai Carrera, 1960a: 153, fig. 6. Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (Tijuca). Distr. – Brazil (Rio de Janeiro). Refs. – Martin & Papavero, 1970: 39 (cat.); Artigas, Papavero & Pimentel, 1988: 243, figs. 6, 13, 35-38. Type, MNRJ.

taperignae Artigas, Papavero & Pimentel, 1988: 243, figs. 8, 15, 39-42. Type-locality: Brazil, Pará, Santarém (Fazenda Taperinha). Distr. – Brazil (Pará). Type, MZUSP.

Tribe **Laphriini** Macquart, 1838

Erecmoninae Hermann, 1912: 21, part.

Genus **Ichneumolaphria** Carrera, 1951

Ichneumolaphria Carrera, 1951: 114. Type-species, *Ichneumolaphria zikani* Carrera, 1912 (orig. des.).

Ref. – Geller-Grimm, 1997 (key to spp.).

fascipennis (Hermann), 1912: 216 (*Rhopalogaster*). Type-locality: Peru, Cuzco, Mishagua, Urubamba river, 350 m. Distr. – Peru (Cuzco). Refs. – Hull, 1962 (1): 339; Martin & Papavero, 1970: 39 (cat.); Lamas M., 1972: 308 (cat.). Syntypes, ZSBS, SMTD [cf. Kallweit, 1992: 195].

schichti Geller-Grimm, 1997: 457, figs. 1a-f. Type-locality: Brazil, Amazonas, rio do Equador. Distr. – Brazil (Amazonas). Type, ZSBS.

zikani Carrera, 1951: 115, figs. 1, 3. Type-locality: Brazil, Minas Gerais, Virgínia, 900 m. Distr. – Brazil (Minas Gerais). Refs. – Hull, 1962 (1): 333; Martin & Papavero, 1970: 39 (cat.).

Genus **Joartigasia** Martínez & Martínez, 1975

Pholidotus Hermann, 1912: 329 (preocc. Brisson, 1762). Type-species, *Pholidotus rubriventris* Hermann, 1912 (sub. des., Martin & Papavero, 1970: 41).

Joartigasia Martínez & Martínez, 1975: 149 *nom. nov.* for *Pholidotus* Hermann, 1912. Type-species, *Pholidotus rubriventris* Hermann, 1912 (aut.).

anceps (Hermann), 1912: 234 (*Pholidotus*). Type-locality: Peru, Huánuco, mouth of Pachitea river. Distr. – Peru (Huánuco). Refs. – Hull, 1962 (1): 336, figs. 618, 1305, 2154; Martin & Papavero, 1970: 41 (cat.; *Pholidotus*); Lamas M., 1972: 308 (cat.; *Pholidotus*). Type, SMTD [cf. Kallweit, 1992: 194].

rubriventris (Hermann), 1912: 229 (*Pholidotus*). Type-locality: “Peru (male), Surinam (female)”. Distr. – Surinam, Guyana, Peru. Refs. – Bromley, 1934b: 346 (*Pholidotus*); Hull, 1962 (1): 336, figs. 268, 616, 1225, 1234, 2148, 2155; Martin & Papavero, 1970: 41 (cat.; *Pholidotus*); Lamas M., 1972: 308 (cat.; *Pholidotus*). Syntypes, ZSBS (male), NMW (female).

ruficaudis (Hermann), 1912: 232 (*Pholidotus*). Type-locality: Peru, Huánuco, mouth of Pachitea river, 150 m. Distr. – Peru (Huánuco). Refs. – Hull, 1962 (1): 336; Martin & Papavero, 1970: 41 (cat.; *Pholidotus*); Lamas M., 1972: 308 (cat.; *Pholidotus*). Type, SMTD [cf. Kallweit, 1992: 197].

Genus **Lampria** Macquart, 1838

Lampria Macquart, 1838: 60 (1839: 178). Type-species, *Laphria clavipes* Fabricius, 1805 (sub. des., Coquillett, 1910: 557). *Lampria*, subg. *Brychomyia* Hull, 1962 (1): 328. Type-species, *Laphria rubriventris* Macquart, 1834 (orig. des.).

aurifex Osten Sacken, 1887: 187. Type-locality: Mexico, Veracruz, Orizaba; Costa Rica, Caché. Distr. – Mexico (Veracruz, Nayarit, San Luis Potosí, Tabasco), Guatemala, Costa Rica. Refs. – Williston, 1901: 318; Aldrich, 1905: 270 (cat.); Kertész, 1909: 204 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 39 (cat.); Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.

bitincta Walker, 1860: 280. Type-locality: “Brazil, Amazon”. Refs. – Williston, 1891: 80 (cat.); Kertész, 1909: 205 (cat.); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 327.

bicincta Martin & Papavero, 1970: 40 (cat.), error.

ciliipes Walker, 1857: 128. Type-locality: “Brazil, Amazon”. Refs. – Williston, 1891: 80 (cat.); Kertész, 1909: 205 (cat.); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.).

circumdata Bellardi, 1861: 115 (15), pl. 1, fig. 17. Type-locality: “Mexico”. Distr. – Mexico. Refs. – Aldrich, 1905: 270 (cat.); Kertész, 1909: 205 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.); Papavero & Ibáñez Bernal, 2001: 105 (hist. Mexican dipterol.). Type, TORO.

clavipes (Fabricius), 1805: 162 (*Laphria*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Mexico (San Luis Potosí), Costa Rica, Panama, Peru (Huánuco, Loreto, Pasco), Brazil (Pará, Espírito Santo). Refs. – Wiedemann, 1821: 237 (*Laphria*), 1828: 513 (*Laphria*); Macquart, 1838: 61, 1848: 22, pl. 2, fig. 7; Rondani, 1848: 91; Walker, 1849: 376, 1855: 510; Schiner, 1866: 691, 1868: 174; Wulp, 1882b: 194; Osten Sacken, 1887: 186; Williston, 1891: 80 (cat.); Aldrich, 1905: 270 (cat.); Kertész, 1909: 205 (cat.); Hermann, 1912: 220, 222; Bromley, 1934: 345; Carrera, 1945a: 160 (preys); Bromley, 1946: 111 (cat.); Carrera & d’Andretta, 1953: 63; Carrera, 1960a: 152; Hull, 1962 (1): 327, figs. 226, 612, 1237, 1248; Carrera & Machado-Allison, 1963: 239; Martin & Papavero, 1970: 40 (cat.); Lamas M., 1972: 308 (cat.). Syntypes, COPN, KIEL [cf. Zimsen, 1964: 469].

corallogaster (Bigot), 1878: 226 (*Laphria*). Type-locality: “North America”. Distr. – U. S. A., Mexico. Refs. – Williston, 1883: 33 (*Laphria*); Aldrich, 1905: 270 (cat.); Kertész, 1909: 205 (cat.); Martin & Wilcox, 1965: 388 (cat.); Martin & Papavero, 1970: 40 (cat.). Type, OXF.

dives (Wiedemann), 1828: 524 (*Laphria*). Type-locality: Brazil, Bahia [Gomez; cf. Papavero, 1971e: 48]. Distr. – Peru, Bolivia, Brazil (Bahia). Refs. – Walker, 1855: 534 (*Laphria*); Schiner, 1866: 692; Williston, 1891: 80 (cat.); Kertész, 1909: 205

- (cat.); Hermann, 1912: 224; Bromley, 1946: 111 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.); Fisher, 1984: 29. Syntypes, BZM [no. 402; cf. Schumann, 1973: 105].
- fulgida** Schiner, 1868: 174. Type-locality: Brazil, São Paulo, Mogi Mirim, Oriçanga (as “Insenga”, error for “Orissanga”). Distr. – Brazil (São Paulo). Refs. – Williston, 1891: 80 (cat.); Kertész, 1909: 205 (cat.); Hermann, 1912: 223; Bromley, 1946: 111 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.). Syntypes, NMW.
- homopoda** (Bellardi), 1862: 217 (20), pl. 3, fig. 16 (*Laphria*). Type-locality: Mexico, Veracruz, Tuxpano near Orizaba. Distr. – Mexico (Veracruz). Refs. – Aldrich, 1905: 272 (cat.; *Laphria*); Kertész, 1909: 187 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 40 (cat.); Papavero & Ibáñez Bernal, 2001: 110 (hist. Mexican dipterol). Type, TORO.
- ichneumon** (Osten Sacken), 1887: 185, pl. 3, fig. 6 (*Laphria*). Type-locality: Guatemala, Purula, 4000 ft. Distr. – Guatemala. Refs. – Aldrich, 1905: 272 (cat.; *Laphria*); Kertész, 1909: 187 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Hull, 1962 (1): 323; Martin & Papavero, 1970: 40 (cat.). Type, BMNH.
- macquartii** (Perty), 1833: 181, pl. 36, fig. 3 (*Laphria*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (as “Sebastianopolis”). Distr. – Brazil (Rio de Janeiro). Refs. – Walker, 1849: 376 (*Laphria*), 1855: 511 (*Laphria*); Williston, 1891: 80 (cat.); Kertész, 1909: 206 (cat.); Bromley, 1934b: 346 (believes *macquarti* is the female of *clavipes* Fabricius, 1805), 1946a: 111 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.). Type, ?ZSBS.
- mexicana** Macquart, 1847: 1847: 53 (1847: 37). Type-locality: “Mexico”. Distr. – Mexico (San Luis Potosí, Veracruz), Costa Rica, Panama, Venezuela. Refs. – Walker, 1855: 510; Bellardi, 1861: 13; Wulp, 1882b: 105; Osten Sacken, 1887: 188; Aldrich, 1905: 270 (cat.); Kertész, 1909: 206 (cat.); Carrera & d’Andretta, 1950: 161, figs. 9, 27; Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.); Papavero & Ibáñez Bernal, 2001: 86 (hist. Mexican dipterol.). Syntypes, MNHNP, OXF.
- parvula** Bigot, 1878: 237. Type-locality: “Brazil”. Refs. – Williston, 1891: 80 (cat.); Kertész, 1909: 206 (cat.); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.). Type, OXF.
- pusilla** (Macquart), 1838: 68 (1839: 184) (*Laphria*). Type-locality: Brazil, Rio Grande do Sul. Refs. – Walker, 1855: 538 (*Laphria*); Williston, 1891: 80 (cat.); Kertész, 1909: 206 (cat.); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.). Type, MNHNP.
- rubriventris** (Macquart), 1834: 284 (*Laphria*). Type-locality: U. S. A., Pennsylvania. Distr. – U. S. A. (Florida, Georgia, Pennsylvania, Texas), Mexico (Coahuila). Refs. – Walker, 1855: 527 (*Laphria*); Bigot, 1878: 217 (*Pogonosoma*); Osten Sacken, 1878: 76, 233 (cat.); Williston, 1883: 32; Aldrich, 1905: 270 (cat.); Kertész, 1909: 206 (cat.); Fattig, 1945: 20; Bromley, 1950: 233; Hull, 1962 (1): 327 [*Lampria* (*Brychomyia*)] Hull, 1962 (1): 327; Martin & Wilcox, 1965: 388 (cat.).
- scapularis** Bigot, 1878: 237. Type-locality: Brazil, Pará. Distr. – Brazil (Pará, São Paulo, Santa Catarina). Refs. – Williston, 1891: 80 (cat.); Kertész, 1909: 206 (cat.); Bromley, 1946: 111 (cat.); Carrera, 1958a: 146, 1960: 153; Carrera & Vulcano, 1961: 70 (prey); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.). Type, OXF.
- spinipes** (Fabricius), 1805: 162 (*Laphria*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Guyana. Refs. – Wiedemann, 1821: 240 (*Laphria*), 1828: 525 (*Laphria*); Walker, 1849: 377 (*Laphria*), 1855: 534 (*Laphria*); Schiner, 1866: 693; Osten Sacken, 1887: 187; Williston, 1891: 80 (cat.); Aldrich, 1905: 270 (cat.); Kertész, 1909: 206 (cat.); Bromley, 1934b: 345; Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.). Type, COPN [cf. Zimsen, 1964: 469].
- affinis** Fabricius, 1805: 163 (*Laphria*). Type-locality: “South America”. Type, COPN [cf. Zimsen, 1974: 469].
- splendens** (Macquart), 1834: 286 (*Laphria*). Type-locality: “Surinam”. Refs. – Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.). Type, lost.
- tolmides** (Walker), 1849: 377 (*Laphria*). Type-locality: “?Brazil”. Refs. – Walker, 1855: 511; Williston, 1891: 80 (cat.); Kertész, 1909: 206 (cat.); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 327; Martin & Papavero, 1970: 40 (cat.).

Genus *Laphria* Meigen, 1803

Lapria Meigen, 1800: 25. Type-species, *Asilus gibbosus* Linnaeus, 1758 (sub. des., Coquillett, 1910: 557). Suppressed by I. C. Z. N., 1963: 359.

Laphria Meigen, 1803: 270. Type-species, *Asilus gibbosus* Linnaeus, 1758 (sub. des.; Latreille, 1810: 443).

Bombomima Enderlein, 1914a: 258. Type-species, *Laphria fulvithorax* Fabricius, 1805 (orig. des.) = *thoracica* (Fabricius, 1805).

componens Walker, 1860: 281. Type-locality: “Mexico”. Distr. – Mexico. Refs. – Aldrich, 1905: 272 (cat.); Kertész, 1909: 179 (cat.); Hull, 1962 (1): 323; Martin & Papavero, 1970: 38 (cat.); Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.).

marginalis Williston, 1901: 318. Type-locality: Mexico, Veracruz, Attoyac. Distr. – Mexico (Veracruz). Refs. – Aldrich, 1905: 272 (cat.); Kertész, 1909: 189 (cat.); Hull, 1962 (1): 323; Martin & Papavero, 1970: 38 (cat.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol). Type, BMNH.

melanogaster Wiedemann, 1821: 236. Type-locality: “Mexico and Savannah”. Distr. – U. S. A. (Georgia, Texas), Mexico,

- ?Colombia. Refs. – Wiedemann, 1828: 507; Macquart, 1846: 75; Walker, 1855: 528; Schiner, 1866: 691; Aldrich, 1905: 272 (cat.); Kertész, 1909: 191 (cat.); Enderlein, 1914a: 252; Hull, 1962 (1): 325 [*Laphria (Bombomima)*]; Martin & Wilcox, 1965: 387 (cat.); Martin & Papavero, 1970: 38 (cat.); Papavero & Ibáñez Bernal, 2001: 64 (hist. Mexican dipterol.).
- thoracica** Fabricius, 1805: 158. Type-locality: “North America”. Distr. – Canada, U. S. A., ?Jamaica. Refs. – Fattig, 1945: 19 (*Bombomima (Dasyllis)*); Farr, 1965: 16 (*Bombomima*); Martin & Wilcox, 1965: 391 (cat.); Hull, 1962 (1): 325 [*Laphria (Bombomima)*]; Martin & Papavero, 1970: 38 (cat.; *Laphria (Bombomima)*). Type, KIEL (lost; only name label remainig; cf. Zimšen, 1964: 469).
- fulvithorax* Fabricius, 1805: 373 (unjustified *nom. nov.* for *thoracica* Fabricius, 1805).
- alcanor* Walker, 1849: 383. Type-locality: U. S. A., Massachusetts. Refs. – Gowdey, 1926: 77.
- trux** McAtee, 1919: 158. Type-locality: U. S. A., California. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Hull, 1962 (1): 322; Martin & Wilcox, 1965: 391 (cat.).
- audax* McAtee, 1919: 158 (as *trux* var.). Type-locality: U. S. A., California.
- willistoniiana** Enderlein, 1914a: 252 (*nom. nov.* for *coerulea* Williston, 1901). Type-locality: Mexico, Veracruz, Atoyac. Distr. – Mexico. Ref. – Hull, 1962 (1): 323.
- coerulea* Williston, 1901: 317 (preocc. Boisduval, 1835). Type-locality: Mexico, Veracruz, Atoyac. Refs. – Aldrich, 1905: 272 (cat.); Kertész, 1909: 179 (cat.); Martin & Papavero, 1970: 38 (cat.); Papavero & Ibáñez Bernal, 2003: 176 (hist. Mexican dipterol.). Type, BMNH.

Genus **Phellopteron** Hull, 1962

Phellopteron Hull, 1962 (1): 341. Type-species, *Phellopteron farri* Hull, 1962 (orig. des.).

- farri** Hull, 1962 (1): 341, figs. 256, 628, 1219, 1228, 2150. Type-locality: Jamaica, Windsor Estate, Trelawny, 12 miles south of Falmouth. Distr. – Jamaica. Refs. – Farr, 1965: 15, figs. 9-10; Martin & Papavero, 1970: 41 (cat.). Type, SIKJ.
- mrazi** Hradský, 1968: 165, figs. 1-2. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. – Brazil (Rio de Janeiro). Ref. – Martin & Papavero, 1970: 41 (cat.). Type, MZUSP.

Genus **Rhopalogaster** Macquart, 1834

Rhopalogaster Macquart, 1834: 279. Type-species, *Laphria longicornis* Wiedemann, 1828 (mon.).

- albidus** Scarbrough & Perez-Gelabert, 2006: 45, fig. 121. Type-locality: Dominican Republic, La Vega, Parque Nacional Armando Bermúdez, km 1.3 along trail west of La Ciénaga, 900-1100 m. Distr. – Dominican Republic (La Vega). Type, USNM.
- araujoi** Carrera, 1952a: 211. Type-locality: Brazil, Goiás, Goiânia (Campinas). Distr. – Brazil (Goiás). Refs. – Hull, 1962 (1): 339; Martin & Papavero, 1970: 40 (cat.). Type, MZUSP.
- aurifer** Hermann, 1912: 211. Type-locality: unknown. Refs. – Hull, 1962 (1): 339; Martin & Papavero, 1970: 40 (cat.). Type, NMW.
- bella** Bromley, 1929a: 283, pl. 1, fig. 5. Type-locality: Cuba, Camaguey. Distr. – Cuba. Refs. – Hull, 1962 (1): 339; Martin & Papavero, 1970: 41 (cat.); Alayo Dalmau & García Ávila, 1983: 46 (cat.). Type, USNM.
- lineata** Hermann, 1912: 212. Type-locality: Brazil, Rio Grande do Sul; Minas Gerais, Mar d’Espanha; Bolivia, Yungas de Coroico, 1800 m. Distr. – Bolivia, Brazil (Minas Gerais, São Paulo, Santa Catarina, Rio Grande do Sul). Refs. – Bromley, 1946: 112 (cat.); Carrera, 1953a: 271; Hull, 1962 (1): 339; Martin & Papavero, 1970: 41 (cat.). Syntypes, NMW, SMTD [cf. Kallweit, 1992: 196].
- longicornis** (Wiedemann), 1828: 519 (*Laphria*). Type-locality: “Brazil”. Refs. – Macquart, 1834: 279, 1838: 72; Walker, 1855: 565; Schiner, 1866: 692; Williston, 1891: 81 (cat.); Kertész, 1909: 208 cat.; Hermann, 1912: 211; Bromley, 1946: 112 (cat.); Hull, 1962 (1): 339, figs. 619, 1221, 1230, 2095, 2098, 2143; Martin & Papavero, 1970: 41 (cat.). Type, NMW.
- micronyx** Tomasovic, 2002: 86, fig. 1. Type-locality: French Guiana, route de Kaw, Camp Patawa, 4o31'N, 52o05'W. Distr. – French Guiana. Type, FUSAGembloix.
- niphardis** Hermann, 1912: 213. Type-locality: Bolivia, La Paz, Mapiri, Sarampiuní, 700 m; San Carlos, 800 m. Ref. – Hull, 1962 (1): 339. Syntypes, NMW, SMTD [cf. Kallweit, 1992: 196].

Tribe **Neophoneini** Artigas, Papavero & Pimentel, 1988

Erecmoneminae Hermann, 1912: 21, part.

Neophoneini Artigas, Papavero & Pimentel, 1988: 214, 244. Type-genus, *Neophoneus* Williston, 1889.

Genus **Neophoneus** Williston, 1889

Phoneus Macquart, 1838: 79 (1839: 195) (preocc. Kaup, 1820). Type-species, *Phoneus servillei* Macquart, 1838 (mon.).
Neophoneus Williston, 1889: 255 (*nom. nov.* for *Phoneus* Macquart, 1838). Type-species, *Phoneus servillei* Macquart, 1838 (aut.).

amandus (Walker), 1849: 373 (*Laphria*). Type-locality: “Brazil”. Refs. – Walker, 1855: 538 (*Laphria*); Williston, 1891: 80 (cat.; *Andrenosoma*); Kertész, 1909: 197 (*Nusa*); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 354; Martin & Papavero, 1970: 45 (cat.). Type, BMNH (abdomen lacking).

amanda Aldrich, 1905: 272 (cat.; *Laphria*), error.

servillei (Macquart), 1838: 79 (1839: 195), pl. 7, figs. 5, 5a (*Phoneus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 573 (*Phoneus*); Williston, 1891: 82 (cat.); Kertész, 1909: 171 (cat.); Bromley, 1946: 111 (cat.); Hull, 1962 (1): 354, figs. 650, 1236, 1245; Martin & Papavero, 1970: 45 (cat.). Syntypes, lost.

?*mustela* Hermann, 1912: 269, fig. 86. Type-locality: “Surinam (female); Brazil, Rio Grande do Sul (male). Refs. – Bromley, 1946: 111 (cat.); Hull, 1962 (1): 354, fig. 273; Artigas, Papavero & Pimentel, 1988: 225, figs. 43-44. Syntypes, MUN.

Unrecognized species of Laphriinae (asterisk (*) indicates that the type is lost)

albibitalis Macquart, 1847: 54 (1847: 38) (*Laphria*). Type-locality: “Brazil”. Refs. – Walker, 1855: 535 (*Laphria*); Williston, 1891: 79 (cat.; *Laphria*); Kertész, 1909: 176 (*Laphria*); Bromley, 1946: 111 (cat.; *Laphria*); Hull, 1962 (1): 323; Martin & Papavero, 1970: 37 (cat.; *Laphria*). Type, OXF.

albibibia Macquart, 1848: 185 (1848: 25), pl. 2, fig. 6 (*Laphria*). Type-locality: “Brazil” and “Rio Negro”.

***appendiculata** Macquart, 1846: 204 (1846: 76), pl. 8, fig. 1 (*Atomosia*). Type-locality: French Guiana, Cayenne. Refs. – Walker, 1855: 569 (*Atomosia*); Williston, 1891: 77 (cat.; *Atomosia*); Kertész, 1909: 163 (cat.; *Atomosia*); Hull, 1962 (1): 373 (*Atomosia*); Martin & Papavero, 1970: 46 (cat.; *Atomosia*).

***bimaculata** Walker, 1855: 534, 550 (*Laphria*). Type-locality: Brazil, “Amazon”. Refs. – Williston, 1891: 79 (cat.; *Laphria*); Bromley, 1946: 111 (cat.; *Laphria*); Martin & Papavero, 1970: 37 (cat.; *Laphria*). [The two specimens under this name in the BMNH do not agree with the original description and are therefore not the types].

dentipes Fabricius, 1805: 162 (*Laphria*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Refs. – Kertész, 1909: 180 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*). Syntypes, COPN, KIEL (in bad condition) [cf. Zimsen, 1964: 469].

dichroa Wiedemann, 1828: 526 (*Laphria*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1855: 693 (*Laphria*); Schiner, 1866: 693 (*Laphria*); Williston, 1891: 79 (cat.; *Laphria*); Bromley, 1946: 111 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 180 (cat.; *Laphria*); Lamas M., 1972: 308 (cat.; *Laphria* (*Laphria*)). Syntypes, IZPAN, SMTD [cf. Kallweit, 1992: 195].

***glauca** Enderlein, 1914a: 251 (*Laphria*). Type-locality: Ecuador, Coca, Archidona and Canelos; Peru, Junín, Chanchamayo; Peru, Loreto, Pebas (as Brazil, Amazonas, in error). Refs. – Bromley, 1946: 111 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 180 (cat.; *Laphria*); Lamas M., 1972: 308 (cat.; *Laphria* (*Laphria*)). Syntypes, IZPAN, SMTD [cf. Kallweit, 1992: 195].

***limbata** Macquart, 1834: 287 (*Laphria*). Type-locality: French Guiana, Cayenne. Refs. – Walker, 1855: 539 (*Laphria*); Williston, 1891: 78 (cat.; *Atomosia*); Kertész, 1909: 164 (cat.; *Atomosia*); Hull, 1962 (1): 373 (*Atomosia*); Martin & Papavero, 1970: 46 (cat.; *Atomosia*).

***modesta** Philippi, 1865: 685, fig. (*Laphria*). Type-locality: Chile, Santiago. Refs. – Schiner, 1866: 706 (*Laphria*); Lynch Arribálzaga, 1882: 144 (*Atomosia*); Williston, 1891: 78 (cat.; *Atomosia*); Kertész, 1909: 164 (cat.; *Atomosia*); Stuardo Ortiz, 1946: 83 (cat.; *Atomosia*); Hull, 1962 (1): 373 (*Atomosia*); Artigas, 1970a: 373; Martin & Papavero, 1970: 47 (cat.; *Atomosia*). Type, lost.

***pilipes** Macquart, 1834: 282 (*Laphria*). Type-locality: “Brazil”. Refs. – Walker, 1855: 537 (*Laphria*); Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 192 (cat.; *Laphria*); Bromley, 1946: 110 (cat.; *Andrenosoma*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*).

***proxima** Walker, 1855: 537 (*Laphria*). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 192 (cat.; *Laphria*); Bromley, 1946: 111 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*).

pusilla Wiedemann, 1828: 532 (*Laphria*). Type-locality: “Brazil”. Refs. – Walker, 1855: 568 (*Atomosia*); Schiner, 1866: 568

- (*Laphria*); Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 192 (cat.; *Laphria*); Bromley, 1946:a 111 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*). Type, SENCK [cf. Geller-Grimm, 1999: 213]. (Unrecognised Atomosiini).
- rubescens** Bigot, 1878: 225 (*Laphria*). Type-locality: "Colombia". Refs. – Osten Sacken, 1891: 119 (*Laphria*); Kertész, 1909: 192 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*).
- *sericans** Walker, 1860: 282 (*Atomosia*). Type-locality: "Mexico". Refs. – Aldrich, 1905: 269 (cat.; *Atomosia*); Kertész, 1909: 155 (cat.; *Atomosia*); Hull, 1962 (1): 373 (*Atomosia*); Martin & Papavero, 1970: 47 (cat.; *Atomosia*); Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.).
- transatlantica** Schiner, 1868: 173 (*Laphria*). Type-locality: "Venezuela". Refs. – Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 195 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*). Syntypes, NMW.
- *triligata** Walker, 1861: 281 (*Laphria*). Type-locality: "Mexico". Refs. – Aldrich, 1905: 273 (cat.; *Laphria*); Kertész, 1909: 195 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*); Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.). Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.).
- *venezuelensis** Macquart, 1846: 202 (1846: 74), pl. 7, fig. 15 (*Laphria*). Type-locality: "Venezuela". Refs. – Walker, 1855: 536 (*Laphria*); Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 196 (*Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*).
- *violacea** Macquart, 1846: 202 (1846: 74) (*Laphria*). Type-locality: "Colombia". Refs. – Walker, 1855: 534 (*Laphria*); Williston, 1891: 80 (cat.; *Laphria*); Kertész, 1909: 196 (cat.; *Laphria*); Hull, 1962 (1): 323 (*Laphria*); Martin & Papavero, 1970: 38 (cat.; *Laphria*).

Subfamily Laphystiinae Hendel, 1936

- Prytaninae Hermann, 1920: 167, 168.
 Laphystiini Hendel, 1936: 1923. Type-genus, *Laphystia* Loew, 1847.
 Hoplistomerinae Enderlein, 1936: 91, 94. Type-genus, *Hoplistomerus* Macquart, 1838.
 Laphystiinae Papavero, 1973a: 232, 243.

Genus *Apoxyria* Schiner, 1866

- Apoxyria** Schiner, 1866: 674. Type-species, *Apoxyria apicata* Schiner, 1866 (orig. des.).

- americana** Carrera, 1955a: 109, figs. 3-6. Type-locality: Brazil, Distrito Federal, Corumbá de Goiás (Fazenda Monjolinho). Distr. – Brazil (Distrito Federal). Refs. – Hull, 1962 (1): 82 (*Helolaphycitis*); Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 8, figs. 7-8, 36, 64-68, 69-70. Type, MZUSP.
- apicata** Schiner, 1866: 682. Type-locality: not stated (Brazil, São Paulo, Ipanema, in TP label). Distr. – Brazil (São Paulo). Refs. – Hermann, 1905: 31; Kertész, 1909: 159 (cat.); Hull, 1962 (1): 103, figs. 35, 454, 859, 868, 1563, 1769; Martin & Papavero, 1970: 11 (cat.); Artigas, Papavero & Costa, 1997: 8. Type, NMW.

Genus *Asicya* Lynch Arribálzaga, 1880

- Asicya** Lynch Arribálzaga, 1880: 224. Type-species, *Asicya fasciata* Lynch Arribálzaga, 1880 (mon.). Ref. – Artigas, Papavero & Costa, 1997: figs. 27-28, 46, 50B, 121-124, 125-126.

- fasciata** Lynch Arribálzaga, 1880: 227. Type-locality: Argentina, Buenos Aires, Baradero. Distr. – Argentina (Buenos Aires). Refs. – Lynch Arribálzaga, 1883a: 145 (cat.); Williston, 1891: 81 (cat.); Hermann, 1908: 154, fig. 3; Kertész, 1909: 158 (cat.; *Laphystia*); Hull, 1962 (1): 76 (*Laphystia*); Martin & Papavero, 1970: 9 (cat.; *Laphystia*); Artigas, Papavero & Costa, 1997: 8. Type, MACN (in very poor condition).

Genus *Chrysotriclis* Artigas, Papavero & Costa, 1997

- Chrysotriclis** Artigas, Papavero & Costa, 1997: 6. Type-species, *Chrysotriclis willinkinorum* Artigas, Papavero & Costa, 1997 (orig. des.).

willinkinorum Artigas, Papavero & Costa, 1997: 6, figs. 31-32, 48, 133-136, 137-138. Type-locality: Argentina, Misiones, Loreto. Distr. – Argentina (Formosa, Misiones). Type, MZUSP.

Genus **Cochleariocera** Artigas, Papavero & Costa, 1997

Cochleariocera Artigas, Papavero & Costa, 1997: 7. Type-species, *Cochleariocera neusae* Artigas, Papavero & Costa, 1997 (orig. des.).

neusae Artigas, Papavero & Costa, 1997: 8, figs. 19-20, 42, 1104-10500-103,. Type-locality: Brazil. Goiás, lago Rico, rio Araguaia. Distr. – Brazil (Distrito Federal, Goiás). Type, MZUSP.

Genus **Cymbipyga** Artigas, Papavero & Costa, 1997

Cymbipyga Artigas, Papavero & Costa, 1997: 8. Type-species, *Apoxyria cymafera* Artigas, 1983 (orig. des.).

cymafera (Artigas), 1983: 28, figs. 1-16 (*Apoxyria*; as *cymafer*, error). Type-locality: Chile, Santiago, Colina, El Portezuelo. Distr. – Chile (Santiago). Ref. – Artigas, Papavero & Costa, 1997: 9, figs. 11-12, 38, 38A, 75-78, 79-80. Type, DZUC.

Genus **Gymnotriclis** Artigas, Papavero & Costa, 1997

Gymnotriclis Artigas, Papavero & Costa, 1997: 9. Type-species, *Gymnotriclis coscaronorum* Artigas, Papavero & Costa, 1997 (orig. des.).

coscaronorum Artigas, Papavero & Costa, 1997: 10, figs. 29-30, 47, 127-130, 131-132. Type-locality: Venezuela, Santa Ana, Tacarigua. Distr. – Panama, Venezuela. Type, MZUSP.

Genus **Helolaphyctis** Hermann, 1920

Helolaphyctis Hermann, 1920: 193. Type-species, *Helolaphyctis modesta* Hermann, 1920 (orig. des.).

Refs. – Hull, 1962 (1): figs. 39, 536, 861, 870, 1676, 1878, 1929; Artigas, Papavero & Costa, 1997: figs. 5-6, 35, 57-61, 62-63.

chrysorhea Hull, 1958f: 97. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (Santa Catarina). Refs. – Martin & Papavero, 1970: 9 (cat.); Artigas, Papavero & Costa, 1997: 10. Type, Hull Coll.

modesta Hermann, 1920: 194, *nomen nudum*. Ref. – Hull, 1962 (1): 82.

nitida Hull, 1958d: 162 (as *nitidus*, error). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (Santa Catarina). Refs. – Martin & Papavero, 1970: 9 (cat.); Artigas, Papavero & Costa, 1997: 10. Type, BMNH.

Genus **Hexameritia** Speiser, 1920

Eutrichodes Hermann, 1920: 190 (preocc. Warren, 1891). Type-species, *Dasypogon micans* Philippi, 1865 (orig. des.).

Hexameritia Speiser, 1920: 447 (*nom. nov.* for *Eutrichodes* Hermann, 1920). Type-species, *Dasypogon micans* Philippi, 1865 (aut.).

micans (Philippi), 1865: 690, fig. (*Dasypogon*). Type-locality: Chile, Santiago. Neotype-locality: Chile, Linares, Cordillera Parral. Distr. – Chile (Aconcagua, Concepción, Coquimbo, Curicó, Linares, Ñuble, O’Higgins, Santiago, Valparaíso). Refs. – Bigot, 1878: 220 (*Dasypeucus*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 90 (*Dasypeucus*); Hermann, 1920: 190 (*Eutrichodes*); Stuardo Ortiz, 1946: 82 (cat.; *Dasypeucus*); Hull, 1962 (1): 83; Artigas, 1970a: 121, figs. 34-41, 474; Martin & Papavero, 1970: 9 (cat.); Artigas, 1971: 16, figs. 26-27; Artigas, Papavero & Costa, 1997: 10, figs. 3-4,

34, 53-56. Type, lost. NT DZUC [des. by Artigas, 1970a: 122].
splendens Philippi, 1865: 690 (*Dasypogon*). Type-locality: Chile, Choapa, Illapel. Refs. – Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 132 (cat.; *Dasypogon*); Hull, 1962 (1): 83, fig. 425; Martin & Papavero, 1970: 9 (syn.). Type, lost.
tricolor Schiner, 1868: 162 (*Triclis*). Type-locality: “Chile”. Refs. – Bigot, 1878: 221 (*Triclis*); Williston, 1891: 72 (cat.; *Triclis*); Kertész, 1909: 98 (cat.; *Triclis*); Hull, 1962 (1): 83, figs. 32, 420, 862, 871; Martin & Papavero, 1970: 9 (syn.). Type, NMW.

Genus **Laphygmolestes** Hull, 1962

Laphygmolestes Hull, 1962 (1): 76. Type-species, *Laphygmolestes flavipes* Hull, 1962 (orig. des.).

flavipes Hull, 1962 (1): 78. Type-locality: Brazil, Mato Grosso do Sul, Corumbá. Distr. – Brazil (Mato Grosso do Sul). Refs. – Martin & Papavero, 1970: 9 (cat.); Artigas, Papavero & Costa, 1997: 11, figs. 9-10, 37, 71-74. Type, not in COR.

Genus **Laphystia** Loew, 1847

Laphystia Loew, 1847: 538. Type-species, *Laphystia sabulicola* Loew, 1847 (mon.).

opaca Coquillett, 1904b: 180. Type-locality: U. S. A., Texas. Distr. – U. S. A. (Louisiana, Texas), Mexico (Tamaulipas, Veracruz). Refs. – Back, 1909: 229; Curran, 1930b: 12 (in key), 14; Bromley, 1935c: 3; Wilcox, 1960: 341, fig. 2; Hull, 1962 (1): 76; Martin & Wilcox, 1965: 376 (cat.); Martin & Papavero, 1970: 9 (cat.). Type, USNM.

sillersi Hull, 1963: 202. Type-locality: U. S. A., Arizona. Distr. – U. S. A. (Arizona), Mexico (Sonora). Ref. – Martin & Wilcox, 1965: 377 (cat.).

sonora Wilcox, 1960: 343, fig. 1. Type-locality: Mexico, Sonora, La Cholla. Distr. – Mexico (Sonora). Refs. – Martin & Papavero, 1970: 9 (cat.); Artigas, Papavero & Costa, 1997: 11. Type, CU.

Genus **Macahyba** Carrera, 1947

Macahyba Carrera, 1947c: 203. Type-species, *Macahyba nordestina* Carrera, 1947 (orig. des.).

nordestina Carrera, 1947c: 205, figs. 1-9. Type-locality: Brazil, Ceará, Icó. Distr. – Brazil (Ceará). Refs. – Hull, 1962 (1): 90, figs. 445, 876, 885; Martin & Papavero, 1970: 9 (cat.); Artigas, Papavero & Costa, 1997: 11, figs. 15-16, 40, 50A, 88-91, 92-93. Type, MZUSP.

Genus **Martinomyia** Özdi̇kmen, 2006

Martinia Hull, 1962 (1): 91 (preocc M'Coy, 1844). Type-species, *Martinia moloch* Hull, 1962 (orig. des.).

Martinomyia Özdi̇kmen, 2006: 93, nom. nov. for *Martinia* Hull, 1962. Type-species, *Martinia moloch* Hull, 1962 (aut.).

Ref. – Artigas, Papavero & Costa, 1997: figs. 17-18, 41, 94, 97, 98-99.

moloch (Hull), 1962 (1): 92 (*Martinia*). Type-locality: Bolivia, Camamina [?]. Distr. – Bolivia. Refs. – Martin & Papavero, 1970: 10 (cat.; *Martinia*); Artigas, Papavero & Costa, 1997: 11 (*Martinia*); Özdi̇kmen, 2006: 94 (n. comb.). Type, USNM.

scalaris (Hermann), 1908: 157, fig. 4 (*Laphystia*). Type-locality: Peru, mouth of Pachitea river, 150 m. Distr. – Peru, Argentina. Refs. – Curran, 1930b: 11 (in key), 18 (*Laphystia*); Engel, 1930: 468; Martin & Papavero, 1970: 10 (cat.; *Macahyba*); Lamas M., 1972: 305 (cat.; *Laphystia*); Fisher, 1984: 30 (*Martinia*); Artigas, Papavero & Costa, 1997: 11 (*Martinia*); Özdi̇kmen, 2006: 94 (n. comb.). Syntypes, MUN, SMTD [cf. Kallweit, 1992: 197].

schnusei (Hermann), 1908: 159, figs. 5-6, pl. 4, fig. 2 (*Laphystia*). Type-locality: Peru, Ucayali river, 300 m. Distr. – Peru. Refs. – Hull, 1962 (1): 76 (*Laphystia*); Martin & Papavero, 1970: 9 (cat.; *Laphystia*); Lamas M., 1972: 306 (cat.; *Macahyba*); Fisher, 1984: 30 (*Martinia*); Artigas, Papavero & Costa, 1997: 11 (?*Macahyba*). Syntypes, MUN, TP SMTD [cf.

Kallweit, 1992: 198]. **N. comb.**

Genus **Perasis** Hermann, 1905

Perasis Hermann, 1905: 37. Type-species, *Perasis sareptana* Hermann, 1905 (orig. des.). Ref. – Artigas, Papavero & Costa, 1997: figs. 25-26, 45, 119-120.

argentifacies (Williston), 1901: 310, pl. 5, fig. 22 (*Triclis*). Type-locality: Mexico, Guerrero, Chilpancingo. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 257 (cat.; *Triclis*); Kertész, 1909: 98 (cat.; *Triclis*); Carrera & d'Andretta, 1950: 159, figs. 8, 24 (*Triclis*); Hull, 1962 (1): 86 (*Triclis*); Martin & Wilcox, 1965: 382 (cat.); Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 11; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Syntypes, BMNH.

Genus **Protometer** Artigas, Papavero & Costa, 1997

Protometer Artigas, Papavero & Costa, 1997: 11. Type-species, *Protometer evae* Artigas, Papavero & Costa, 1997 (orig. des.).

bokermanni Artigas, Papavero & Costa, 1997: 121, fig. 110. Type-locality: Brazil, Mato Grosso, São Domingos. Distr. – Brazil (Mato Grosso). Type, MZUSP.

evae Artigas, Papavero & Costa, 1997: 12, figs. 21-22, 43, 106-109, 110, 111-112. Type-locality: Brazil, Amazonas, Paraná da Eva. Distr. – Brazil (Amazonas). Type, MZUSP.

Genus **Psilocurus** Loew, 1874

Psilocurus Loew, 1874: 373. Type-species, *Psilocurus nudiusculus* Loew, 1874 (mon.).

Orthoneuromyia Williston, 1893a: 67. Type-species, *Orthoneuromyia modesta* Williston, 1893 (mon.).

Ref. – Artigas, Papavero & Costa, 1997: figs. 13-14, 39, 81-85, 86-87.

camposi Curran, 1931b: 8. Type-locality: Ecuador, Guayas, Guayaquil. Distr. – Ecuador. Refs. – Hull, 1962 (1): 91; Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 53. Type, AMNH.

caudatus Williston, 1901: 308, pl. 5, fig. 20. Type-locality: Mexico, Guerrero: Chilpancingo, Amula and Tepetlapa. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 257 (cat.); Kertész, 1909: 86 (cat.); Curran, 1930b: 8 (in key); Carrera & d'Andretta, 1950: 160, figs. 6, 20; Hull, 1962 (1): 91; Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 53; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Syntypes, BMNH.

puellus Bromley, 1934a: 92. Type-locality: U. S. A., Texas, Bexar County. Distr. – U. S. A. (Arizona, New Mexico, Texas), Mexico (Sonora). Refs. – Hull, 1962 (1): 91; Martin & Wilcox, 1965: 382 (cat.); Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 53.

reinhardi Bromley, 1951: 16. Type-locality: U. S. A., Texas, Bexar County. Distr. – U. S. A. (Texas). Refs. – Hull, 1962 (1): 91; Martin & Wilcox, 1965: 382 (cat.). Type, AMNH.

Genus **Triclioscelis** Roeder, 1900

Triclioscelis Roeder, 1900: 337. Type-species, *Triclioscelis burmeisteri* Roeder, 1900 (sub. des., Hermann, 1920: 183).

Ref. – Curran, 1934a: 2 (key to spp.).

burmeisteri Roeder, 1900: 337. Type-locality: Argentina, Tucumán. Distr. – Argentina (Tucumán). Refs. – Kertész, 1909: 99 (cat.); Engel, 1930: 468; Curran, 1934a: 2 (in key); Hull, 1962 (1): 99, figs. 1670, 1689; Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 53.

femorata Roeder, 1900: 339. Type-locality: Argentina, Santa Fe. Distr. – Paraguay, Argentina (Santa Fe). Refs. – Kertész, 1909: 99 (cat.); Engel, 1930: 468; Curran, 1934a: 2 (in key); Carrera, 1947a: 44; Hull, 1962 (1): 99; Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 53, figs. 23-24, 44, 49, 50, 113-116, 117-118.

- perfecta** Curran, 1934a: 2. Type-locality: Ecuador, Guayas, Golfo de Guayaquil, Puerto Grande, Isla Puná. Distr. – Ecuador. Refs. – Hull, 1962 (1): 99, figs. 41, 427, 893, 902, 1914, 1976; Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 53. Type, AMNH.
- salti** (Curran), 1931b: 11 (in key), 18 (*Laphystia*). Type-locality: Colombia, Magdalena, Ciénaga. Distr. – Colombia. Refs. – Curran, 1934a: 2; Hull, 1962 (1): 76 (*Laphystia*); Martin & Papavero, 1970: 10 (cat.); Artigas, Papavero & Costa, 1997: 53. Type, AMNH.

Genus **Zabrops** Hull, 1958

Zabrops Hull, 1958g: 90. Type-species, *Triclis tagax* Williston, 1883 (orig. des.).

Ref. – Fisher, 1977 (rev.).

- janiceae** Fisher, 1977: 211, figs. 55-57, 65, 72-73. Type-locality: Mexico, Baja California, 20 miles west of Bahía de los Ángeles. Distr. – Mexico (Baja California). Ref. – Artigas, Papavero & Costa, 1997: 53. Type, CAS.
- tagax** (Williston), 1883: 9, pl. 1, figs. 6, 6a (*Triclis*). Type-locality: U. S. A., California, Kern County. Distr. – U. S. A. (California). Refs. – Williston, 1886: 37 (*Triclis*); Back, 1909: 231, pl. 9, fig. 3 (*Triclis*); Hull, 1958g: 90, figs. 58, 444, 883, 892 (n. comb.), 1962 (1): 100, figs. 58, 444, 883, 892; Martin & Wilcox, 1965: 386 (cat.); Fisher, 1977: 200, figs. 33-42, 59, 66, 73; Artigas, Papavero & Costa, 1997: 53, figs. 1-2, 33, 51-52. Type, KU.
- ssp. **argutus** Fisher, 1977: 204, figs. 46, 60, 67, 73. Type-locality: Mexico, Baja California, 7 miles northwest of El Rosario. Distr. – Mexico (Baja California). Ref. – Artigas, Papavero & Costa, 1997: 53. Type, CAS.
- thologaster** Fisher, 1977: 210, figs. 52-54, 64, 71, 73. Type-locality: Mexico, Baja California, 0.5 mile northeast of El Molino, Bahía San Quintín. Distr. – Mexico (Baja California). Ref. – Artigas, Papavero & Costa, 1997: 53. Type, CAS.

Unrecognized species of Laphystiinae

- columbina** Schiner, 1868: 175 (*Laphystia*). Type-locality: “Colombia”. Refs. – Hull, 1962 (1): 76 (*Laphystia*); Martin & Papavero, 1970: 9 (cat.; *Laphystia*). Type, NMW.
- robusta** Hermann, 1908: 161, fig. 7 (*Laphystia*). Type-locality: Peru, Cuzco, Rosalina, Urubamba river, 700 m. Refs. – Hull, 1962 (1): 76 (*Laphystia*); Martin & Papavero, 1970: 9 (cat.; *Laphystia*); Lamas M., 1972: 305 (*Laphystia*). Syntypes, MUN, SMTD [cf. Kallweit, 1992: 197].

Subfamily Leptogastrinae Schiner, 1862

- Leptogastrinae Schiner, 1862b: liv. Type-genus, *Leptogaster*
Leptogasterinae G. H. Hardy, 1926: 306.
Leptogastridae Martin, 1968a: 70.

Genus **Apachekolos** Martin, 1957

Apachekolos Martin, 1957a: 352. Type-species, *Leptogaster scapularis* Bigot, 1878 (orig. des.).

- clavipes** (Johnson), 1894: 273 (*Leptogaster*; *nom. nov.* for *longipes* Johnson, 1894). Type-locality: “Jamaica”. Distr. – Bahama Is., Jamaica, Dominican Republic. Refs. – Aldrich, 1905: 254 (cat.; *Leptogaster*); Kertész, 1909: 50 (cat.; *Leptogaster*); Johnson, 1919: 430 (*Leptogaster*); Gowdey, 1926: 77 (cat.; *Leptogaster*); Hull, 1962 (1): 299 (*Leptogaster*); Farr, 1963: 18, fig. 11 (n. comb.); Martin, 1968e: 5 (cat.); Scarbrough, Perez-Gelabert & Page, 2005: 32, figs. 1, 7, 13-18, 62. Type, MCZ.
- longipes* Johnson, 1894: 273 (*Leptogaster*; preocc. Walker, 1859). Type-locality: “Jamaica”. Type, MCZ.
- acus** Martin, 1957b: 1, 6 (in key), figs. 1-2. Type-locality: Bahamas, Berry Islands, Little Harbor Cay. Ref. – Farr, 1963: 18 (syn.). Type, AMNH.
- crinita** Martin, 1957a: 354, fig. 1. Type-locality: U. S. A., New Mexico, 8 miles east of Pietown, near the Arizona eastern border. Distr. – U. S. A. (California, New Mexico, Arizona), Mexico (Baja California, Sonora, San Luis Potosí). Refs. – Hull, 1962

- (1): 301 (*Leptogaster (Apachekulos)*); Martin & Wilcox, 1965: 361 (cat.); Martin, 1968e: 5 (cat.). Type, AMNH.
- flaventis** Scarbrough & Perez-Gelabert in Scarbrough, Perez-Gelabert & Page, 2005: 32 (in key), 33, figs. 8, 19-21, 62. Type-locality: Dominican Republic, Azua, Loma El Cucuricho, 814 m. Distr. – Dominican Republic. Type, USNM.
- invasus** Scarbrough & Perez-Gelabert in Scarbrough, Perez-Gelabert & Page, 2005: 32 (in key), 34, figs. 10, 25-27, 62. Type-locality: Dominican Republic, Montecristi, road 186, road north to Playa Buen Hombre. Distr. – Dominican Republic. Type, USNM.
- magnus** Scarbrough & Perez-Gelabert in Scarbrough, Perez-Gelabert & Page, 2005: 32 (in key), 35, figs. 10, 25-27, 62. Type-locality: Dominican Republic, Montecristi, road 186, road to Playa Buen Hombre, 132 m. Distr. – Dominican Republic. Type, USNM.
- scapularis** (Bigot), 1878: 444 (*Leptogaster*). Type-locality: U. S. A., California. Distr. – U. S. A. (California, Texas), Mexico (Baja California Norte, Baja California Sur, Tamaulipas). Refs. – Williston, 1883: 33 (*Leptogaster*); Back, 1909: 169 (*Leptogaster*); Martin, 1957a: 353, fig. 2 (n. comb.); Hull, 1962 (1): 301 (*Leptogaster (Apachekulos)*); Martin & Wilcox, 1965: 361 (cat.). Type, ?.
- volubilis** Scarbrough & Perez-Gelabert in Scarbrough, Perez-Gelabert & Page, 2005: 32 (in key), 37, figs. 11, 28-33, 62. Type-locality: Dominican Republic, Peravia, Matadero, start of Hills, 450 m. Distr. – Dominican Republic. Type, USNM.
- vultus** Scarbrough & Perez-Gelabert in Scarbrough, Perez-Gelabert & Page, 2005: 31 (in key), 38, figs. 34-39, 62. Type-locality: Dominican Republic, Plata, road 221, 8 km south of Bombita, Puerto. Distr. – Dominican Republic. Type, USNM.

Genus **Beameromyia** Martin, 1957

- Beameromyia** Martin, 1957a: 355. Type-species, *Leptogaster pictipes* Loew, 1862 (orig. des.).
- chrysops** Martin, 1957a: 356, fig. 12. Type-locality: U. S. A., Florida, Royal Palm Park. Distr. – U. S. A. (Florida), Jamaica. Refs. – Refs. – Hull, 1962 (1): 301 (*Leptogaster (Beameromyia)*); Farr, 1963: 16, fig. 8; Martin & Wilcox, 1965: 361 (cat.); Martin, 1968e: 5 (cat.). Type, KU.
- cubensis** (Bigot), 1857a: 332 (*Gonypes*). Type-locality: “Cuba”. Distr. – Cuba, Puerto Rico (Mona Island). Refs. – Osten Sacken, 1874: 176 (*Leptogaster*); Roeder, 1885: 340 (*Leptogaster*); Aldrich, 1905: 254 (cat.; *Leptogaster*); Kertész, 1909: 51 (cat.; *Leptogaster*); Curran, 1928b: 22; Bromley, 1929a: 274 (*Leptogaster*); Hull, 1962 (1): 299 (*Leptogaster*); Martin, 1968e: 5 (cat.); Alayo Dalmau & García Ávila, 1983: 50; Scarbrough, 1997b: 239, figs. 7-12. Type, MNHNP.
- dicrana** Scarbrough, 1997b: 241, figs. 13-18. Type-locality: Dominican Republic, Pedernales, ca. 35 km north of Cabo Rojo, Las Abejas, 1250 m. Distr. – Dominican Republic. Ref. – Scarbrough, Perez-Gelabert & Page, 2005: 31 (in key), 39, figs. 2, 40-42, 62. Type, USNM.
- dorsalis** (Williston), 1901: 301 (*Leptogaster*). Type-locality: Mexico, Guerrero, Xucumanatlán. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 254 (cat.; *Leptogaster*); Kertész, 1909: 52 (cat.; *Leptogaster*); Hull, 1962 (1): 299 (*Leptogaster*); Martin, 1968e: 5 (cat.); Papavero & Ibáñez Bernal, 2003: 174 (hist. Mexican dipterol). Type, BMNH.
- graminicola** Farr, 1963: 16, figs. 9-10. Type-locality: Jamaica, Kingston, Palisadoes. Distr. – Jamaica. Refs. – Martin, 1968e: 5 (cat.); Scarbrough, 1997b: 241. Type, USNM.
- incisuralis** Scarbrough & Page in Scarbrough, Perez-Gelabert & Page, 2005: 31 (in key), 41, figs. 3, 43-50, 63. Type-locality: Dominican Republic, Pedernales, Upper Las Abejas, 38 km north-northwest of Cabo Rojo, 1350 m. Distr. – Dominican Republic. Type, CMNH.
- insulara** Martin, 1957b: 3, 6 (in key), fig. 3. Type-locality: Bahama Islands, Turks and Caicos Island, West Caicos Island. Distr. – Bahama Islands. Ref. – Scarbrough, 1997b: 242, figs. 19-20. Type, AMNH.
- insularia* Martin, 1968e: 5 (cat.), error.
- macula** Martin, 1957: 360, fig. 15. Type-locality: U. S. A., Arizona, Ruby . Distr. – U. S. A. (Arizona), Mexico (Coahuila). Refs. – Linsley & Cazier, 1963: 117 (biol.); Martin & Wilcox, 1965: 362 (cat.). Type, KU.
- melana** Scarbrough, 1997b: 238, figs. 1-6. Type-locality: Dominican Republic, Jarabacoa. Distr. – Dominican Republic, Haiti. Ref. – Scarbrough, Perez-Gelabert & Page, 2005: 30 (in key), 43, fig. 63. Type, USNM.
- punicea** Martin, 1957a: 362, fig. 13. Type-locality: U. S. A., Arizona, Santa Rita Mountains, Madera Canyon. Distr. – U. S. A. (Arizona), Mexico (Chiapas). Refs. – Martin & Wilcox, 1965: 362 (cat.); Martin, 1968e: 5 (cat.). Type, KU.
- quaterna** Scarbrough, 1997b: 243, figs. 21-29. Type-locality: Puerto Rico, Mona Island, Los Pinos. Distr. – Puerto Rico. Type, USNM.
- virginensis** Scarbrough, 1997: 244, figs. 30-37. Type-locality: Virgin Islands, Guana Island Distr, – Virgin Islands. Type, USNM.

Genus **Eurhabdus** Aldrich, 1923

- Eurhabdus** Aldrich, 1923: 2. Type-species, *Eurhabdus zephyreus* Aldrich, 1923 (orig. des.).

- jamaicensis** Farr, 1973: 324, figs. 1-4. Type-locality: Jamaica, Portland, Hardwar Gap, Green Hills. Distr. – Jamaica. Type, USNM.
- longissimus** Tomasovic, 2002: 88, fig. 2. Type-locality: French Guiana, Montagne de Kaw, Relais Patawa. Distr. – French Guiana. Type, FUSAGembloix.
- sororius** Scarbrough & Perez-Gelabert in Scarbrough, Perez-Gelabert & Page, 2005: 30 (in key), 44, figs. 4, 51-52, 63. Type-locality: Dominican Republic, Pedernales, 26 km north of Cabo Rojo, 730 m. Distr. – Dominican Republic. Type, CMNH.
- sp.** Farr, 1963: 29, fig. 12. Jamaica.
- zephyreus** Aldrich, 1923: 3, figs. 1-3. Type-locality: Costa Rica, San Mateo, Higuito. Distr. – Costa Rica (San Mateo). Refs. – Hull, 1962 (1): 317, figs. 587, 1196, 1205, 2037; Martin, 1968e: 6 (cat.). Type, USNM.

Genus **Leptogaster** Meigen, 1803

Leptogaster Meigen, 1803: 269. Type-species, *Asilus tipuloides* Linnaeus, 1758 (mon.; misident.) = *Leptogaster cylindrica* (De Geer, 1776).

Gonypes Latreille, 1805: 309. Type-species, *Asilus tipuloides* Linnaeus, 1758 (mon.; misident.) = *Leptogaster cylindrica* (De Geer, 1776).

One species known as fossil: *Leptogaster prior* Melander (1947: 44) (Oligocene; U. S. A., Colorado).

aegra Martin, 1957a: 381, fig. 53. Type-locality: U. S. A., North Carolina, Raleigh. Distr. – U. S. A. (Missouri, North Carolina, South Carolina, Florida), Mexico (San Luis Potosí). Refs. – Hull, 1962 (1): 298; Martin & Wilcox, 1965: 362 (cat.). Type, NCDA.

annulipes Walker, 1855: 309. Type-locality: Brazil, Pará [i. e., Pará, Belém]. Distr. – Brazil (Pará). Refs. – Williston, 1891: 68 (cat.); Kertész, 1909: 50 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 2 (cat.).

apicalis Enderlein, 1914b: 153. Type-locality: Ecuador, Tungurahua: Hacienda Santa Inés, 1244 m (1°25'S, 78°12'W) [cf. Brown, 1941: 846]. Distr. – Ecuador. Refs. – Hull, 1962 (1): 299; Martin, 1968e: 2 (cat.). Type, IZPAN.

arenicola James, 1937: 13. Type-locality: U. S. A., Colorado. Distr. – U. S. A. (Colorado, Nebraska), Mexico (Coahuila). Ref. – Baker, 1939: 167; Martin, 1957a: 370, fig. 41; Hull, 1962 (1): 299; Martin & Wilcox, 1965: 362 (cat.).

bahamiensis Scarbrough, 1996: 194, figs. 1-8. Type-locality: Bahama Islands, Grande Bahama Island, Freeport. Distr. – Bahamas, Cuba. Type, USNM.

basalis Walker, 1855: 770. Type-locality: Brazil, Amazonas, Tefé (as Ega). Distr. – Brazil (Amazonas). Refs. – Williston, 1891: 68 (cat.); Kertész, 1909: 50 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 2 (cat.).

bengryi Farr, 1963: 10, figs. 4, 5b. Type-locality: Jamaica, Portland, Hardwar Gap, trail to Caledonia Peak. Distr. – Jamaica. Ref. – Martin, 1968e: 3 (cat.). Type, USNM.

bicolor (Macquart), 1848: 191 (1848: 31) (*Gonypes*). Type-locality: “Rio Negro”. Refs. – Walker, 1855: 771; Williston, 1891: 68 (cat.); Kertész, 1909: 50 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.).

cingulipes Walker, 1849: 131. Type-locality: Brazil, “Amazon”. Refs. – Williston, 1891: 68 (cat.); Kertész, 1909: 50 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, ? lost.

clavistylus (Rondani), 1848: 96 (*Gonypes*, as *clavistilus*). Type-locality: “Brazil” [Belém, Pará; cf. Papavero, 1973d: 343]. Refs. – Williston, 1891: 68 (cat.); Kertész, 1909: 51 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.).

concinnata Williston, 1901: 300. Type-locality: Mexico, Guerrero, Omiltemi. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 254 (cat.); Kertész, 1909: 51 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.); Papavero & Ibáñez Bernal, 2003: 174 (hist. Mexican dipterol). Type, BMNH.

crocea Williston, 1901: 300. Type-locality: Mexico, Guerrero, Omiltemi. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 254 (cat.); Kertész, 1909: 51 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.); Papavero & Ibáñez Bernal, 2003: 174 (hist. Mexican dipterol). Type, BMNH.

distincta Schiner, 1867: 375. Type-locality: “Brazil”. Refs. – Williston, 1891: 68 (cat.); Kertész, 1909: 52 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, NMW.

ferruginea Walker, 1855: 771. Type-locality: Brazil, Pará [i. e., Pará, Belém]. Distr. – Brazil (Pará). Refs. – Williston, 1891: 68 (cat.); Kertész, 1909: 52 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, BMNH.

fervens Wiedemann, 1830: 646. Type-locality: “Mexico”. Refs. – Schiner, 1866: 694; Osten Sacken, 1874: 176; Aldrich, 1905: 254 (cat.); Kertész, 1909: 52 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.); Papavero & Ibáñez Bernal, 2001: 75 (hist. Mexican dipterol). Type, BZM [no. 646; cf. Schumann, 1973: 110].

truquii Bellardi, 1861: 187 (87), pl. 2, fig. 18. Type-locality: Mexico, Morelos, Yantepec (as Jantepet). Refs. – Osten Sacken, 1874: 176 (*Leptogaster*); Aldrich, 1905: 254 (cat.); Kertész, 1909: 57 (cat.); Hull, 1962 (1): 299; Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipterol). Type, TORO.

hyacinthina Scarbrough, 1996: 197, figs. 9-16. Type-locality: Dominican Republic, Pedernales, 3.3 km northeast of Los Arroyos. Distr. – Dominican Republic. Ref. – Scarbrough, Perez-Gelabert & Page, 2005: 31 (in key), 45, fig. 63. Type, CMNH.

- intima** Williston, 1901: 301. Type-locality: Mexico, Guerrero, Xucumanatlán. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 254 (cat.); Kertész, 1909: 54 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.); Papavero & Ibáñez Bernal, 2003: 174 (hist. Mexican dipteron). Type, BMNH.
- jamaicensis** Farr, 1963: 9, figs. 3, 5c. Type-locality: Jamaica, St. Thomas, Morant Bay road, 14.5 miles east of Kingston. Distr. – Jamaica. Ref. – Martin, 1968e: 3 (cat.). Type, USNM.
- kamerlacheri** Schiner, 1867: 358. Type-locality: “Brazil”. Refs. – Kertész, 1909: 54 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, NMW (abdomen missing). Type, NMW.
kamerlocheri Williston, 1891: 68 (cat.), error.
- lernerri** Curran, 1953: 1. Type-locality: Bahama Islands, South Bimini Island. Distr. – U. S. A. (Florida), Bahamas (South Bimini). Refs. – Hull, 1962 (1): 299; Martin, 1957b: 4, 6 (in key), fig. 4, 1968: 3 (cat.). Type, AMNH.
- lineata** Scarbrough, 1996: 200, figs. 17-22 (as *lineatus*, error). Type-locality: Dominican Republic, Pedernales, 35 km north-northwest of Cabo Rojo, 1430 m. Distr. – Dominican Republic. Ref. – Scarbrough, Perez-Gelabert & Page, 2005: 31 (in key), 45, figs. 12, 53-55, 63. Type, USNM.
- martini** Farr, 1963: 13, figs. 6-7. Type-locality: Jamaica, St. Thomas, 12.5 miles east of Kingston, Morant Bay road. Distr. – Jamaica. Ref. – Martin, 1968e: 3 (cat.). Type, USNM.
- micropygialis** Williston, 1901: 301. Type-locality: Mexico, Guerrero, La Venta. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 254 (cat.); Kertész, 1909: 54 (cat.); Hull, 1962 (1): 302 (*Psilonyx*); Martin, 1968e: 3 (cat.); Papavero & Ibáñez Bernal, 2003: 174 (hist. Mexican dipteron). Type, BMNH.
- multicincta** Walker, 1851: 155. Type-locality: “South America”. Type, BMNH. Refs. – Walker, 1855: 772; Williston, 1891: 68 (cat.); Kertész, 1909: 54 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, BMNH.
- nigra** Wiedemann, 1828: 533. Type-locality: “Brazil”. Refs. – Walker, 1855: 770; Schiner, 1866: 694, 1867: 356; Williston, 1891: 68 (cat.); Kertész, 1909: 54 (cat.); Bromley, 1946: 105 (cat.); Carrera, 1947d: 266 (preys); Carrera & d’Andretta, 1961: 67 (misident?; preys); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, SENCK [cf. Geller-Grimm, 1999: 212].
- nubeculosa** Bigot, 1878: 455. Type-locality: “Colombia”. Refs. – Williston, 1891: 68 (cat.); Kertész, 1909: 55 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, OXF.
- obscuripes** Loew, 1862: 191 (Cent. 2, no. 13). Type-locality: “Cuba” [Gundlach; cf. Papavero, 1971e: 181-183]. Distr. – U. S. A. (Florida), Cuba. Refs. – Osten Sacken, 1874: 176; Aldrich, 1905: 254 (cat.); Kertész, 1909: 55 cat.); Bromley, 1929a: 273, fig. 1, pl. 1, fig. 16; Martin, 1957a: 366; Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 50 (cat.); Scarbrough, 1996: 202. Syntypes, MCZ, BZM [no. 6492; cf. Schumann, 1973: 140].
ramoni Jaennicke, 1867: 354. Type-locality: “Cuba”. Type, SENCK [cf. Geller-Grimm, 1999: 213].
- pubescens** Curran, 1934c: 323. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica. Distr. – Guyana. Refs. – Hull, 1962 (1): 299; Martin, 1968e: 3 (cat.). Type, ANSP.
- roederi** Williston, 1896a: 305, pl. 11, fig. 80. Type-locality: West Indies, St. Vincent Island. Distr. – St. Vincent Island. Refs. – Aldrich, 1905: 254 (cat.); Kertész, 1909: 56 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 4 (cat.); Scarbrough, 1996: 203, fig. 23. Lectotype BMNH [des. by Scarbrough, 1996: 203].
- rubida** Wiedemann, 1821: 242. Type-locality: “Brazil” [Freyreiss; cf. Papavero, 1971e: 57-58]. Refs. – Wiedemann, 1828: 533; Walker, 1855: 649; Schiner, 1866: 694; Williston, 1891: 69 (cat.); Kertész, 1909: 56 (cat.); Martin, 1968e: 4 (cat.); Geller-Grimm, 1999: 214 (material in SENCK). Syntypes, NMW, BZM [no. 9975; cf. Schumann, 1973: 152].
testacea Loew of Williston, 1901: 299, misident.
- spinitarsis** Bromley, 1951: 3. Type-locality: Panama, Canal Zone, Barro Colorado Island. Distr. – Panama (Canal Zone). Refs. – Hull, 1962 (1): 299; Martin, 1968e: 4 (cat.). Type, USNM.
- titanus** Carrera, 1958a: 147, figs. 6-7. Type-locality: Brazil, São Paulo, Salesópolis. Distr. – Brazil (São Paulo, Santa Catarina). Ref. – Martin, 1968e: 4 (cat.). Type, MZUSP.
- tornowii** Brèthes, 1904: 337. Type-locality: “Argentina”. Refs. – Kertész, 1909: 56 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 4 (cat.).
- triangulata** Williston, 1901: 299, pl. 5, figs. 13, 13a. Type-locality: Mexico, Guerrero, Amula. Distr. – Mexico (Guerrero, Morelos). Refs. – Aldrich, 1905: 254 (cat.); Kertész, 1909: 57 (cat.); Hull, 1962 (1): 299; Papavero & Ibáñez Bernal, 2003: 174 (hist. Mexican dipteron). Type, BMNH.
triangulata Martin, 1968e: 4 (cat.), error.
- tropica** Curran, 1934c: 329. Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Distr. – Guyana. Refs. – Hull, 1962 (1): 299; Martin, 1968e: 4 (cat.). Type, AMNH.
- vitripennis** Schiner, 1867: 356. Type-locality: “Brazil”. Distr. – Brazil (Santa Catarina). Refs. – Williston, 1891: 69 (cat.); Kertész, 1909: 57 (cat.); Bromley, 1946: 105 (cat.); Hull, 1962 (1): 299; Martin, 1968e: 4 (cat.). Type, NMW.
- vorax** Curran, 1934c: 329. Type-locality: Guyana, Potaro-Siparuni, Tumatumari. Distr. – Guyana. Refs. – Hull, 1962 (1): 299; Martin, 1968e: 4 (cat.). Type, AMNH.

Genus *Leptopteromyia* Williston, 1907

Leptopteromyia Williston, 1907: 1. Type-species, *Leptopteromyia gracilis* Williston, 1907 (sub. mon., Williston, 1908: 195,

figs. 35, 77).

Ref. – Martin, 1971 (key to spp.).

- argentinae** Martin, 1972: 270. Type-locality: Argentina, Tucumán, Dique El Cadillal [as Dioovo Codillal, error]. Distr. – Argentina (Tucumán). Type, IML.
- argentinae* Martin, 1971: 266 (only in key, without description), *nomen nudum*.
- brasiliæ** Martin, 1971: 266 (in key), 268. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (São Paulo, Santa Catarina). Type, MZUSP.
- colombiae** Martin, 1971: 266 (in key), 269. Type-locality: Colombia, Cundinamarca, 11 miles west of Congueza, 1340 m. Distr. – Colombia (Cundinamarca). Type, CAS.
- gracilis** Williston, 1907: 195, figs. 35, 77. Type-locality: “Brazil”. Distr. – Brazil (Rio de Janeiro). Refs. – Carrera, 1947b: 91, figs. 1-7, 94 (puparium); D. E. Hardy, 1947: 72, fig. 2; Hull, 1962 (1): 316; Martin, 1968e: 6 (cat.), 1971: 266 (in key), 267. *willistoni* Hermann, 1926a: 143, *nomen nudum*.
- lopesi** Martin, 1971: 266 (in key), 267. Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (Manguinhos). Distr. – Brazil (Rio de Janeiro).
- mexicanae** Martin, 1971: 266 (in key), 269. Type-locality: Mexico, Sonora, Bahía San Carlos. Distr. – Mexico (Sonora). Type, CAS.
- peruæ** Martin, 1971: 266 (mentions specimen bred from Embioptera nests), 266 (in key), 267. Type-locality: Peru, Huánuco, Tingo María. Distr. – Peru (Huánuco). Type, CAS.

Genus **Psilonyx** Aldrich, 1923

Psilonyx Aldrich, 1923: 5. Type-species, *Leptogaster annulatus* Say, 1823 (orig. des.).

- annulatus** (Say), 1823: 75 (1859: 75) (*Leptogaster*). Type-locality: U. S. A., Pennsylvania. Distr. – U. S. A., Central America, Puerto Rico, northern South America. Refs. – Back, 1909: 159 (*Leptogaster*); Aldrich, 1923: 5 (n. comb.); Engel, 1930: 457, fig. 1 (Bolivia, Argentina; misident.); Bromley, 1934c: 164 (*Leptogaster (Psilonyx)*); Bromley, 1950: 230 (*Leptogaster (Psilonyx)*); Martin, 1957a: 352, fig. 25; 1957b: 6 (in key), 7; Hull, 1962 (1): 302; Martin, 1965a: 18; Martin & Wilcox, 1965: 363 (cat.); Martin, 1968e: 4 (cat.).
- histrio* Wiedemann, 1828: 535 (*Leptogaster*). Type-locality: U. S. A., Pennsylvania.
- arawak** Farr, 1963: 7, fig. 1. Type-locality: Jamaica, St. Andrew, Ferry. Distr. – Jamaica. Ref. – Martin, 1968e: 4 (cat.). Type, USNM.
- macropygialis** (Williston), 1901: 301, pl. 5, fig. 14 (*Leptogaster*). Type-locality: Mexico, Tabasco, Teapa. Distr. – Mexico (Tabasco). Refs. – Aldrich, 1905: 254 (cat.; *Leptogaster*); Kertész, 1909: 54 (cat.; *Leptogaster*); Hull, 1962 (1): 302; Martin, 1968e: 4 (cat.); Papavero & Ibáñez Bernal, 2003: 174 (hist. Mexican dipterol). Type, BMNH.
- magnicauda** (Curran), 1934c: 328 (*Leptogaster*). Type-locality: Guyana, Cuyuni-Mazaruni, Bartica, Kartabo. Distr. – Guyana. Refs. – Hull, 1962 (1): 302; Martin, 1968e: 4 (cat.). Type, AMNH.
- venustus** (Bromley), 1929a: 274, fig. 2 (*Leptogaster*). Type-locality: “Cuba”. Distr. – Cuba. Refs. – Hull, 1962 (1): 299 (*Leptogaster*); Martin, 1968e: 4 (cat.); Alayo Dalmau & García Ávila, 1983: 50 (cat.). Type, MCZ.
- zephyrus** Scarbrough & Page in Scarbrough, Perez-Gelabert & Page, 2005: 30 (in key), 45, figs. 6, 56-61, 63. Type-locality: Dominican Republic, Pedernales, Upper Las Abejas, 38 km north-northwest of Cabo Rojo, 1350 m. Distr. – Dominican Republic. Type, CMNH.

Genus **Schildia** Aldrich, 1923

Schildia Aldrich, 1923: 4. Type-species, *Schildia microthorax* Aldrich, 1823 (orig. des.).

Shannomyioleptus Carrera, 1944: 87. Type-species, *Shannomyioleptus fragilis* Carrera, 1944 (orig. des.).

Ref. – Martin, 1975b (key to spp.).

- alphus** Martin, 1975b: 190 (in key and text). Type-locality: Brazil, Ceará, Cascavel (Sítio B. Vista). Distr. – Brazil (Ceará). Type, MZUSP.
- fragilis** (Carrera), 1944: 87, figs. 1-3 (*Shannomyioleptus*). Type-locality: Brazil, Mato Grosso do Sul, Maracaju. Distr. – Brazil (Mato Grosso do Sul). Refs. – Hull, 1962 (1): 314, figs. 205, 589, 1194, 1203, 1584 (*Schildia (Shannomyioleptus)*); Martin, 1968e: 5 (cat.), 1975b: 190 (in key). Type, MZUSP.
- gracillima** (Walker), 1855: 722 (*Leptogaster*). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Distr. – Brazil (Pará). Refs. –

Williston, 1891: 68 (cat.; *Leptogaster*); Kertész, 1909: 53 (cat.; *Leptogaster*); Bromley, 1946: 105 (cat.; *Leptogaster*); Hull, 1962 (1): 299 (*Leptogaster*); Martin, 1968e: 5 (cat.), 1975b: 190 (in key). Type, BMNH.
guatemalae Martin, 1975b: 190 (in key), 191. Type-locality: Guatemala, Santa Rosa, Guazacápan. Distr. – Guatemala (Santa Rosa). Type, USNM.
jamaicensis Farr, 1962: 191, figs. 1-8. Type-locality: Jamaica, St. Andrew, Long Mountain., Distr. – Jamaica. Refs. – Farr, 1963: 19; Martin, 1968e: 5 (cat.), 1975b: 190 (in key). Type, USNM.
microthorax Aldrich, 1923: 4. Type-locality: Costa Rica, San Mateo, Higuito. Distr. – Costa Rica (San Mateo), Peru. Refs. – Hull, 1962 (1): 314, figs. 201, 588, 1192, 1201; Martin, 1968e: 5 (cat.), 1975b: 190 (in key); Fisher, 1984: 289. Type, USNM.
ocellata Martin, 1975b: 189 (in key), 192. Type-locality: Brazil, Amapá, rio Amapari. Distr. – Brazil (Amapá). Type, MZUSP.
zona Martin, 1975b: 190 (in key), 192. Type-locality: Panama, Canal Zone, Piña area. Distr. – Panama (Canal Zone). Type, UUEC.

Genus **Systelogaster**, nom. nov.

Systelogaster Hermann, 1926a: 149 (preocc. Gahan, 1917). Type-species, *Euscelidia fascipennis* Schiner, 1867 (orig. des.).
Systelogaster, nom. nov. for *Systelogaster* Hermann, 1912. Type-species, *Euscelidia fascipennis* Schiner, 1867 (aut.).

alba (Martin), 1975a: 2 (in key and text) (*Systelogaster*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (Santa Catarina). Type, ZSBS. **N. comb.**
?breviventris (Rondani), 1848: 96 (*Gonyptes*). Type-locality: “Brazil” [Pará, Belém; cf. Papavero, 1973d: 343]. Refs. – Williston, 1891: 68 (*Leptogaster*); Kertész, 1909: 50 (*Leptogaster*); Bromley, 1946: 105 (cat.; *Leptogaster*); Hull, 1962 (1): 311 (*Systelogaster*); Martin, 1968e: 6 (cat.; *Systelogaster*). Type, **N. comb.**
calceolata (Bigot), 1878: 430 (*Lasiocnemus*). Type-locality: Brazil, “Amazon”. Refs. – Williston, 1891: 69 (cat.; *Lasiocnemus*); Kertész, 1909: 57 (*Lasiocnemus*); Bromley, 1946: 105 (cat.; *Lasiocnemus*); Hull, 1962 (1): 311 (*Systelogaster*); Martin, 1968a: 6 (cat.; *Systelogaster*), 1975a: 2 (in key), 3 (*Systelogaster*). Type, OXF. **N. comb.**
fascipennis (Schiner), 1867: 359 (*Euscelidia*). Type-locality: “Brazil”. Distr. – Brazil (São Paulo, Santa Catarina, Rio Grande do Sul). Refs. – Williston, 1891: 69 (cat.; *Euscelidia*); Bromley, 1946: 105 (cat.; *Euscelidia*); Hull, 1962 (1): 311, figs. 206B, 602, 1180, 2052, 2059 (*Systelogaster*); Martin, 1968e: 6 (cat.; *Systelogaster*), 1975a: 2 (in key), 3 (*Systelogaster*). Type, NMW. **N. comb.**
parva (Martin), 1975a: 2 (in key), 4 (*Systelogaster*). Type-locality: Paraguay, Cordillera, San Bernardino. Distr. – Paraguay (Cordillera). Type, ZSBS. **N. comb.**
silacea (Martin), 1975a: 2 (in key), 4 (*Systelogaster*). Type-locality: Peru, Pasco, Puerto Yessup, río Pichis. Distr. – Peru (Pasco). Type, ZSBS. **N. comb.**
aurantiaca Hermann, 1926a: 151 (*Systelogaster*), *nomen nudum*. Ref. – Hull, 1962 (1): 311 (*Systelogaster*); Martin, 1975a: 4 (*Systelogaster*).

Genus **Tipulogaster** Cockerell, 1913

Tipulogaster Cockerell, 1913: 214. Type-species, *Leptogaster badius* Loew, 1862 (orig. des.) = *glabrata* (Wiedemann, 1828).

Tipulogaster glabrata (Wiedemann, 1828), an extant species, is also known as fossil (Oligocene; U. S. A., Colorado).

lancea Tomasovic, 2002: 89, figs. 3a-d. Type-locality: French Guiana, Saül, Mont Belvédère. Distr. – French Guiana. Type, FUSA Gembloix.

Subfamily **Stenopogoninae** Hull, 1962

Stenopogonini Hull, 1962 (1): 118. Type-genus, *Stenopogon* Loew, 1847.
 Stenopogoninae Paramonov, 1966: 141.

Tribe **Acronychini** Artigas & Papavero, 1991

Acronychini Artigas & Papavero, 1991a: 141, 142. Type-genus, *Acronyches* Williston, 1908.

Genus **Acronyches** Williston, 1866

Acronyches Williston, 1908: 388, fig. 103. Type-species, *Acronyches willistoni* Hermann, 1921 (sub. des., Hermann, 1921: 121).

Megonyx Hull, 1962 (1): 312. Type-species, *Megonyx giganteus* Hull, 1962 (orig. des.) = *Acronyches fenestratus* Hermann, 1921.

Ref. – Papavero, 1971a (key to spp.).

alexanderi Papavero, 1971a: 147. Type-locality: Brazil, São Paulo, Mogi das Cruzes. Distr. – Brazil (Rio de Janeiro, São Paulo). Ref. – Artigas & Papavero, 1991b: 248. Type, MZUSP.

fenestratus Hermann, 1921: 122. Type-locality: Paraguay, Itapúa, Santísima Trinidad (as Santa Trinidad), and Argentina. Distr. – Peru and Brazil (Amazonas) to Paraguay and Argentina. Refs. – Hull, 1962 (1): 312; Fisher, 1984: 27; Artigas & Papavero, 1991b: 248, figs. 1-2.

giganteus Hull, 1962 (1): 313, fig. 204 (*Megonyx*). Type-locality: Brazil, Amazonas, Manaus. Type, AMNH.

geosarginus Papavero, 1971a: 148. Type-locality: Surinam, Paramaribo. Distr. – Surinam. Ref. – Artigas & Papavero, 1991b: 248. Type, MZUSP.

imitator Hermann, 1921: 122. Type-locality: “Paraguay”. Distr. – Brazil (Mato Grosso), Paraguay. Refs. – Hull, 1962 (1): 312, figs. 190, 601, 1178, 1186, 2062; Artigas & Papavero, 1991b: 248. Type, ZSBS.

maya Martin, 1968d: 180. Type-locality: Mexico, Yucatán, Chichén Itzá. Distr. – Mexico (Yucatán). Ref. – Artigas & Papavero, 1991b: 248. Holotype, FSCA.

meruuna Papavero, 1971a: 150. Type-locality: Costa Rica, Turrialba. Distr. – Costa Rica. Ref. – Artigas & Papavero, 1991b: 248. Type, MCZ.

plutactites Papavero, 1971a: 150. Type-locality: Costa Rica, Turrialba. Distr. – Costa Rica. Ref. – Artigas & Papavero, 1991b: 248. Type, MCZ.

rarus Martin, 1968d: 181. Type-locality: Mexico, Morelos, Yautepec. Distr. – Mexico (Morelos). Ref. – Artigas & Papavero, 1991b: 248. Type, CAS.

westcotti Martin, 1968d: 182. Type-locality: Mexico, Nayarit, 20 miles southeast of Ixtlán del Río. Distr. – Mexico (Nayarit). Type, CAS. Ref. – Artigas & Papavero, 1991b: 248. Type, CAS.

willistoni Hermann, 1921: 121. Type-locality: “Surinam”. Distr. – Surinam. Refs. – Hull, 1962 (1): 312; Artigas & Papavero, 1991b: 248.

Tribe **Bathypogonini** Artigas & Papavero, 1991

Bathypogonini Artigas & Papavero, 1991a: 140, 142. Type-genus, *Bathypogon* Loew, 1851.

Genus **Carebaricus** Artigas & Papavero, 1991

Carebaricus Artigas & Papavero, 1991b: 249. Type-species, *Stenopogon rionegrensis* Lamas M., 1971 (orig. des.).

rionegrensis (Lamas), 1971a: 15, figs. 1-5 (*Stenopogon*). Type-locality: Argentina, Río Negro, río Colorado. Distr. – Argentina (Río Negro, Neuquén). Ref. – Artigas & Papavero, 1991b: 249, figs. 3-10 (n. comb.). Type, MZUSP.

Tribe **Cyrtopogonini** Artigas & Papavero, 1991

Cyrtopogonini Artigas & Papavero, 1991a: 141, 142. Type-genus, *Cyrtopogon* Loew, 1847.

Genus **Ablautus** Loew, 1866

Ablautus Loew, 1866: 37 (Cent. 7, no. 63). Type-species, *Ablautus trifarius* Loew, 1866 (mon.).

Albautatus Loew, 1874: 377, *emend.*

- arnaudi** Wilcox, 1966d: 675. Type-locality: Mexico, Baja California Norte, San Felipe. Distr. – U. S. A. (California), Mexico (Baja California Norte) TP CAS.
- californicus** Wilcox, 1935: 226. Type-locality: U. S. A., California. Distr. – U. S. A. (California), Mexico. Refs. – Hull, 1962 (1): 204; Martin & Wilcox, 1965: 364 (cat.); Wilcox, 1966d: 677. Type, CAS.
- coachellus** Wilcox, 1966d: 677. Type-locality: U. S. A., California, Indio. Distr. – U. S. A. (Arizona, California, New Mexico), Mexico (Sonora). Type, CAS.
- coquilletti** Wilcox, 1935: 226. Type-locality: U. S. A., California. Distr. – U. S. A. (California, Utah), Mexico. Refs. – Hull, 1962 (1): 204; Martin & Wilcox, 1965: 364 (cat.); Nelson, 1987: 59. Type, USNM.
- flavipes** Coquillett, 1904b: 178. Type-locality: U. S. A., California. Distr. – U. S. A. (California, Arizona, New Mexico, Utah), Mexico (Baja California, Sonora, Durango). Refs. – Kertész, 1909: 86 (cat.); Hull, 1962 (1): 204; Martin & Wilcox, 1965: 364 (cat.); Martin & Papavero, 1970: 21 (cat.); Nelson, 1987: 59; Artigas & Papavero, 1991f: 59.
- mimus** (Osten Sacken), 1877: 290 (*Ablautatus*). Type-locality: U. S. A., California. Distr. – U. S. A. (California, Arizona, New Mexico, Utah), Mexico (Baja California). Type. . Refs. – Williston, 1885: 53 (*Ablautatus*); Aldrich, 1905: 255 (cat.); Kertész, 1909: 86 (cat.); Hull, 1962 (1): 204; Martin & Wilcox, 1965: 364 (cat.); Nelson, 1987: 59.
- squamipes** Cole, 1924: 11. Type-locality: U. S. A., Arizona, Chiricahua Mountains. Refs. – Hull, 1962 (1): 204, figs. 141, 1050, 1059, 1795, 100; Martin & Wilcox, 1965: 364 (cat.). Type, CAS.
- trifarius** Loew, 1866: 36 (Cent. 7, no. 63). Type-locality: U. S. A., California. Distr. – U. S. A. (California), ?Mexico (Osten Sacken, 1887: 168). Refs. – Osten Sacken, 1887: 168 (*Ablautatus*); Aldrich, 1905: 155 (cat.); Kertész, 1909: 87 (cat.); Hull, 1962 (1): 204, fig. 540; Martin & Wilcox, 1965: 364 (cat.); Martin & Papavero, 1970: 21 (cat.); Artigas & Papavero, 1991f: 59. Type, MCZ.

Genus **Callinicus** Loew, 1872

- Callinicus** Loew, 1872a: 71. Type-species, *Callinicus calcaneus* Loew, 1872 (mon.).
Chrysoceria Williston, 1907: 1. Type-species, *?Laparus pictitarsis* Bigot, 1878 (orig. des.).

- pictitarsis** (Bigot), 1878: 417 (*?Laparus*) Type-locality: U. S. A., California. Distr. – U. S. A. (California, Arizona, Utah), Mexico (Baja California). Refs. – Hull, 1962 (1): 168; Martin & Wilcox, 1965: 365 (cat.).

Genus **Cyrtopogon** Loew, 1847

- Cyrtopogon** Loew, 1847: 516. Type-species, *Asilus ruficornis* Fabricius, 1794 (sub. des., Rondani, 1856:).
Euarmostus Walker, 1851: 102. Type-species, *Euarmostus bimacula* Walker, 1851 (mon.).
Enarmostus Walker, 1855: 423, error.
Philammosius Rondani, 1856: 156. Type-species, *Dasytopogon fimbriatus* Meigen, 1820 (orig. des.).

- basingeri** Wilcox & Martin, 1936a: 49. Type-locality: U. S. A., California, San Bernardino County, Forest Home. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Hull, 1962 (1): 172; Martin & Wilcox, 1965: 368 (cat.). Type, CAS.
vandykei Wilcox & Martin, 1936a: 27, pl. 3. Type-locality: U. S. A., California, San Bernardino County, Forest Home. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Hull, 1962 (1): 172; Martin & Wilcox, 1965: 368 (cat.). Type, CAS.

Genus **Dasycyrton** Philippi, 1865

- Dasycyrton** Philippi, 1865: 701. Type-species, *Dasycyrton gibbosus* Philippi, 1865 (mon.).

- arrayanensis** Artigas, 1970a: 93, figs. 70, 83, 478. Type-locality: Chile, Santiago, Arrayán. Distr. – Chile (Santiago, Biobío). Ref. – Artigas & Papavero, 1991f: 59. Type, DZUC.
coquimbensis Artigas, 1970a: 95, figs. 78, 477. Type-locality: Chile, Choapa, Illapel. Distr. – Chile (Coquimbo). Ref. – Artigas & Papavero, 1991f: 59. Type, DZUC.
gibbosus Philippi, 1865: 701, pl. 26, figs. 30, 30a. Type-locality: Chile, Santiago and Aconcagua: Hacienda Catemú. Neotype-locality: Chile, Coquimbo [as Atacama, error], Cerro Caquis. Distr. – Chile (Coquimbo, Santiago). Refs. – Williston, 1891: 71 (cat.); Kertész, 1901: 75 (cat.); Hull, 1962 (1): 139; Martin & Papavero, 1970: 16 (cat.); Artigas, 1970a: 97, figs. 82, 407, 492; Artigas & Papavero, 1991f: 59. Syntypes, lost. NT DZUC [des. by Artigas, 1970a: 97].
medinai Artigas, 1970a: 99, fig. 86 (as *medinae*, error). Type-locality: “Chile”. Distr. – Chile. Ref. – Artigas & Papavero,

1991f: 59. Type, MNHNS.
sucinopedis Artigas, 1970a: 105, figs. 80, 480. Type-locality: Chile, Valparaíso, Valparaíso. Distr. – Chile (Valparaíso). Ref. – Artigas & Papavero, 1991f: 59. Type, DZUC.

Genus **Dasyppecus** Philippi, 1865

Dasyppecus Philippi, 1865: 692. Type-species, *Dasyppecus heteroneurus* Philippi, 1865 (sub. des., Hull, 1962 (1): 139).

heteroneurus Philippi, 1865: 692, pl. 28, figs. 54, 54a, 54b. Type-locality: “Chile”. Neotype-locality: Chile, Quebrada de Paihuano. Distr. – Chile (Coquimbo, Santiago). Refs. – Williston, 1891: 72 (cat.); Kertész, 1909: 90 (cat.); Bromley, 1932: 166; Stuardo Ortiz, 1946: 81 (cat.); Hull, 1962 (1): 139, figs. 2525–2526; Oldroyd, 1965: 351; Artigas, 1970a: 110, figs. 63–64, 66, 463; Martin & Papavero, 1970: 16 (cat.); Artigas, 1971: 22, fig. 45; Artigas & Papavero, 1991f: 59. Type, lost. NT BMNH [des. by Oldroyd, 1965: 351].

?**latus** (Philippi), 1965: 686 (*Dasypogon*). Type-locality: Chile, Choapa, Illapel. Refs. – Bigot, 1878: 220; Williston, 1891: 68 (*Dasypogon*); Kertész, 1909: 90 (cat.); Hull, 1962 (1): 139; Martin & Papavero, 1970: 16 (cat.); Artigas & Papavero, 1991f: 59. Type, lost.

Genus **Graptostylus** Hull, 1962

Graptostylus Hull, 1962 (1): 207. Type-species, *Graptostylus dolosus* Hull, 1962 (orig. des.).

dolosus Hull, 1962 (1): 208. Type-locality: Chile, Valparaíso. Distr. – Chile (Coquimbo, Santiago). Refs. – Artigas, 1970a: 119, figs. 65, 67–69, 71, 74, 457, 1971: 23, figs. 43–44; Artigas & Papavero, 1991f: 59. Type, USNM.

Genus **Hadrokolos** Martin, 1959

Hadrokolos Martin, 1959: 3. Type-species, *Holopogon texanus* Bromley, 1934 (orig. des.).

notialis Martin, 1967b: 194. Type-locality: Mexico, Oaxaca, Tetalapán. Distr. – Mexico (Oaxaca). Ref. – Martin & Papavero, 1970: 18 (cat.). Type, KU.

Genus **Heteropogon** Loew, 1847

Dasypogon, subg. **Heteropogon** Loew, 1847: 488. Type-species, *Dasypogon manicatus* Meigen, 1820 (sub. des., Back, 1909: 318).

Anisopogon Loew, 1874: 377 (unjustified *nom. nov.* for *Heteropogon* Loew, 1847). Type-species, *Dasypogon manicatus* Meigen, 1820 (aut.).

Dasypogon, subg. *Pycnopogon* Loew, 1847: 526. Type-species, *Dasypogon (Pycnopogon) mixtus* Loew, 1847 (sub. des., Rondani, 1856: 157).

dejectus (Williston), 1901: 306 (*Holopogon*). Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 261 (cat.); Kertész, 1909: 59 (cat.); Hull, 1962 (1): 177 (*Holopogon*); Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 59; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Type, BMNH.

divisus (Coquillett), 1902: 139 (*Pycnopogon*). Type-locality: Mexico, Chihuahua. Distr. – Mexico (Chihuahua). Refs. – Aldrich, 1905: 262 (*Pycnopogon*); Kertész, 1909: 89 (cat.; *Pycnopogon*); Hull, 1962 (1): 189 (*Pycnopogon*); Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 59. Type, USNM.

dorothyae Martin, 1962c: 373. Type-locality: Mexico, Guerrero, 32 miles north of Chilpancingo. Distr. – Mexico (Guerrero, Morelos). Refs. – Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 59. Type, KU.

duncani Wilcox, 1941: 54. Type-locality: U. S. A., Arizona, Baboquivari Mountains. Distr. – U. S. A. (Arizona), Mexico (Sonora). Ref. – Martin & Wilcox, 1965: 374 (cat.). Type, CAS.

ludius (Coquillett), 1893a: 20 (*Anisopogon*). Type-locality: U. S. A., California, Los Angeles and San Bernardino Counties; Canada, British Columbia. Distr. – Canada, U. S. A. (California), Mexico (Baja California). Refs. – Brown, 1929: 305,

- pl. 29, figs. 24-25; Pritchard, 1935: 2 (in key); Martin & Wilcox, 1965: 374 (cat.).
- paurosomus** Pritchard, 1935: 3 (in key), 4. Type-locality: U. S. A., Arizona, Tucson, Santa Catalina Mts. Distr. – U. S. A. (Arizona), Mexico (Sonora, Sinaloa). Refs. – Martin & Wilcox, 1965: 374 (cat.); Lavigne & Bollington, 1999: 9 (ethology). Type, AMNH.
- rejectus** Williston, 1901: 307. Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. – U. S. A. (Arizona), Mexico (Guerrero). Refs. – Aldrich, 1905: 263 (cat.); Kertész, 1909: 66 (cat.); Martin & Wilcox, 1965: 374 (cat.); Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 59; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipteral). Type, BMNH.
- spatulatus** Prichard, 1935: 3 (in key), 5. Type-locality: U. S. A., Arizona, Hayden. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Martin & Wilcox, 1965: 374 (cat.); Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 59. Type, AMNH.
- willistoni** Martin, 1962c: 375. Type-locality: Mexico, Morelos, highway 136, near Cuernavaca. Distr. – Mexico (Morelos, Guerrero). Refs. – Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 59. Type, KU.

Genus **Holopogon** Loew, 1847

Dasypogon, subg. **Holopogon** Loew, 1847: 473. Type-species, *Dasypogon nigripennis* Meigen, 1820 (sub. des., Coquillett, 1910: 552).

One species known as fossil: *Holopogon archilestes* Hull (1962: 272) (Oligocene; U. S. A.).

- atrifrons** Cole, 1924: 8. Type-locality: U. S. A., California, San Bernardino County, Mill Creek Canyon. Distr. – U. S. A. (Oregon, California, Nevada), Mexico (Baja California). Refs. – Hull, 1962 (1): 177; Martin & Wilcox, 1965: 375 (cat.). Type, CAS.
- currani** Martin, 1959: 17. Type-locality: U. S. A., Arizona, Winona. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Martin & Wilcox, 1965: 375 (cat.); Martin & Papavero, 1970: 19 (cat.); Artigas & Papavero, 1991f: 60. Type, AMNH.
- fisheri** Martin, 1967b: 195. Type-locality: Mexico, Coahuila, Saltillo, Canyon de la Carbonera. Distr. – Mexico (Coahuila). Refs. – Martin & Papavero, 1970: 19 (cat.); Artigas & Papavero, 1991f: 60. Type, CAS.
- pulcher** Williston, 1901: 306, pl. 5, fig. 18. Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. – Mexico (Guerrero, Morelos). Refs. – Aldrich, 1905: 262 (cat.); Kertész, 1909: 61 (cat.); Hull, 1962 (1): 177; Martin & Papavero, 1970: 19 (cat.); Artigas & Papavero, 1991f: 60; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipteral). Type, BMNH.
- violaceus** Williston, 1901: 306, pl. 5, fig. 17. Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 262 (cat.); Kertész, 1909: 62 (cat.); Hull, 1962 (1): 177; Martin & Papavero, 1970: 19 (cat.); Artigas & Papavero, 1991f: 60; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipteral). Type, BMNH.

Genus **Itolia** Wilcox, 1936

Itolia Wilcox, 1936b: 201. Type-species, *Itolia maculata* Wilcox, 1936 (orig. des.).

- atripes** Wilcox, 1949: 191 (in key), 193. Type-locality: U. S. A., Arizona, Mohawk. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Hull, 1962 (1): 200; Martin & Wilcox, 1965: 376 (cat.); Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 60. Type, CAS.
- fascia** Martin, 1966b: 213. Type-locality: Mexico, Puebla, Petlalcingo. Distr. – Mexico (Puebla). Refs. – Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991g: 60. Type, UCD.
- maculata** Wilcox, 1936b: 202, fig. 1. Type-locality: U. S. A., Arizona, Santa Rita Mts. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Wilcox, 1949: 191 (in key); Hull, 1962 (1): 200, figs. 70, 593, 989, 1832-1833, 1954; Martin & Wilcox, 1965: 376 (cat.); Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 60. Type, CAS.
- pilosa** Martin, 1966b: 214. Type-locality: Mexico, Sonora, Vicam. Distr. – Mexico (Sonora). Refs. – Martin & Papavero, 1970: 20 (cat.); Artigas & Papavero, 1991f: 60. Type, CAS.
- timberlakei** Wilcox, 1949: 191 (in key), 192. Type-locality: U. S. A., California, Riverside County, White Water. Distr. – U. S. A. (California, Arizona), Mexico (Baja California). Refs. – Hull, 1962 (1): 200; Martin & Wilcox, 1965: 376 (cat.). Type, CAS.

Genus **Ivettea** Artigas & Papavero, 1991

Ivettea Artigas & Papavero, 1991f: 60. Type-species, *Dasycyrton minusculus* Artigas, 1970 (orig. des.).

minuscula (Artigas), 1970a: 101, figs. 77, 425 (*Dasycyrton*). Type-locality: Chile, Linares, El Radal. Distr. – Chile (Linares, Valdivia). Refs. – Artigas, 1971: 20, figs. 39-40 (*Dasycyrton*); Artigas & Papavero, 1991f: 60 (n. comb.). Type, DZUC.

Genus **Lonquimayus** Artigas & Papavero, 1991

Lonquimayus Artigas & Papavero, 1991f: 60. Type-species, *Holopogon tener* Bigot, 1878 (orig. des.).

notocinereatus (Artigas), 1970a: 102, figs. 85, 494 (*Dasycyrton*). Type-locality: Chile, Malleco, Angol. Distr. – Chile (Malleco). Ref. – Artigas & Papavero, 1991g: 60 (n. comb.). Type, DZUC.

papaveroi (Artigas), 1970a: 104, figs. 81, 491. Type-locality: Chile, Biobío, Los Ángeles, Fundo San José. Distr. – Chile (Biobío, Cautín, Malleco). Ref. – Artigas & Papavero, 1991f: 60 (n. comb.). Type, DZUC.

tener (Bigot), 1878: 437 (*Holopogon*; as *tenerum*). Type-locality: “Chile”. Distr. – Chile (Arauco, Concepción, Malleco, Ñuble, Santiago). Refs. – Williston, 1891: 71 (cat.; *Holopogon*, as *tenerus*); Kertész, 1909: 61 (cat.; *Holopogon*); Stuardo Ortiz, 1946: 80 (cat.; *Holopogon*; as *tenerum*); Hull, 1962 (1): 177 (*Holopogon*); Artigas, 1970a: 107, figs. 72-73, 75-76, 84, 487 (*Dasycyrton*), 1971: 21, fig. 4 (*Dasycyrton*); Martin & Papavero, 1970: 19 (cat.); Artigas & Papavero, 1991f: 60 (n. comb.). Type, OXF.

Genus **Metapogon** Coquillett, 1904

Metapogon Coquillett, 1904b: 181. Type-species, *Metapogon trifarius* Coquillett, 1901 (orig. des.).

carinatus Wilcox, 1964: 192 (in key), 193. Type-locality: U. S. A., California, Riverside County, White Water. Distr. – U. S. A. (California, Utah), Mexico (Baja California). Type, CAS.

gibber (Williston), 1883: 14 (?*Cyrtopogon*). Type-locality: U. S. A., California. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Back, 1909: 293 (*Cyrtopogon*); Wilcox & Martin, 1936: 2; Hull, 1962 (1): 203, figs. 124, 570, 987, 996; Wilcox, 1964: 192 (in key), 193; Martin & Wilcox, 1965: 378 (cat.). Type, KU.

leechi Wilcox, 1964: 192 (in key), 197. Type-locality: Mexico, Baja California Sur, 3 miles southwest of Misión San Javier. Distr. – Mexico (Baja California). Ref. – Artigas & Papavero, 1991f: 61. Type, CAS.

punctipennis Coquillett, 1904b: 182. Type-locality: U. S. A., New Mexico, Organ. Distr. – U. S. A. (California, Arizona, New Mexico), Texas, Mexico (Coahuila). Refs. – Back, 1909: 305; Hull, 1962 (1): 203; Wilcox, 1964: 192 (in key), 198; Martin & Wilcox, 1965: 378 (cat.). Type, USNM.

tricellus Wilcox, 1964: 192 (in key), 199. Type-locality: U. S. A., Riverside County, White Water. Distr. – U. S. A. (California, Nevada), Mexico (Baja California). Type, CAS.

Genus **Nannocyrtopogon** Wilcox & Martín, 1936

Nannocyrtopogon Wilcox & Martín, 1936b: 449. Type-species, *Cyrtopogon cerussatus* Osten Sacken, 1877 (orig. des.).

lestomyiformis Wilcox & Martín, 1936b: 454, pl. 1. Type-locality: U. S. A., California, San Diego County, Desert edge. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Wilcox & Martín, 1957: 385; Hull, 1962 (1): 175; Martín & Wilcox, 1965: 379 (cat.). Type, OSU.

Genus **Nothopogon** Artigas & Papavero, 1991

Nothopogon Artigas & Papavero, 1991f: 61. Type-species, *Nothopogon triangularis* Artigas & Papavero, 1991 (orig. des.).

triangularis Artigas & Papavero, 1991f: 61, figs. 14-15. Type-locality: Argentina, Salta, El Carmen, 27 km south of Molinos, 1900 m. Distr. – Argentina (Salta). Type, DZUC.

Genus **Raulcortesia** Artigas & Papavero, 1991

Raulcortesia Artigas & Papavero, 1991f: 62. Type-species, *Dasycyrton lanosus* Artigas, 1970 (orig. des.).

lanosa (Artigas), 1970a: 98, figs. 79, 490 (*Dasycyrton*). Type-locality: Chile, Curicó, Palos Negros. Distr. – Chile (Curicó). Refs. – Artigas, 1971: 20, figs. 37–38 (*Dasycyrton*); Artigas & Papavero, 1991f: 62 (n. comb.). Type, DZUC.

Genus **Sintoria** Hull, 1962

Sintoria Hull, 1962 (1): 197. Type-species, *Sintoria emeralda* Hull, 1962 (orig. des.).

Ref. – Wilcox, 1972 (rev.).

emeralda Hull, 1962 (1): 199. Type-locality: Mexico, México, [San Miguel de] Neplanta. Distr. – Mexico (Aguascalientes, Guanajuato, Jalisco, México, San Luis Potosí, Veracruz). Refs. – Wilcox, 1972: 55; Artigas & Papavero, 1991f: 62. Type, not in USNM.

lagunae Wilcox, 1972: 55. Type-locality: Mexico, Baja California Sur, La Laguna, Sa. Victoria. Distr. – Mexico (Baja California Sur). Ref. – Artigas & Papavero, 1991f: 62. Type, CSCLB.

rossi Wilcox, 1972: 58. Type-locality: Mexico, Mt. Popocatépetl, north slope, 11,000 ft. Distr. – Mexico. Ref. – Artigas & Papavero, 1991f: 62. Type, CAS.

Genus **Wilcoxia** James, 1941

Wilcoxia James, 1941: 38. Type-species, *Wilcoxia cinerea* James, 1941 (orig. des.).

†sp. Poinar Jr., 1992: 178. Dominican Republic (Oligocene/Miocene).

Unrecognized species of Cyrtopogonini

bullatus Wulp, 1882b: 100 (*Holopogon*). Type-locality: “Argentina”. Refs. – Williston, 1891: 71 (cat.); Kertész, 1909: 59 (cat.); Hull, 1962 (1): 177 (*Holopogon*); Martin & Papavero, 1970: 19 (cat.). Type, ZMA.

Tribe Dioctriini Hendel, 1936

Dioctriini Hendel, 1936: 1923. Type-genus, *Dioctria* Meigen, 1803.

Dioctriinae Enderlein, 1936: 92.

Dioctrini Lehr, 1977: 48, 51.

Diochiini Arnett, 1985: 675.

Genus **Nannodioctria** Wilcox & Martin, 1942

Dioctria, subg. *Neodioctria* Wilcox & Martin, 1941: 7 (preocc. Ricardo, 1912). Type-species, *Dioctria* (*Neodioctria*) *albicornis* Wilcox & Martin, 1941 (orig. des.).

Dioctria, subg. **Nannodioctria** Wilcox & Martin, 1942: 35 (nom. nov. for *Neodioctria* Wilcox & Martin, 1941). Type-species, *Dioctria* (*Neodioctria*) *albicornis* Wilcox & Martin, 1941 (aut.).

australis (Adisoemarto & Wood), 1975: 515 (*Dioctria* (*Nannodioctria*)). Type-locality: Mexico, Chiapas, 8 milles northeast of San Cristóbal. Distr. – Mexico (Chiapas, Distrito Federal). Ref. – Artigas & Papavero, 1991c: 262. Holotype, CNC.

Tribe Echthodopini Adisoemarto & Wood, 1975

Echthodopini Adisoemarto & Wood, 1975: 523. Type-genus, *Echthodopa* Loew, 1866.

Genus **Bohartia** Hull, 1958

Bohartia Hull, 1958: 317. Type-species, *Bohartia bromleyi* Hull, 1958 (orig. des.).

senecta Adisoemarto & Wood, 1975: 532. Type-locality: Mexico, Baja California, 15 miles north of Colonia Guerrero. Distr. – Mexico (Baja California). Ref. – Artigas & Papavero, 1991c: 263. Holotype, CAS.

Tribe Enigmomorphini Hull, 1962

Enigmomorphini Hull, 1962 (1): 291. Type-genus, *Enigmomorphus* Hermann, 1912.

Genus **Alyssomyia** Hull, 1962

Alyssomyia Hull, 1962 (1): 143. Type-species, *Clavator brevicornis* Philippi, 1865 (orig. des.).

Ref. – Artigas, 1973 (key to spp.).

brevicornis (Philippi), 1865: 700, fig. 3 (*Clavator*). Type-locality: Chile, Santiago. Neotype-locality: Chile, Choapa, Illapel. Distr. – Peru (Lima), Chile (Aconcagua, Biobío, Choapa, Santiago, Valparaíso). Refs. – Williston, 1891: 71 (cat.); *Hypenetes*; Kertész, 1909: 74 (cat.; *Hypenetes*); Bromley, 1932: 264 (*Hypenetes*); Stuardo Ortiz, 1946: 80 (cat.; *Pritchardia*); Hull, 1962 (1): 144, figs. 154, 480, 1013, 1022, 1830; Carrera & Papavero, 1965: 51; Artigas, 1970a: 59, figs. 42–44, 482; Martin & Papavero, 1970: 17 (cat.); Artigas, 1971: 17, fig. 31; Lamas M., 1972: 306 (cat.); Artigas & Papavero, 1991d: 30. Type, lost. NT DZUC [des. by Artigas, 1970a: 59].

bulbosa Artigas, 1970a: 61, figs. 45–48, 436. Type-locality: Chile, Atacama, Carrizal Bajo, Quebrada Algodones. Distr. – Chile (Aconcagua, Atacama, Coquimbo, Santiago, Valparaíso). Refs. – Artigas, 1971: 17, figs. 28–29; Artigas & Papavero, 1991d: 30. Type, DZUC.

misera Artigas, 1973: 125, figs. 7–11. Type-locality: Chile, Coquimbo, Baños del Toro, 3400–4000 m. Distr. – Chile (Coquimbo). Type, Ref. – Artigas & Papavero, 1991e: 30. DZUC.

pampina Artigas, 1970a: 63, figs. 49, 473. Type-locality: Chile, Antofagasta, Cordillera de Antofagasta, Talabre (Toconao), Chinina. Distr. – Chile (Antofagasta, Atacama). Refs. – Artigas, 1971: 18, fig. 30; Artigas & Papavero, 1991d: 30. Type, DZUC.

quinquemaculata Artigas, 1973: 124, figs. 1–5. Type-locality: Chile, Santiago, La Ollita, Cantillana, Piche-Alhué. Distr. – Chile (Colchagua, Santiago). Ref. – Artigas & Papavero, 1991d: 30. Type, DZUC.

Genus **Araujoa** Arrtigas & Papavero, 1991

Araujoa Artigas & Papavero, 1991d: 30. Type-species, *Araujoa pernambucana* Artigas & Papavero, 1991 (orig. des.).

pernambucana Artigas & Papavero, 1991d: 30, figs. 3–10. Type-locality: Brazil, Pernambuco, 10–20 km northwest of Petrolina. Distr. – Brazil (Pernambuco). Type, MZUSP.

Genus **Archilestris** Loew, 1874

Archilestes Schiner, 1866: 672 (1868: 136, second erection of genus) (preoc. Selys-Longchamps, 1862). Type-species, *Dasypogon capnopterus* Wiedemann, 1828 (sub. des., Schiner, 1868: 168).

Archilestris Loew, 1874a: 377 (*nom. nov.* for *Archilestes* Schiner, 1866). Type-species, *Dasypogon capnopterus* Wiedemann, 1828 (aut.).
Pseudarchilestes Bigot, 1889: clxxxiii (unnecessary *nom. nov.* for *Archilestes* Schiner, 1866). Type-species, *Dasypogon capnopterus* Wiedemann, 1828 (aut.).
Pseudoarchilestes Bigot, 1890: 96, emend.

Ref. – Papavero & Bernardi, 1974 (key to spp.).

capnoptera (Wiedemann), 1828: 376 (*Dasypogon*). Type-locality: “Brazil”. Distr. – Brazil (Mato Grosso, Goiás, Minas Gerais, Espírito Santo, Rio de Janeiro, São Paulo, Santa Catarina), Paraguay. Refs. – Walker, 1854: 444 (*Dasypogon*); Schiner, 1866: 677 (*Archilestes*), 1867: 377 (*Archilestes*); Lynch Arribálzaga, 1880: 28, 1882: 132, 1883a: 142 (cat.); Williston, 1889: 259; Kertész, 1909: 156 (cat.); Carrera, 1945a: 160 (preys), 1958a: 147, 1960a: 161; Carrera & d’Andretta, 1961: 69 (prey); Hull, 1962 (1): 141, figs. 151, 473, 1015, 1024; Papavero & Bernardi, 1974: 166, figs. 1, 4-6; Artigas & Papavero, 1991d: 31; Geller-Grimm, 1999: 207 (specimen in SENCK). Lectotype NMW [des. by Papavero & Bernardi, 1974: 266].
albitarsis Macquart, 1846: 194 (1846: 66), pl. 7, fig. 7 (*Dasypogon*). Type-locality: Brazil, Minas Gerais. Refs. – Walker, 1854: 410 (*Dasypogon*); Bigot, 1889: clxxxiii (*Pseudarchilestes*), 1890: 96 (*Pseudoarchilestes*). Type, OXF.
capnopterus Bromley, 1946: 108 (cat.); Martin & Papavero, 1970: 16 (cat.), error.
excellens Enderlein, 1914b: 174. Type-locality: Ecuador, Bolívar, Balzapamba. Distr. – Ecuador, Bolivia. Refs. – Hull, 1962 (1): 141; Martin & Papavero, 1970: 16 (cat.); Papavero & Bernardi, 1974: 268, figs. 7-9; Artigas & Papavero, 1991d: 31. Type, IZPAN.
geijskesi Papavero & Bernardi, 1974: 270, figs. 10-12. Type-locality: Surinam, Brokopondo, Phedra. Distr. – Surinam. Ref. – Artigas & Papavero, 1991d: 31. Type, MZUSP.
magnifica (Walker), 1854: 427 (*Dasypogon*). Type-locality: “Mexico”. Distr. – U. S. A. (Arizona), Mexico (Chihuahua, Coahuila, Durango, San Luis Potosí, Nayarit, Jalisco, Colima, Guanajuato, Querétaro, Veracruz, Morelos, Guerrero, Oaxaca). Refs. – Bellardi, 1861: 79, pl. 1, fig. 11 (*Dasypogon*); Schiner, 1868: 168 (*Archilestes*); Loew, 1872a: 68 (*Microstylum*; note); Osten Sacken, 1887: 169; Bigot, 1889: clxxxiii (*Archilestes*); Williston, 1889: 258, 1891: 71 (cat.); Aldrich, 1905: 256 (cat.); Back, 1909: 216; Kertész, 1909: 157 (cat.); Wilcox, 1936b: 204; Hull, 1962 (1): 141, figs. 118, 1826, 1851; Martin, 1965a: 18; Martin & Wilcox, 1965: 364 (cat.); Martin, 1968: 90, fig. 41; Martin & Papavero, 1970: 16 (cat.); Papavero & Bernardi, 1974: 271, figs. 3, 13-15; Artigas & Papavero, 1991d: 3; Papavero & Ibáñez Bernal, 2001: 95 (hist. Mexican dipterol.). Type, BMNH.
wenzeli Papavero & Bernardi, 1974: 273, figs. 2, 16-18. Type-locality: Guatemala, Chimaltenango, Yepocapa (Finca Panajabel). Distr. – Guatemala. Ref. – Artigas & Papavero, 1991d: 31. Type, FMNH.

Genus **Archilestroides** Artigas & Papavero, 1991

Archilestroides Artigas & Papavero, 1991d: 31. Type-species, *Archilestroides guimaraesi* Artigas & Papavero, 1991 (orig. des.).

guimaraesi Artigas & Papavero, 1991d: 32, figs. 24-31. Type-locality: Brazil, São Paulo, Salesópolis (Estação Biológica de Boracéia). Distr. – Brazil (Rio de Janeiro, São Paulo). Ref. – Castillo, Jerez & Artigas, 1994 (egg chorion). Type, MZUSP.

Genus **Aymarasilus** Artigas, 1974

Aymarasilus Artigas, 1974a: 227. Type-species, *Aymarasilus inti* Artigas, 1974 (orig. des.).

inti Artigas, 1974a: 228, figs. 1-10. Type-locality: Chile, Parinacota, Putre. Distr. – Chile (Parinacota). Ref. – Artigas & Papavero, 1991d: 32. Type, DZUC.

Genus **Creolestes** Hull, 1962

Bathypogon, subg. **Creolestes** Hull, 1962 (1): 150. Type-species, *Bathypogon hirtuosus* Schiner, 1868 (orig. des.) = *Creolestes nigribarbis* (Philippi, 1865).

keiseri (Carrera & Papavero), 1965: 42 (*Alyssomyia*). Type-locality: Chile, Valdivia, Panguipulli. Distr. – Chile (Arauco,

- Cautín, Concepción, Linares, Llanquihue, Malleco, Ñuble, O'Higgins, Osorno, Valdivia). Refs. – Artigas, 1971: 19, figs. 34-35; Artigas & Papavero, 1991d: 33. Type, MZUSP.
- keisseri* Artigas, 1970a: 79, figs. 78, 470, error.
- nigribarbis** (Philippi), 1865: 699, fig. 2 (*Clavator*). Type-locality: "Chile". Distr. – Chile (Coquimbo, Curicó, Maule, Ñuble, O'Higgins, Santiago, Valparaíso). Refs. – Williston, 1891: 71 (cat.; *Hypenetes*); Kertész, 1909: 74 (cat.; *Hypenetes*); Bromley, 1932: 264 (*Hypenetes*); Stuardo Ortiz, 1946: 80 (cat.; *Pritchardia*); Hull, 1962 (1): 154 (*Hypenetes*); Carrera & Papavero, 1965: 51 (*Alyssomyia*); Martin & Papavero, 1970: 18 (n. comb.); Artigas, 1971: 19, figs. 32-33; Artigas & Papavero, 1991d: 33. Type, MNHNS.
- hirtuosus* Schiner, 1868: 160 (*Bathypogon*). Type-locality: "Chile". Refs. – Williston, 1891: 72 (cat.; *Bathypogon*); Kertész, 1909: 82 (cat.; *Bathypogon*); Stuardo Ortiz, 1946: 81 (cat.; *Bathypogon*); Hull, 1962 (1): 150, figs. 512, 1019, 1028 (*Bathypogon (Creolestes)*). Syntypes, NMW.
- negribarbis* Artigas, 1970a: 82, figs. 54-57, 59-60, 62, 497 (n. comb.), error.
- parvus** (Bigot), 1878: 432 (*Bathypogon*). Type-locality: "Chile". Distr. – Chile. Refs. – Williston, 1891: 72 (cat.; *Bathypogon*); Kertész, 1909: 82 (cat.; *Bathypogon*); Stuardo Ortiz, 1946: 81 (cat.; *Bathypogon*); Hull, 1962: 150 (*Bathypogon*); Artigas, 1970a: 85, fig. 61; Artigas & Papavero, 1991d: 33. Type, OXF.
- rubricornis** (Philippi), 1865: 700 (*Clavator*). Type-locality: "Chile". Neotype-locality: Chile, Angol, Cerros de Nahuelbuta. Distr. – Chile (Malleco, Maule, Ñuble). Refs. – Williston, 1891: 71 (cat.; *Hypenetes*); Kertész, 1909: 74 (cat.; *Hypenetes*); Bromley, 1932: 264 (*Hypenetes*); Stuardo Ortiz, 1946: 80 (cat.; *Pritchardia*); Hull, 1962 (1): 154 (*Hypenetes*); Carrera & Papavero, 1965: 51 (*Alyssomyia*); Artigas, 1970a: 89, figs. 50, 448; Martin & Papavero, 1970: 18 (cat.; n. comb.); Artigas, 1971: 18, fig. 36; Artigas & Papavero, 1991d: 33. Type, lost. NT DZUC [des. by Artigas, 1970a: 89].
- rufescens* (Philippi), 1865: 700 (*Clavator*). Type-locality: "Chile" [Valparaíso in ST labels]. Neotype-locality: Chile, Santiago, Manquehue. Distr. – Chile (Santiago, Valparaíso). Refs. – Williston, 1891: 71 (cat.; *Hypenetes*); Kertész, 1909: 74 (cat.; *Hypenetes*); Bromley, 1932: 264 (*Hypenetes*); Stuardo Ortiz, 1946: 80 (cat.; *Pritchardia*); Hull, 1962 (1): 154 (*Hypenetes*); Artigas, 1970a: 86, figs. 51-53, 498; Martin & Papavero, 1970: 18 (cat.; n. comb.); Artigas & Papavero, 1991d: 33. Type, lost. NT DZUC [des. by Artigas, 1970a: 86].
- claripennis* Schiner, 1868: 160 (*Bathypogon*). Type-locality: "Chile". Type, NMW. Refs. – Williston, 1891: 71 (cat.; *Bathypogon*); Kertész, 1909: 82 (cat.; *Bathypogon*); Martin & Papavero, 1970: 18 (cat.; syn.). Type, NMW.

Genus **Cylicomera** Lynch Arribálzaga, 1881

Cylicomera Lynch Arribálzaga, 1881: 115. Type-species, *Cylicomera rubrofasciata* Lynch Arribálzaga, 1881 (sub. des., Hull, 1962 (1): 135).

Ref. – Lamas M., 1973a (key to spp.).

- dissona** Lamas M., 1973a: 28, figs. 17, 25, 40, 60, 98-100. Type-locality: Argentina, Tucumán, San Pedro de Colalao. Distr. – Argentina (Tucumán, Buenos Aires, Santiago del Estero). Ref. – Artigas & Papavero, 1991d: 33. Type, MZUSP.
- rubrofasciata** Lynch Arribálzaga, 1881: 119. Type-locality: Argentina, Buenos Aires, Chascomús and Chacabuco. Distr. – Brazil (Tocantins, Rio Grande do Sul), Paraguay, Argentina. Refs. – Lynch Arribálzaga, 1883a: 141 (cat.); Williston, 1891: 70 (cat.); Kertész, 1909: 148 (cat.); Hull, 1962 (1): 137, figs. 478, 1034, 1043; Martin & Papavero, 1970: 15 (cat.; *Cylicomera*); Lamas M., 1973a: 25; Artigas & Papavero, 1991c: 33. Lectotype MACN [des. by Lamas M., 1973a: 25].
- fraterna** Lynch Arribálzaga, 1881: 117. Type-locality: Argentina, Buenos Aires, Chacabuco. Distr. – Argentina (Buenos Aires). Refs. – Lynch Arribálzaga, 1883a: 141 (cat.); Williston, 1891: 70 (cat.); Kertész, 1909: 148 (cat.); Hull, 1962 (1): 137; Martin & Papavero, 1970: 15 (cat.; *Cylicomera*); Artigas & Papavero, 1991d: 33. Lectotype MACN [des. by Lamas M., 1973: 25].

Genus **Dicranus** Loew, 1851

Microstylum, subg. **Dicranus** Loew, 1851: 13. Type-species, *Dasypogon rutilus* Wiedemann, 1821 (orig. des.).
Macronix Bigot, 1857: 549 (in key). Type-species, *Dasypogon longiungulatus* Macquart, 1850 (mon.) = *Dicranus schtottkyi* Bezzii, 1910.

Ref. – Carrera, 1955b (key to spp.).

- jaliscoensis** Williston, 1901: 302, pl. 5, fig. 15. Type-locality: Mexico, Nayarit, San Blás; Santiago Izcuintla; Guadalajara, Jalisco. Distr. – Mexico (Jalisco, Nayarit). Refs. – Aldrich, 1905: 255 (cat.); Kertész, 1909: 155 (cat.); Carrera, 1955b:

- 244; Hull, 1962 (1): 143, figs. 163, 948, 957, 1696, 1826; Martin & Papavero, 1970: 17 (cat.); Artigas & Papavero, 1991d: 33; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipteral). Syntypes, BMNH.
- nigerrimus** Carrera, 1955b: 245. Type-locality: Argentina, [?], Estancia Siete de Abril. Distr. – Argentina. Refs. – Hull, 1962 (1): 143; Artigas & Papavero, 1991d: 33. Type, IML.
- rutilus** (Wiedemann), 1821: 231 (*Dasypogon*). Type-locality: “Brazil” [Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-102]. Distr. – Brazil (Mato Grosso and Goiás to Santa Catarina). Refs. – Wiedemann, 1828: 370 (*Dasypogon*); Loew, 1851: 23; Walker, 1854: 445 (*Dasypogon*); Schiner, 1866: 676; Williston, 1891: 70 (cat.); Kertész, 1909: 155 (cat.); Bromley, 1946: 108 (cat.); Carrera, 1955b: 236, fig. 1-5; Hull, 1962 (1): 143; Martin & Papavero, 1970: 17 (cat.); Artigas & Papavero, 1991d: 33. Syntypes, BZM [no. 489; cf. Schumann, 1973: 154].
- schröttkyi** Bezzi, 1910a: 67 (nom. nov. for *longiungulatus* Macquart, 1850). Type-locality: Bolivia, Yungas. Distr. – Venezuela (Aragua), Peru, Bolivia, Brazil (Amazonas, Goiás, Santa Catarina), Paraguay, Argentina (Misiones). Refs. – Carrera, 1947a: 43, 1955b: 242, 1959: 2; Hull, 1962 (1): 143, figs. 476, 1934; Carrera & Machado-Allison, 1963: 238; Lamas M., 1972: 306 (cat.); Fisher, 1984: 27; Artigas & Papavero, 1991d: 33.
- longiungulatus* Macquart, 1850: 371 (1850: 67), pl. 6, fig. 14 (*Dasypogon*; preocc. Macquart, 1838). Type-locality: Bolivia, Yungas (as Brazil, error). Refs. – Bigot, 1857b: 549 (*Macronix*); Schiner, 1866: 676; Williston, 1891: 70 (cat.); Kertész, 1909: 155 (cat.); Bromley, 1946: 108 (cat.).
- tucma** Lynch Arribálzaga, 1880: 26. Type-locality: Argentina, Tucumán. Distr. – Argentina (Córdoba, Jujuy, Salta, Tucumán). Refs. – Lynch Arribálzaga, 1881: 124; Wulp, 1882b: 98; Lynch Arribálzaga, 1883a: 142 (cat.); Williston, 1891: 70 (cat.); Kertész, 1909: 156 (cat.); Carrera, 1955b: 240; Hull, 1962 (1): 143; Martin & Papavero, 1970: 17 (cat.); Artigas & Papavero, 1991d: 33.

Genus **Enigmomorphus** Hermann, 1912

- Enigmomorphus** Hermann, 1912: 272. Type-species, *Enigmomorphus paradoxus* Hermann, 1912 (orig. des.).
Enigmomorpheus Hull, 1962 (1): 291, error.

- paradoxus** Hermann, 1912: 274, fig. 87, pl. 5, figs. 64-65. Type-locality: Paraguay, Asunción. Distr. – Brazil (São Paulo), Paraguay. Refs. – Hull, 1962 (1): 292, figs. 107, 567, 1159, 1168, 1607, 1920, 1990; Artigas & Papavero, 1991d: 33. Type, BUD.

Genus **Leptochelina** Artigas, 1970

- Leptochelina** Artigas, 1970a: 143. Type-species, *Leptochelina jaujensis* Artigas, 1970 (orig. des.).

- jaujensis** Artigas, 1970a: 144, figs. 128, 130-132, 426. Type-locality: Chile, Tarapacá, Jauja, Mamiña, Iquique. Distr. – Chile (Tarapacá). Ref. – Artigas & Papavero, 1991d: 34. Type, DZUC.

Genus **Microstylum** Macquart, 1838

- Microstylum** Macquart, 1838: 26 (1839: 142). Type-species, *Dasypogon venosus* Wiedemann, 1821 (sub. des., Back, 1909: 213).

Two species known as fossils: *Microstylum destructum* Cockerell (1909: 254) and *Microstylum wheeleri* Cockerell (1908: 173) (Oligocene; U. S. A., Colorado).

- catastygnum** Papavero, 1971d: 25, figs. 4-6. Type-locality: French Guiana, Maroni. Distr. – French Guiana. Ref. – Artigas & Papavero, 1991d: 34. Type, MNHNP.
- fulvigaster** Bigot, 1878: 410. Type-locality: “Mexico”. Distr. – Mexico. Refs. – Williston, 1883: 33; Aldrich, 1905: 256 (cat.); Kertész, 1909: 152 (cat.); Hull, 1962 (1): 141 (*Archilestris*), 157 [*Microstylum*]; Artigas & Papavero, 1991d: 34; Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipteral). Type, OXF.
- galactodes** Loew, 1866: 25 (Cent. 7, no. 44). Type-locality: U. S. A., New Mexico. Distr. – U. S. A. (New Mexico, Arizona, Kansas, Texas, Oklahoma, Colorado), Mexico (Coahuila). Refs. – Williston, 1883: pl. 1, fig. 7; Aldrich, 1905: 256 (cat.); Kertész, 1909: 152 (cat.); Back, 1909: 213; Hine, 1918: 3; Bromley, 1935c: 3; Baker, 1939: 168; Carrera & d’Andretta, 1950: 167; Hull, 1962 (1): 157; Martin & Wilcox, 1965: 379 (cat.). Type, MCZ.
- mexicanum** Martin, 1960: 44. Type-locality: Mexico, Nuevo León, Monterrey. Distr. – Mexico (Nuevo León). Ref. – Artigas & Papavero, 1991d: 34. Type,

morusum Loew, 1872a: 67 (Cent. 10, no. 27). Type-locality: U. S. A., Texas. Distr. – U. S. A. (Texas, Arizona, New Mexico, Oklahoma, Kansas, Colorado), Mexico (Coahuila, Tamaulipas). Refs. – Baker, 1939: 168; Hull, 1962 (1): 157; Martin & Wilcox, 1965: 379 (cat.). Type, MCZ.
pollens Osten Sacken, 1878: 230 (cat.; note 100). Type-locality: U. S. A., Texas. Ref. – Hull, 1962 (1): 157.

Genus **Pritchardia** Stuardo Ortiz, 1946

Pritchardia Stuardo Ortiz, 1946: 80. Type-species, *Dasypogon hirtipes* Macquart, 1838 (orig. des.).
Strobilopygius Hull, 1956: 70. Type-species, *Dasypogon hirtipes* Macquart, 1838 (orig. des.).

curicoensis Artigas, 1970a: 157, fig. 464. Type-locality: Chile, Curicó, La Jaula. Distr. – Chile (Curicó). Refs. – Artigas, 1971: 27, fig. 58; Artigas & Papavero, 1991d: 34. Type, DZUC.

hirtipes (Macquart), 1838: 46 (1839: 162) (*Dasypogon*). Type-locality "Chile". Distr. – Chile Arauco, Cautín, Chiloé, Concepción, Llanquihue, Maule, Malleco, Ñuble, Santiago, Talca, Valdivia, Valparaíso), Argentina (Tucumán, Neuquén, Chubut). Refs. – Blanchard, 1852: 368 (*Dasypogon*); Walker, 1854: 438 (*Dasypogon*); Philippi, 1865: 686 (*Dasypogon*); Williston, 1891: 71 (cat.; *Bathypogon*); Kertész, 1909: 82 (cat.; *Bathypogon*); Bromley, 1932: 265 (*Hypenetes*); Stuardo Ortiz, 1946: 80 (cat.; n. comb.); Hull, 1956: 72 (*Strobilopygius*), 1962 (1): 36, figs. 22, 422, 809, 818, 1659, 1909, 2002; Carrera & Papavero, 1965: 49; Artigas, 1970a: 159, figs. 16-23, 412, 1971: 27, figs. 61-62; Artigas & Papavero, 1991d: 34. Type, MNHNP.

lopesi Carrera & Papavero, 1965: 49, figs. 1-2. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. – Brazil (Rio de Janeiro, São Paulo). Ref. – Artigas & Papavero, 1991d: 34. Type, MZUSP.

puella (Bromley), 1932: 264 (*Hypenetes*). Type-locality: Chile, Llanquihue, Peulla. Distr. – Chile (Coquimbo, Valparaíso), Argentina (Neuquén). Refs. – Stuardo Ortiz, 1946: 80 (cat.; n. comb.); Hull, 1962 (1): 36, figs. 45, 1020, 1029, 1884; Carrera & Papavero, 1965: 49; Artigas, 1970a: 163, fig. 472; Artigas & Papavero, 1991d: 34. Type, BMNH.

tertialis (Bromley), 1932: 265 (*Hypenetes*). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. – Chile (Arauco, Llanquihue, Malleco), Argentina (Río Negro, Neuquén). Refs. – Stuardo Ortiz, 1946: 80 (n. comb.); Hull, 1962 (1): 154 (*Hypenetes*); Carrera & Papavero, 1965: 49; Artigas, 1970a: 161, figs. 24-27, 481, 1971: 28, figs. 59-60; Artigas & Papavero, 1991d: 34. Type, BMNH.

Genus **Prolepsis** Walker, 1851

Prolepsis Walker, 1851: 101. Type-species, *Prolepsis fumiflamma* Walker, 1851 (mon.) = *Prolepsis lucifer* (Wiedemann, 1828).
Cacodaemon Schiner, 1866: 671, 678, 702 (preocc. Thomson, 1857). Type-species, *Dasypogon lucifer* Wiedemann, 1828 (orig. des.).

Dizonias Loew, 1861: 29. Type-species, *Dizonias phoenicurus* Loew, 1866 (sub. des., Coquillett, 1910: 534) = *Prolepsis tristis* (Walker, 1851). Ref. – Lamas M., 1973a: 30 (syn.).

Sphageus Loew, 1861: 32. Type-species, *Sphageus chalcoproctus* Loew, 1866 (mon.). Ref. – Lamas M., 1973a: 30 (syn.).
Tolmerolestes Lynch Arribálzaga, 1881: 27. Type-species, *Tolmerolestes fax* Lynch Arribálzaga, 1881 (sub. des., Kirby, 1882: 243, as *lax*, error). Ref. – Lamas M., 1973a: 30 (syn.).

Cacodaemonides Strand, 1927: 48 (unnecessary nom. nov. for *Cacodaemon* Schiner, 1866). Type-species, *Dasypogon lucifer* Wiedemann, 1828 (aut.).

Ref. – Lamas M., 1973a (rev.).

chalcoprocta (Loew), 1866: 32 (*Sphageus*). Type-locality: Cuba, La Habana, Playa del Chivo. Distr. – Cuba. Refs. – Aldrich, 1905: 256 (cat.; *Sphageus*); Kertész, 1909: 82 (cat.; *Sphageus*); Bromley, 1929a: 277 (*Sphageus*); Hull, 1962 (1): 137, figs. 117, 474, 912, 921, 1576A-B, 1740, 1901, 2051, 2055 (*Sphageus*); Martin & Papavero, 1970: 15 (cat.; *Sphageus*); Lamas M., 1973a: 53, figs. 30, 48, 68-70; Alayo Dalmau & García Ávila, 1983: 45 (cat.; *Sphageus*); Artigas & Papavero, 1991d: 34. Type, BZM [no. 6491; cf. Schumann, 1973: 96].

colalao Lamas M., 1973a: 41, figs. 19-20, 36, 54, 74-76. Type-locality: Argentina, Tucumán, San Pedro Colalao. Distr. – Argentina (Catamarca, Jujuy, Mendoza, Salta, Tucumán). Ref. – Artigas & Papavero, 1991d: 34. Type, MZUSP.

costaricensis Lamas M., 1973a: 52, figs. 12, 23, 31, 49. Type-locality: Costa Rica, Liberia, 400 ft. Distr. – Costa Rica. Ref. – Artigas & Papavero, 1991d: 34. Type, CAS.

crabroniformis (Schiner), 1866: 702 (*nomen nudum*), 1867: 375 (*Cacodaemon*). Type-locality: unknown. Distr. – Brazil (Paraná), Argentina (Córdoba, Santa Fe). Refs. – Williston, 1891: 70 (cat.); Kertész, 1909: 147 (cat.); Hull, 1962 (1): 139; Martin & Papavero, 1970: 16 (cat.); Lamas M., 1973a: 55, figs. 5, 22, 28, 46; Artigas & Papavero, 1991d: 34. Type, NMW.

- elotensis** (Martin), 1966: 216, figs. 6, 8 (*Sphageus*). Type-locality: Mexico, Sinaloa, Elota. Distr. – Mexico (Sinaloa, Sonora). Refs. – Martin & Papavero, 1970: 15 (cat.; *Sphageus*); Lamas M., 1973a: 55, figs. 18, 20, 47, 65–67; Artigas & Papavero, 1991d: 34. Type, CAS.
- fax** (Lynch Arribálzaga), 1881: 30 (*Tolmerolestes*). Type-locality: Argentina, Buenos Aires, Baradero. Distr. – Argentina (San Luís, Córdoba, Buenos Aires). Refs. – Lynch Arribálzaga, 1883a: 141 (cat.; *Tolmerolestes*); Williston, 1891: 70 (cat.; *Tolmerolestes*); Hull, 1962 (1): 135 (*Tolmerolestes*); Martin & Papavero, 1970: 15 (cat.; *Tolmerolestes*); Lamas M., 1973a: 36, figs. 14, 39, 55, 77–70; Artigas & Papavero, 1991d: 34. Type, lost.
- lax* Kirby, 1882: 243 (*Tolmerolestes*), error.
- brethesi* Gemignani, 1936: 43, pl. 4 (*Tolmerolestes*). Type-locality: Argentina, San Luís, Potrero de los Funes. Refs. – Hull, 1962 (1): 135 (*Tolmerolestes*); Martin & Papavero, 1970: 15 (cat.; *Tolmerolestes*). Type, MACN.
- fenestrata** (Macquart), 1838: 53 (1839: 169) (*Senobasis*). Type-locality: Brazil, western Minas Gerais. Distr. – Brazil (Goiás, Minas Gerais, São Paulo), Paraguay (Guaira), Argentina (Chacxo, Misiones). Refs. – Walker, 1854: 500 (*Senobasis*); Williston, 1891: 68 (*Dasytopogon*); Kertész, 1909: 124 (cat.; *Stenobasis*); Bromley, 1946: 106 (cat.; *Senobasis*); Hull, 1962 (1): 233 (*Blepharepium*); Lamas M., 1973a: 46, figs. 9–10, 41, 51, 89–91; Artigas & Papavero, 1991d: 35. Type, MNHNP.
- funebris** Lamas M., 1973a: 58, figs. 4, 26, 44. Type-locality: Brazil, Goiás, Jataí (Fazenda Aceiro). Distr. – Brazil (Goiás). Ref. – Artigas & Papavero, 1991d: 35. Type, MZUSP.
- huatayata** Lamas M., 1972: 306. Type-locality: Bolivia, La Paz, Huatajata, lago Titicaca. Distr. – Peru (Cuzco, Puno), Bolivia (La Paz). Ref. – Artigas & Papavero, 1991d: 35. Type, MZUSP.
- huatajata* Lamas M., 1973a: 33, figs. 13, 24, 42, 53, 80–92, error.
- indecisa** Lamas M., 1973a: 31, figs. 7, 38, 56, 71–73. Type-locality: Argentina, Tucumán, Tucumán. Distr. – Argentina (Tucumán, Catamarca, La Rioja, Jujuy, Santiago del Estero). Ref. – Artigas & Papavero, 1991d: 35. Type, MZUSP.
- lucifer** (Wiedemann), 1828: 388 (*Dasytopogon*). Type-locality: Uruguay, Montevideo. Distr. – Brazil (Rio Grande do Sul), Argentina (Buenos Aires, Catamarca, Córdoba, Chubut, La Pampa, Mendoza, Neuquén, Río Negro, San Luís, Santa Fé, Santiago del Estero, Tucumán), Uruguay (Colonia, Lavalleja, San José); ?Chile. Refs. – Walker, 1854: 432 (*Dasytopogon*); Schiner, 1866: 678 (*Cacodaemon*); Lynch Arribálzaga, 1879: 152 (*Cacodaemon*), 1881: 26; Wulp, 1882b: 97; Lynch Arribálzaga, 1883a: 140 (cat.); Williston, 1891: 70 (cat.); Kertész, 1909: 147 (cat.); Bromley, 1946: 108 (cat.); Carrera, 1950b: 86, figs. 1–4, 1953a: 277; Carrera & d'Andretta, 1961: 69 (prey); Hull, 1962 (1): 139, figs. 187, 459, 919, 1763, 1766, 1947; Martin & Papavero, 1970: 16 (cat.); Lamas M., 1973a: 58, figs. 3, 27, 45, 62–64; Artigas & Papavero, 1991d: 35; Soria & Mello, 1998: 279 (biol., larva, pupa); Soria, Mello & Oliveira, 2004: 323 (larvae preying upon nymphs of *Eurhizococcus brasiliensis* (Hempel, 1922) (Hemiptera, Margarodidae)); Foldi, 2005: 120 (brief mention as predator of *Eurhizococcus brasiliensis*); Soria & Dal Conte, 2005: 3 (brief mention as predator of *Eurhizococcus brasiliensis*). Type, BZM.
- satanas* Wiedemann, 1828: 401 (*Dasytopogon*). Type-locality: Uruguay, Montevideo. Refs. – Loew, 1851: 13 (*Microstylum*); Walker, 1854: 442 (*Dasytopogon*); Schiner, 1866: 679 (*Cacodaemon*); Bigot, 1866: 320 (*Dasytopogon*). Type, BZM.
- rufipennis* Macquart, 1838: 45 (1839: 161) (*Dasytopogon*). Type-locality: Uruguay, “depuis de l’embouchure de l’Uruguay jusqu’aux Missions”. Ref. – Walker, 1854: 438 (*Dasytopogon*); Lynch Arribálzaga, 1880: 29 (*Dasytopogon*). Type, MNHNP.
- fumiflamma* Walker, 1851: 101, pl. 3, figs. 6, 6a. Type-locality: “Brazil”. Refs. – Walker, 1854: 437. Type, BMNH.
- quadrinotatum* Bigot, 1878: 431 (*Cacodaemon*). Type-locality: “Chile”. Refs. – Williston, 1891: 70 (cat.); Kertész, 1909: 148 (cat.); Stuardo Ortiz, 1946: 83 (cat.); Hull, 1962: 139; Artigas, 1970a: 266, figs. 142–147, 412. Type OXF.
- martini** Lamas M., 1973a: 34, figs. 33, 59. Type-locality: Argentina, Córdoba, Villa de María. Distr. – Argentina (Córdoba). Ref. – Artigas & Papavero, 1991d: 35. Type, MZUSP.
- ?pluto** (Lynch Arribálzaga), 1881: 112 (*Tolmerolestes*). Type-locality: Argentina, Buenos Aires, Baradero. Distr. – Brazil (Rio Grande do Sul), Argentina (Buenos Aires, Catamarca, Córdoba, Jujuy, La Rioja, Salta, San Luís, Tucumán). Refs. – Lynch Arribálzaga, 1883a: 141 (cat.; *Tolmerolestes*); Williston, 1891: 70 (cat.; *Tolmerolestes*); Kertész, 1909: 148 (cat.; *Tolmerolestes*); Carrera, 1959: 2 (*Tolmerolestes*); Hull, 1962 (1): 135, figs. 472, 913, 922 (*Tolmerolestes*); Martin & Papavero, 1970: 15 (cat.; *Tolmerolestes*); Lamas M., 1973a: 38, figs. 15, 37, 57 (may be female of *fax* Lynch Arribálzaga, 1881); Artigas & Papavero, 1991d: 35. Lectotype MACN.
- rubripes* Lynch Arribálzaga, 1881: 114 (*Tolmerolestes*). Type-locality: Argentina, San Luís. Refs. – Lynch Arribálzaga, 1883a: 141 (cat.; *Tolmerolestes*); Williston, 1891: 70 (cat.; *Tolmerolestes*); Kertész, 1909: 148 (cat.; *Tolmerolestes*); Hull, 1962 (1): 134, fig. 113 (*Tolmerolestes*); Martin & Papavero, 1970: 15 (cat.; *Tolmerolestes*). Type, MACN.
- pseudopluto** Lamas M., 1973a: 40, figs. 16, 32, 61. Type-locality: Argentina, Tucumán, El Infernillo. Distr. – Argentina (Tucumán). Ref. – Artigas & Papavero, 1991d: 35. Type, MZUSP.
- rosariana** (Carrera), 1959: 2, fig. 1 (*Tolmerolestes*). Type-locality: Argentina, Rosario. Distr. – Brazil (São Paulo, Paraná, Santa Catarina), Argentina (Córdoba, Santa Fe). Refs. – Martin & Papavero, 1970: 15 (cat.; *Tolmerolestes*); Lamas M., 1973a: 43, figs. 6, 11, 43, 52, 92–94; Artigas & Papavero, 1991d: 35. Type, ZSBS.
- ?sandaraca** (Martin), 1966: 217, fig. 7 (*Sphageus*). Type-locality: Mexico, Sinaloa, Elota. Distr. – Mexico (Sinaloa). Refs. – Martin & Papavero, 1970: 15 (cat.; *Sphageus*); Lamas M., 1973a: 55 (may be female of *elotensis* Martin, 1966); Artigas & Papavero, 1991d: 35. Type, FSCA.
- tristis** (Walker), 1851: 93 (*Dasytopogon*). Type-locality: “U. S. A.”. Distr. – U. S. A., Mexico (Colima, Guerrero, Jalisco, Michoacán, Morelos, Nayarit), Belize. Refs. – Walker, 1854: 423 (*Dasytopogon*); Osten Sacken, 1874: 185 (*Dizonias*);

Williston, 1891: 69 (cat.; *Dizonias*); Johnson, 1903: 112, fig. 4 (*Dizonias*); Aldrich, 1905: 257 (cat.; *Dizonias*); Kertész, 1909: 148 (cat.; *Dizonias*); Hine, 1922: 3 (*Dizonias*); Fattig, 1945: 16; Bromley, 1950: 231 (*Dizonias*); Hull, 1962 (1): 134, fig. 1927 (*Dizonias*); Martin, 1965a: 23 (*Dizonias*); Martin & Wilcox, 1965: 372 (cat.; *Dizonias*); Martin & Papavero, 1970: 14 (cat.; *Dizonias*); Lamas M., 1973a: 47, figs. 8, 35, 50, 83-88; Artigas & Papavero, 1991d: 35. Type, BMNH.

quadrimaculatus Bellardi, 1861: 180 (80), pl. 1, fig. 8 (*Dasypogon*). Type-locality: "Mexico". Refs. – Loew, 1866: 31 (*Dizonias*; note), 1872: 68 (*Dizonias*, note); Osten Sacken, 1874: 180 (*Dizonias*); Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipterol). Type, TORO.

lucasi Bellardi, 1861: 181 (81), pl. 1, fig. 7 (*Dasypogon*). Type-locality: "Mexico". Refs. – Loew, 1872: 68 (*Dizonias*; note); Osten Sacken, 1874: 180 (*Dizonias*); Williston, 1901: 304 (*Dizonias*); Aldrich, 1905: 257 (cat.; *Dizonias*); Kertész, 1909: 148 (cat.; *Dizonias*); Hull, 1962 (1): 134, figs. 1770, 1773 (*Dizonias*); Martin & Papavero, 1970: 14 (cat.; *Dizonias*); Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipterol). Type, TORO.

phoenicurus Loew, 1866: 29 (*Dizonias*). Type-locality: Mexico, Tamaulipas. Refs. – Osten Sacken, 1874: 180 (*Dizonias*); Williston, 1901: 304 (*Dizonias*); Aldrich, 1905: 257 (cat.; *Dizonias*); Kertész, 1909: 148 (cat.; *Dizonias*); Hull, 1962 (1): 134, figs. 110, 471, 951, 960 (*Dizonias*). Type, MCZ.

bicinctus Loew, 1866: 30 (*Dizonias*). Type-locality: U. S. A., New Mexico. Refs. – Osten Sacken, 1874: 180 (*Dizonias*), 1878: 68, 230 (cat.; *Dizonias*); Wulp, 1882: 96 (*Dizonias*); Martin & Wilcox, 1965: 372 (cat.; *Dizonias*). Type, MCZ.

pilatei Johnson, 1903: 112, fig. 5 (*Dizonias*). Type-locality: U. S. A., New Mexico, Tifton. Refs. – Aldrich, 1905: 257 (cat.; *Dizonias*); Kertész, 1909: 148 (cat.; *Dizonias*); Fattig, 1945: 15 (*Dizonias*); Hull, 1962 (1): 134 (*Dizonias*); Martin & Wilcox, 1965: 372 (cat.; *Dizonias*).

albifasciatus Back, 1904: 292 (*Ospriocerus*). Type-locality: U. S. A., Florida, Indian River. Type, ?MCZ.

bromleyi Carrera & d'Andretta, 1950: 164, fig. 26 (*Dizonias*). Type-locality: Mexico, Michoacán, Apatzingán. Ref. – Hull, 1962 (1): 134 (*Dizonias*). Type, FMNH.

Genus **Taperigna** Artigas & Papavero, 1991

Taperigna Artigas & Papavero, 1991d: 36. Type-species, *Taperigna diogmitiformis* Artigas & Papavero, 1991 (orig. des.).

diogmitiformis Artigas & Papavero, 1991d: 36, figs. 15-18. Type-locality: Brazil, Pará, Santarém (Fazenda Taperinha). Distr. – Brazil (Amazonas, Pará). Ref. – Castillo, Jerez & Artrigas, 1994 (egg chorion). Type, MZUSP.

Tribe **Phellini** Hardy, 1926

Phellini G. H. Hardy, 1926: 306, 307, 309. Type-genus, *Phellus* Walker, 1851.
Phelini Papavero, 1973a: 345, error.

Genus **Obelophorus** Schiner, 1868

Obelophorus Schiner, 1866: 672 (1868: 167; second erection of genus). Type-species, *Dasypogon terebratus* Macquart, 1850 (orig. des.).

Refs. – Artigas, 1970a: 146-158, figs. 28-33, 476, 1971: figs. 23, 25.

landbecki (Philippi), 1865: 686, pl. 25, fig. 24 (*Dasypogon*). Type-locality: Chile, Valdivia, Colchagua. Neotype-locality: "Chile". Distr. – Chile (Arauco, Concepción, Maule, Ñuble, O'Higgins, Valdivia). Refs. – Schiner, 1866: 672; Williston, 1891: 70 (cat.); Kertész, 1909: 156 (cat.); Bromley, 1932: 46, figs. 28, 805, 814, 1654, 1877; Hull, 1962 (1): 46, figs. 28, 805, 814, 1654, 1877; Artigas, 1970a: 147, fig. 485; Martin & Papavero, 1970: 11 (cat.); Artigas, 1971: 15, fig. 25; Artigas & Papavero, 1993: 310. Type, lost. NT MHNHS [des. by Artigas, 1970a: 147].

terebratus (Macquart), 1850: 370 (1850: 66), pl. 6, fig. 11 (*Dasypogon*). Type-locality: "Chile". Distr. – Chile (O'Higgins, Santiago, Valparaíso). Refs. – Blanchard, 1852: 366 (*Dasypogon*); Philippi, 1865: 685 (*Dasypogon*); Schiner, 1866: 672, 1868: 168; Williston, 1891: 70 (cat.); Kertész, 1909: 156 (cat.); Bromley, 1932: 263; Stuardo Ortiz, 1946: 83 (cat.); Hull, 1962 (1): 46, fig. 401; Artigas, 1970a: 149, figs. 28-33, 476; Martin & Papavero, 1970: 11 (cat.); Artigas, 1971: 15, figs. 34-34; Artigas & Papavero, 1993: 310.

Tribe Plesiommatini Artigas & Papavero, 1991

Plesiommatini Artigas & Papavero, 1991a: 141, 142. Type-genus, *Plesiomma* Macquart, 1838.

Genus **Cystoprosopa** Hull, 1962

Plesiomma, subg. **Cystoprosopa** Hull, 1962 (1): 220. Type-species, *Plesiomma (Cystoprosopa) sepia* Hull, 1962 (orig. des.). Refr. – Artigas & Papavero, 1993: 310 (generic status).

semirufa (Wiedemann), 1828: 393 (*Dasypogon*). Type-locality: Uruguay. Montevideo (as Brazil, error) [Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-101]. Distr. – Uruguay. Refs. – Walker, 1854: 443 (*Dasypogon*); Schiner, 1866: 678; Wulp, 1882b: 96; Lynch Arribalzaga, 1882: 132, 1883a: 143 (cat.); Williston, 1891: 69 (cat.); Bromley, 1946: 108 (cat.; *Plesiomma*); Hull, 1962 (1): 219; Artigas & Papavero, 1993: 310 (n. comb.). Syntypes, BZM [no. 494; cf. Schumann, 1973: 157].
sepia (Hull), 1962 (1): 221 (*Plesiomma (Cystoprosopa)*). Type-locality: Mexico, Morelos, Alpuyeca. Distr. – Mexico (Morelos). Refs. – Martin & Papavero, 1970: 22 (cat.); Artigas & Papavero, 1993: 310 (n. comb.). Type, CAS.

Genus **Dapsilochetus** Hull, 1962

Plesiomma, subg. **Dapsilochetus** Hull, 1962 (1): 219. Type-species, *Plesiomma (Dapsilochetus) inflatum* Hull, 1952 (orig. des.). Ref. – Artigas & Papavero, 1993: 311 (generic status).

inflatus (Hull), 1962 (1): 219 (*Plesiomma (Dapsilochetus)*). Type-locality: Haiti, Mount Basil, 4700 ft. Distr. – Dominican Republic (La Altagracia, Montecristi, Pedernales), Haiti. Refs. – Martin & Papavero, 1970: 22 (cat.; *Plesiomma*); Artigas & Papavero, 1993: 311 (n. comb.); Scarbrough & Perez-Gelabert, 2006: 73 (in key), 76, figs. 98-104 (*Plesiomma*). Type, MCZ.

Genus **Plesiomma** Macquart, 1838

Plesiomma Macquart, 1838: 54 (1839: 170). Type-species, *Plesiomma testaceum* Macquart, 1838 (sub. des., Back, 1909: 306) = *caedens* (Wiedemann, 1828).

Dolichodes Macquart, 1838: 49 (1839: 165). Type-species, *Dolichodes ferrugineus* Macquart, 1838 (orig. des.).

Plesioma Schiner, 1866: 702, error.

Plesiomma, subg. *Boropis* Hull, 1962 (1): 219. Type-species, *Asilus lineatus* Fabricius, 1781 (orig. des.).

angustum (Macquart), 1848: 20 (1848: 180), pl. 1, figs. 11, 11a (*Dasypogon*). Type-locality: “Haiti”. Distr. – Dominican Republic (La Altagracia, Montecristi, Pedernales), Haiti. Refs. – Walker, 1854: 429 (*Dasypogon*); Osten Sacken, 1874: 185; Bigot, 1878: 221; Aldrich, 1905: 167 (*Dasypogon*); Hull, 1962 (1): 219 (n. comb.); Martin & Papavero, 1970: 21 (cat.); Artigas & Papavero, 1993: 311 (as unplaced sp.); Scarbrough & Perez-Gelabert, 2006: 73 (in key and text), figs. 91-97. Type, lost.

augusta Kertész, 1909: 157 (cat.; *Dolichodes*), error.

atrum Bromley, 1929b: 45. Type-locality: Mexico, Veracruz, Coatepec. Distr. – Mexico (Veracruz). Refs. – Hull, 1962 (1): 219; Martin & Papavero, 1970: 21 (cat.); Artigas & Papavero, 1993: 311. Type, USNM.

caedens (Wiedemann), 1828: 584 (*Dasypogon*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Distr. – Brazil, Paraguay. Refs. – Walker, 1854: 442 (*Dasypogon*); Schiner, 1866: 681; Williston, 1891: 69 (cat.); Kertész, 1909: 145 (cat.); Bromley, 1946: 108 (cat.); Carrera, 1947a: 43, 1959: 2; Hull, 1962 (1): 219; Martin & Papavero, 1970: 21 (cat.); Artigas & Papavero, 1993: 311. Type, SENCK [cf. Geller-Grimm, 1999: 207].

testaceum Macquart, 1838: 55 (1839: 171), pl. 5, fig. 2 (preocc. Fabricius, 1805). Type-locality: unknown. Refs. – Walker, 1854: 501; Kertész, 1909: 146 (cat.); Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.). Type, MHNHP.

ferrugineum (Macquart), 1838: 49 (1839: 165), pl. 4, fig. 1 (*Dolichodes*). Type-locality: “Brazil”. Distr. – Brazil (Santa Catarina). Refs. – Enderlein, 1924b: 174 (*Dolichodes*); Hull, 1962 (1): 219. Type, lost.

jungens Schiner, 1867: 375. Type-locality: “Brazil”. Refs. – Walker, 1854: 495 (*Dolichodes*); Williston, 1891: 71 (cat.; *Dolichodes*); Kertész, 1909: 157 (cat.; *Dolichodes*); Bromley, 1946: 108 (cat.; *Dolichodes*); Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.); Artigas & Papavero, 1993: 311 (syn.). Type, NMW.

fuliginosum (Wiedemann), 1821: 218 (*Dasypogon*). Type-locality: “Brazil” [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Wiedemann, 1828: 380 (*Dasypogon*); Walker, 1854: 446 (*Dasypogon*); Schiner, 1866: 677; Jaennicke, 1867: 359;

Williston, 1891: 69 (cat.); Kertész, 1909: 145 (cat.); Bristowe, 1925: 483-484 ("One day [...] I had just finished watching a long-bodied Sphegid wasp, *Sphex* (sub-gen. *Isodontia*) *costipennis* Spin. (*chrysobapta* Sm. and *petiolata* Sm.) depositing a Locustid grasshopper in a hollow bamboo, when I saw, as I thought, another specimen flying close to me. I got my net ready, and then I saw the insect fly without any undue haste at an Epeirid spider, *Epeira grayi* Bl., which was sitting in the middle of its web. I seized the spider, which immediately collapsed without showing any resistance, and calmly sat in the center of the web with its proboscis buried in the spider's abdomen. As I watched I saw a male spider, which, according to custom, had been sitting on the outskirts of the web, come quietly climbing down to see what was happening. He stopped an inch or two away and then, as though realizing his own danger, he turned round and crept quietly away. Before I left this locality (which was between Rio and Petropolis) I saw another of these wasp-like Asilid flies, *Plesiomma fuliginosa* Wied. male, as they turned out to be, attack another spider of the same species in very much the same way, thus showing that this interesting reversal of the usual role of the spider and the fly was no accidental occurrence. The invitation 'Come into my parlour said the spider to the fly' has in this case been made once too often"); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.); Artigas & Papavero, 1993: 311. Type, SENCK [cf. Geller-Grimm, 1999: 209].

?*caminarius* Wiedemann, 1828: 585 (*Dasypogon*). Type-locality: "Brazil" [Freireiss; cf. Papavero, 1971e: 57-58]. Refs. – Walker, 1854: 443 (*Dasypogon*); Schiner, 1866: 681 (as syn. of *caedens* Wiedemann, 1828); Williston, 1891: 69 (cat.); Kertész, 1909: 145 (cat.); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 219; Martin & Papavero, 1970: 21 (cat.); Artigas & Papavero, 1993: 311 (syn.). Type, SENCK [cf. Geller-Grimm, 1999: 207].

nigrum Macquart, 1838: 55 (1839: 171). Type-locality: "Brazil". Ref. – Walker, 1854: 501. Type, ?

funestum Loew, 1861: 35 (Cent. 7, no. 31). Type-locality: "Cuba". Distr. – Cuba. Refs. – Loew, 1866: 18; Osten Sacken, 1874: 177; Aldrich, 1905: 260 (cat.); Kertész, 1909: 145 (cat.); Bromley, 1929a: 276; Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 45 (cat.); Artigas & Papavero, 1993: 311. Type, MCZ.

lugubris Jaennicke, 1867: 356 (*Dioctria*). Type-locality: "Cuba". Type, SENCK [cf. Geller-Grimm, 1999: 210].

haemorrhoum (Fabricius), 1805: 161 (*Laphria*). Type-locality: "South America (Smidt)" [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Refs. – Loew, 1851: 20 (*Dasylysis*); Hansen, 1883: pl. 4, fig. 13; Kertész, 1909: 146 (cat.); Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.); Artigas & Papavero, 1993: 311. Type, COPN [cf. Zimsen, 1964: 469].

haemorrhoea Zimsen, 1964: 469 (*Laphria*), error.

indecorum Loew, 1866: 19 (Cent. 7, no. 33). Type-locality: "Cuba". Distr. – Cuba. Refs. – Johnson, 1894: 274; Aldrich, 1905: 260 (cat.); Kertész, 1909: 146 (cat.); Bromley, 1929a: 276, pl. 1, fig. 12; Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 45 (cat.); Artigas & Papavero, 1993: 311. Type, MCZ.

longiventris Schiner, 1867: 375. Type-locality: "Cuba". Ref. – Aldrich, 1905: 261 (cat.); Kertész, 1909: 146 (cat.). Type, NMW.

leptogaster (Perty). 1833: 180, pl. 36, fig. 2 (*Laphria*). Type-locality: Brazil, Bahia. Refs. – Walker, 1854: 504 (*Dasypogon*); Williston, 1891: 67 (cat.; *Dasypogon*); Kertész, 1909: 129 (cat.; *Dasypogon*); Bromley, 1946: 108 (cat.); Hull, 1962 (1): 219, 228 (*Dasypogon*); Maldonado-Capriles & Navarro, 1967: 52; Alayo Dalmau & Garcia Ávila, 1983: 45 (cat.); Artigas & Papavero, 1993: 311. Type, ZSBS.

lineatum (Fabricius), 1781: 465 (*Asilus*). Type-locality: "St. Croix Island". Distr. – Mexico (Yucatán), Virgin Islands (St. Croix). Refs. – Fabricius, 1787: 360 (*Asilus*); Olivier, 1789: 269 (*Asilus*); Gmelin, 1792: 2900 (*Asilus*); Fabricius, 1794: 386 (*Asilus*); 1805: 167 (*Dasypogon*); Wiedemann, 1821: 221 (*Dasypogon*), 1828: 385 (*Dasypogon*); Walker, 1854: 428 (*Dasypogon*); Schiner, 1866: 677, 1867: 374; Aldrich, 1905: 260 (cat.); Kertész, 1909: 146 (cat.; as *leptogastrum*); Bromley, 1929a: 276; Carrera & d'Andretta, 1950: 166; Hull, 1962 (1): 219 [*Plesiomma (Boropis)*]]; Farr, 1965: 7, figs. 1-2; Martin & Papavero, 1970: 22 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 45 (cat.); Artigas & Papavero, 1993: 311; Scarbrough & Perez-Gelabert, 2006: 80. Type, KIEL (only wings remaining) [cf. Zimsen, 1964: 470].

loewi Artigas & Papavero, 1993: 311 (*nom. nov* for *leptogaster* Loew, 1866). Type-locality: "Cuba".

leptogaster Loew, 1866: 18 (preocc. Perty, 1833). Type-locality: "Cuba". Refs. – Aldrich, 1905: 146 (cat.); Kertész, 1909: 146 (cat.); Martin & Papavero, 1970: 22 (cat.). Type, ?

macrum Loew, 1861: 35 (Cent. 7, no. 34). Type-locality: "Cuba". Distr. – Cuba. Refs. – Loew, 1866: 20; Aldrich, 1905: 261 (cat.); Kertész, 1909: 146 (cat.); Scarbrough & Perez-Gelabert, 2006: 81. Type, MCZ.

salti Bromley, 1929a: 277. Type-locality: Cuba, Soledad. Distr. – Cuba. Refs. – Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.); Alayo Dalmau & Garcia Ávila, 1983: 45 (cat.); Artigas & Papavero, 1993: 311. Type, MCZ.

simile Scarbrough & Perez-Gelabert, 2006: 73 (in key), 81, figs. 105-111, 122. Type-locality: Dominican Republic, Barahona, nr. Filipinas Larimar Mine. Distr. – Dominican Republic (Barahona, San Cristóbal). Type, FSCA.

testaceum (Fabricius), 1805: 167 (*Dasypogon*). Type-locality: South America (Smidt) [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Guyana. Refs. – Wiedemann, 1821: 222 (*Dasypogon*), 1828: 386 (*Dasypogon*); Walker, 1854: 431 (*Dasypogon*); Schiner, 1866: 677; Williston, 1891: 68 (cat.); Kertész, 1909: 146 (cat.); Curran, 1934c: 331; Hull, 1962 (1): 219; Martin & Papavero, 1970: 22 (cat.); Artigas & Papavero, 1993: 311. Syntypes, COPN, KIEL (in bad condition) [cf. Zimsen, 1964: 470].

Tribe Stenopogonini Hull, 1962

Genus **Ospriocerus** Loew, 1866

Ospriocerus Loew, 1866: 29 (Cent. 7, no. 51). Type-species, *Dasypogon aecus* Wiedemann, 1828 (sub. des., Back, 1909: 184) = *Ospriocerus abdominalis* (Say, 1824).

abdominalis (Say), 1824: 50 (!859: 64) (*Asilus*). Type-locality: U. S. A., "Northwest Territory". Distr. – U. S. A., Mexico (Coahuila, Chihuahua, Durango, Nuevo León, San Luis Potosí). Refs. – Coquillett, 1898: 37; Aldrich, 1905: 255 (cat.); Kertész, 1909: 81 (cat.); Brown, 1929: 301, pl. 29, figs. 12-15; Baker, 1939: 167; Hull, 1962 (1): 129, figs. 89, 1722, 1728; Martin, 1965a: 23; Martin & Wilcox, 1965: 381 (cat.); Martin & Papavero, 1970: 13 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312.

aecus Wiedemann, 1828: 390 (*Dasypogon*; nom. nov. for *abdominalis* Say, 1824, when preocc. in *Dasypogon*). Type-locality: U. S. A., "Northwest Territory". Refs. – Walker, 1854: 425 (*Dasypogon*); Schiner, 1866: 678 (*Dasypogon*); Loew, 1866: 29; Osten Sacken, 1877: 290; Williston, 1883: pl. 1, figs. 5, 5a; Osten Sacken, 1887: 168.

spathulatus Bellardi, 1861: 182 (82), pl. 1, fig. 9 (*Dasypogon*). Type-locality: "Mexico". Refs. – Loew, 1872: 68 (note); Aldrich, 1905: 255 (cat.); Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipteral). Type, TORO.

aeacides Loew, 1866: 29 (Cent. 7, no. 51). Type-locality: U. S. A., California. Ref. – Artigas & Papavero, 1993: 312. Type, MCZ.
ventralis Coquillett, 1898: 37. Type-locality: U. S. A., Arizona. Refs. – Aldrich, 1905: 255 (cat.); Kertész, 1909: 81 (cat.); Martin & Wilcox, 1965: 381 (cat., syn.).

aeacidinus (Williston), 1886: 289 (*Stenopogon*). Type-locality: U. S. A., Kansas. Distr. – U. S. A. (Kansas, Montana, Wyoming, Colorado, Nevada, Kansas, Oklahoma, Texas), Mexico (Coahuila, Tamaulipas). Refs. – Bromley, 1935c: 2 (*Stenopogon*); Baker, 1939: 167 (*Stenopogon*); Martin & Wilcox, 1965: 381 (cat.; n. comb.).

alamoensis Martin, 1968c: 409, figs. 17, 50, 60. Type-locality: Mexico, Sinaloa, Ellota. Distr. – Mexico (Sinaloa). Refs. – Martin & Papavero, 1970: 13 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, CAS.

arizonensis (Bromley), 1937a: 305 (*Stenopogon*). Type-locality: U. S. A., New Mexico, Albuquerque. Distr. – U. S. A. (New Mexico), Mexico (Sonora, Chihuahua). Refs. – Martin & Wilcox, 1965: 381 (cat.; n. comb.); Martin & Papavero, 1970: 13 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, USNM.

brevis Martin, 1968c: 406 (in key), 411, figs. 49, 58. Type-locality: U. S. A., California, Kern County, Randsburg. Distr. – U. S. A. (Arizona, California, Nevada), Mexico (Baja California, Sonora). Type, UCD.

diversus Williston, 1901: 303, pl. 5, fig. 16. Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. – Mexico (Guerrero, Oaxaca). Refs. – Aldrich, 1905: 255 (cat.); Kertész, 1909: 81 (cat.); Hull, 1962 (1): 129, fig. 427, 898, 907; Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipteral). Type, BMNH.

evansi Martin, 1968c: 423, figs. 16, 51, 64. Type-locality: Mexico, Guerrero, Iguala. Distr. – Mexico (Morelos, Guerrero). Refs. – Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, CAS.

latipennis (Loew), 1866: 28 (Cent. 7, no. 49) (*Stenopogon*). Type-locality: U. S. A., New Mexico. Distr. – U. S. A. (New Mexico), Mexico (Tamaulipas). Refs. – Baker, 1939: 167 (*Stenopogon*); Carrera & d'Andretta, 1950: 166 (*Stenopogon*); Martin & Wilcox, 1965: 381 (cat.; n. comb.); Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, MCZ.

consanguineus Loew, 1866: 27 (Cent. 7, no. 49) (*Stenopogon*). Type-locality: U. S. A., Nebraska. Ref. – Martin & Wilcox, 1965: 381 (cat.; syn.). Type, MCZ.

longulus (Loew), 1866: 28 (Cent. 7, no. 50) (*Stenopogon*). Type-locality: U. S. A., New Mexico. Distr. – U. S. A. (New Mexico), Mexico (Chihuahua, Tamaulipas, San Luis Potosí). Refs. – Bromley, 1935c: 2 (*Stenopogon*); Martin & Wilcox, 1965: 381 (cat.; n. comb.); Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, MCZ.

minos Osten Sacken, 1887: 291. Type-locality: Mexico, Sonora. Distr. – U. S. A., Mexico (Coahuila, Sonora). Refs. – Coquillett, 1898: 37; Aldrich, 1905: 255 (cat.); Kertész, 1909: 81 (cat.); Baker, 1939: 167; Hull, 1962 (1): 129; Martin & Wilcox, 1965: 382 (cat.); Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, BMNH.

nitens (Coquillett), 1904c: 34 (*Stenopogon*). Type-locality: U. S. A., Texas. Distr. – U. S. A. (Texas), Mexico (Tamaulipas). Refs. – Hull, 1962 (1): 129; Martin & Wilcox, 1965: 382 (cat.); Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, USNM.

monki Bromley, 1934d: 225. Type-locality: U. S. A., Texas.

painterorum Martin, 1968c: 417, figs. 25, 37. Type-locality: Mexico, Coahuila, 58 miles north of Monclova. Distr. – Mexico (Coahuila). Refs. – Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, CAS.

sinaloensis Martin, 1968c: 418, figs. 27, 43. Type-locality: Mexico, Sonora, Navojoa. Distr. – Mexico (Coahuila, Sonora). Refs. – Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, CAS.

- tenebrosus** (Coquillett), 1904c: 33 (*Stenopogon*). Type-locality: U. S. A., Texas, Brownville. Distr. – U. S. A. (Texas), Mexico (Chihuahua). Refs. – Kertész, 1909: 80 (cat.; *Stenopogon*); Schaeffer, 1916: 65 (*Stenopogon*); Martin & Wilcox, 1965: 381 (cat.; n. comb.); Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, USNM.
- tequilae** Martin, 1968c: 419, figs. 4-5, 46, 60. Type-locality: Mexico, Jalisco, Tequila. Distr. – U. S. A. (Arizona), Mexico (Jalisco, Nayarit, Sinaloa, Oaxaca, Campeche). Refs. – Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, CAS.
- youngi** Martin, 1968c: 421, figs. 20, 36. Type-locality: Mexico, Tamaulipas, Monté, 800 ft. Distr. – Mexico (Tamaulipas). Refs. – Martin & Papavero, 1970: 14 (cat.; *Stenopogon (Ospriocerus)*); Artigas & Papavero, 1993: 312. Type, CAS.

Genus *Scleropogon* Loew, 1866

Scleropogon Loew, 1866: 26 (Cent. 7, no. 45). Type-species, *Scleropogon picticornis* Loew, 1866 (orig. des.).

- dispar** (Bromley), 1937a: 306 (*Stenopogon*). Type-locality: U. S. A., Arizona, Patagonia. Distr. – U. S. A. (Arizona), Mexico. Refs. – Hull, 1962 (1): 124; Wilcox, 1971: 97, figs. 236-239. Type, KU.
- duncani** (Bromley), 1937a: 307 (*Stenopogon*). Type-locality: U. S. A., New Mexico, Silver City. Distr. – U. S. A. (New Mexico), Mexico (Sonora). Refs. – Hull, 1962 (1): 124; Martin & Papavero, 1970: 12 (cat.; *Stenopogon*); Wilcox, 1971: 98, figs. 212-215; Artigas & Papavero, 1993: 312. Type, CAS.
- haigi** Wilcox, 1971: 100, figs. 240-243. Type-locality: U. S. A., Arizona, 10 miles east of Nogales. Distr. – U. S. A. (Arizona), Mexico (Durango, Sinaloa). Type, UCD.
- indistinctus** (Bromley), 1937a: 308 (*Stenopogon*). Type-locality: U. S. A., Arizona, White Mts. Distr. – U. S. A. (Arizona), Mexico (Zacatecas, Chihuahua, Tamaulipas). Refs. – Carrera & d'Andretta, 1950: 166, figs. 21, 41; Hull, 1962 (1): 124; Martin & Papavero, 1970: 12 (cat.; *Stenopogon*); Wilcox, 1971: 105, figs. 254-257; Artigas & Papavero, 1993: 312. Type, CAS.
- lugubris** Williston, 1901: 304. Type-locality: Mexico, Guerrero, Dos Arroyos. Distr. – Mexico (Guerrero). Refs. – Aldrich, 1905: 256 (cat.); Kertész, 1909: 76 (cat.); Hull, 1962 (1): 124 (*Stenopogon*); Martin & Papavero, 1970: 12 (cat.; *Stenopogon*); Artigas & Papavero, 1993: 312; Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Type, BMNH.
- oaxacensis** (Martin), 1968b: 390, fig. 4 (*Stenopogon*). Type-locality: Mexico, Oaxaca, El Camarón. Distr. – Mexico (Oaxaca). Refs. – Martin & Papavero, 1970: 12 (cat.; *Stenopogon*); Artigas & Papavero, 1993: 312. Type, CAS.
- petilus** (Martin), 1968b: 302, figs. 11, 18, 30 (*Stenopogon*). Type-locality: Mexico, Sonora, Guaymas. Distr. – Mexico (Sonora). Refs. – Martin & Papavero, 1970: 12 (cat.; *Stenopogon*); Wilcox, 1971: 110 (as probable syn. of *picticornis* Loew, 1866); Artigas & Papavero, 1993: 312. Type, CAS.
- picticornis** Loew, 1866: 26. Type-locality: U. S. A., California. Distr. U. S. A. (Arizona, California, Colorado, Kansas, Montana, Nevada, New Mexico, South Dakota, Texas, Wyoming), Mexico (Baja California Sur, Chihuahua, Sonora). – Refs. – Coquillett, 1904b: 179; Aldrich, 1905: 256 (cat.); Back, 1909: 209 (*Stenopogon (Scleropogon)*); Kertész, 1909: 76 (cat.); Hull, 1962 (1): 124; Wilcox, 1971: 108, figs. 260-263. Type, MCZ.
- texanus** (Bromley), 1931b: 431 (*Stenopogon*). Type-locality: U. S. A., Texas, Dilley. Distr. – U. S. A. (Texas), Mexico (Tamaulipas). Refs. – Hull, 1962 (1): 124; Martin & Papavero, 1970: 13 (cat.; *Stenopogon*); Wilcox, 1971: 111, figs. 228-231; Artigas & Papavero, 1993: 312. Type, OSU.

Genus *Stenopogon* Loew, 1847

Dasypogon, subg. **Stenopogon** Loew, 1847: 453. Type-species, *Asilus sabaudus* Fabricius, 1794 (sub. des., Coquillett, 1910: 179).

- albalulus** Martin, 1968b: 374. Type-locality: Mexico, Michoacán, Apatzingán. Distr. – Mexico (Michoacán). Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, CAS.
- indistinctus* Bromley of Carrera & d'Andretta, 1950: 166, misident.
- arnaudi** Martin, 1968b: 375, figs. 13, 22. Type-locality: Mexico, Baja California Norte, Sierra San Pedro Martir, Rancho Viejo, 7000 ft. Distr. – U. S. A. (Arizona, California), Mexico (Baja California). Refs. – Martin & Papavero, 1970: 12 (cat.); Wilcox, 1971: 16; Artigas & Papavero, 1993: 313. Type, CAS.
- blaisdelli** Wilcox, 1971: 41, figs. 72-75. Type-locality: U. S. A., California, San Diego County, Coronado. Distr. – U. S. A. (California), Mexico. Type, CAS.
- boharti** Bromley, 1951: 6. Type-locality: U. S. A., Arizona, Yuma. Distr. – U. S. A. (California, Arizona), Mexico. Refs. – Hull, 1962 (1): 124; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1971: 27, figs. 36-39. Type, AMNH.
- rossi** Martin, 1968b: 393, fig. 2. Type-locality: Mexico, Baja California Sur, Coyote Cove, Concepción Bay. Ref – Martin

- & Papavero, 1970: 13 (cat.); Wilcox, 1971: 27 (syn.); Artigas & Papavero, 1993: 313. Type, CAS.
- breviusculus** Loew, 1872a: 68 (Cent. 10, no. 28). Type-locality: U. S. A., California. Distr. – U. S. A. (California), Mexico. Refs. – Aldrich, 1905: 256 (cat.); Back, 1909: 1906, pl. 12, fig. 3; Kertész, 1909: 77 (cat.); Bromley, 1937a: 299, fig. 1; Hull, 1962 (1): 124; Martin & Wilcox, 1965: 383 (cat.); Wilcox, 1971: 30, figs. 40-43. Type, MCZ.
- colimae** Martin, 1968b: 377, fig. 16. Type-locality: México, Colima, Vulcano. Distr. – Mexico (Colima). Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, USNM.
- dorothyae** Martin, 1968b: 378, fig. 26. Type-locality: Mexico, Guerrero, Chilpancingo. Distr. – Mexico (Guerrero, Morelos, Oaxaca). Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, CAS.
- elizabethae** Martin, 1968b: 379, figs. 15, 20, 33. Type-locality: Mexico, Chihuahua. Distr. – Mexico (Chihuahua). Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, CAS.
- felis** Bromley, 1931b: 429. Type-locality: U. S. A., California, Coronado. Distr. – U. S. A. (California), Mexico. Refs. – Hull, 1962 (1): 124; Martin & Wilcox, 1965: 384 (cat.); Wilcox, 1971: 65, figs. 144-147. Type, OSU.
- flavotibialis** Martin, 1968b: 380, figs. 8, 31. Type-locality: Mexico, Baja California Norte, Sierra San Pedro Martir, La Grulla, 6900 ft. Distr. – Mexico (Baja California Norte). Refs. – Martin & Papavero, 1970: 12 (cat.); Wilcox, 1971: 75, figs. 140-143; Artigas & Papavero, 1993: 313. Type, CAS.
- fuscolimbatus** Bigot, 1878: 421. Type-locality: “Mexico”. Distr. – Mexico (Guerrero, Morelos, Oaxaca). Refs. – Williston, 1883: 33; Aldrich, 1905: 256 (cat.); Kertész, 1909: 78 (cat.); Hull, 1962 (1): 124; Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313; Papavero & Ibáñez Bernal, 2001: 121 (hist. Mexican dipterol). Type, OXF.
- galbinus** Martin, 1968b: 385, fig. 24. Type-locality: “Mexico”. Distr. – Mexico. Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, USNM.
- hamus** Martin, 1968b: 385, fig. 1. Type-locality: Mexico, Baja California Norte, Sierra San Pedro Martir, La Grulla, 6900 ft. Distr. – Mexico (Baja California). Refs. – Martin & Papavero, 1970: 12 (cat.); Wilcox, 1971: 55, figs. 104-107; Artigas & Papavero, 1993: 313. Type, CAS.
- hiemalis** Martin, 1968b: 386, fig. 29. Type-locality: Mexico, Oaxaca, Tangola Bay. Distr. – Mexico (Oaxaca). Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, AMNH.
- jubatus** (Coquillett) in Baker, 1904: 38 (*Scleropogon*). Type-locality: U. S. A., California, Claremont. Distr. – U. S. A. (California), Mexico. Refs. – Back, 1909: 195; Kertész, 1909: 76 (cat., *Scleropogon*); Hull, 1962 (1): 124; Martin & Wilcox, 1965: 384 (cat.). Type, USNM.
- nigroverticellus* Bromley, 1937a: 298. Type-locality: U. S. A., California, San Diego. Refs. – Hull, 1962 (1): 124; Wilcox, 1971: 56 (syn.). Type, KU.
- nigritulus** Coquillett, 1904b: 179. Type-locality: U. S. A., California, Los Angeles County. Distr. – U. S. A. (California), Mexico (Baja California Norte). Refs. – Back, 1909: 197; Kertész, 1909: 79 (cat.); Hull, 1962 (1): 124; Martin & Wilcox, 1965: 384 (cat.); Wilcox, 1971: 22, figs. 24-27. Type, USNM.
- nigrolimbatus** Martin, 1968b: 389, fig. 9. Type-locality: Mexico, Nuevo León, Galeana. Distr. – Mexico (Nuevo León). Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, CAS.
- ortegai** Martin, 1968b: 391, figs. 10, 28, 32. Type-locality: Mexico, Guerrero, Chilpancingo, 1600 ft. Distr. – Mexico (Guerrero). Refs. – Martin & Papavero, 1970: 12 (cat.); Artigas & Papavero, 1993: 313. Type, CAS.
- setosus** Bezark, 1984: 53, figs. 1-4. Type-locality: Costa Rica, Guanacaste, 4 km northwest of Canas. Distr. – Costa Rica (Guanacaste). Type, CAS.
- silaceus** Martin, 1968b: 394, figs. 5, 23, 35. Type-locality: Mexico, Guanajuato, Celaya. Distr. – Mexico (Guanajuato, Morelos, San Luis Potosí, Zacatecas). Refs. – Martin & Papavero, 1970: 13 (cat.); Artigas & Papavero, 1993: 313. Type, CAS.
- stonei** Bromley, 1937a: 298. Type-locality: Mexico, Baja California Norte, Ensenada. Distr. – Mexico (Baja California). Refs. – Hull, 1962 (1): 124; Martin, 1968b: 372, 394; Martin & Papavero, 1970: 13 (cat.); Wilcox, 1971: 63, figs. 136-139; Artigas & Papavero, 1993: 313. Type, CAS.
- surrufus** Martin, 1968b: 395, figs. 12, 19. Type-locality: Mexico, Guerrero, Mexcala. Distr. – Mexico (Guerrero). Refs. – Martin & Papavero, 1970: 13 (cat.); Artigas & Papavero, 1993: 313. Type, CU.
- truquii** (Bellardi), 1861: 176 (76), pl. 1, fig. 10 (*Dasyopogon*). Type-locality “Mexico”. Distr. – Mexico (Morelos, Veracruz). Refs. – Osten Sacken, 1887: 256 (*Scleropogon*); Williston, 1901: 256 (*Scleropogon*); Kertész, 1909: 76 (cat.; *Scleropogon*); Hull, 1962 (1): 124 (*Scleropogon*); Martin & Papavero, 1970: 13 (cat.); Artigas & Papavero, 1993: 313; Papavero & Ibáñez Bernal, 2001: 107 (hist. Mexican dipterol). Type, TORO.
- wilcoxi** Bromley, 1937a: 300. Type-locality: U. S. A., California, San Diego Co. Distr. – U. S. A. (California), Mexico. Refs. – Hull, 1962 (1): 124; Martin & Wilcox, 1965: 384 (cat.); Wilcox, 1971: 87, figs. 200-203. Type, KU.
- williamsi** Wilcox, 1971: 38, figs. 68-71. Type-locality: U. S. A., California, San Diego County, La Mesa. Distr. – U. S. A. (California), Mexico. Type, CAS.
- xochimilcae** Martin, 1968b: 396, figs. 6, 25. Type-locality: Mexico, Distrito Federal, Xochimilco. Distr. – Mexico (Distrito Federal, Jalisco). Refs. – Martin & Papavero, 1970: 13 (cat.); Artigas & Papavero, 1993: 313. Type, USNM.
- truquii* Bellardi of Williston, 1901: 256, misident.
- zebra** Martin, 1968b: 397, figs. 7, 27, 34. Type-locality: Mexico, Guerrero, Chilpancingo. Distr. – Mexico (Guerrero). Refs. – Martin & Papavero, 1970: 13 (cat.); Artigas & Papavero, 1993: 313. Type, CAS.

Tribe Tillobromatini Artigas & Papavero, 1991

Tillobromini Artigas & Papavero, 1991a: 141, 143. Type-genus, *Tillobroma* Hull, 1962.
Tillobromatini Sabrosky, 1999: 308, corrected spelling.

Genus **Euthrixius** Artigas, 1971

Euthrix Philippi, 1865: 690 (preocc. Meigen, 1930). Type-species, *Dasypogon venustus* Philippi, 1865 (sub. des., Martin & Papavero, 1970: 17).

Euthrixius Artigas, 1971: 22 (*nom. nov.* for *Euthrix* Philippi, 1865). Type-species, *Dasypogon venustus* Philippi, 1865 (aut.).

distinguendus (F. Lynch Arribálzaga) *in* E. Lynch Arribálzaga, 1881: 121 (*Scylaticus*). Type-locality: Argentina, Buenos Aires, Chacabuco. Distr. – Argentina (Buenos Aires). Refs. – Lynch Arribálzaga, 1883a: 142 (cat.; *Scylaticus*); Osten Sacken, 1891: 418 (*Scylaticus*); Kertész, 1909: 103 (cat.; *Scylaticus*); Hull, 1962 (1): 145 (*Scylaticus*); Martin & Papavero, 1970: 17 (cat.; *Scylaticus*); Artigas & Papavero, 1991e: 18 (n. comb.). Type, lost.

rufipes (Philippi), 1865: 691 (*Dasypogon*). Type-locality: “Chile”. Neotype-locality: Chile, Curicó, Estero La Jaula. Distr. – Chile (Aconcagua, Curicó, Valparasí). Refs. – Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 131 (cat.; *Dasypogon*); Stuardo Ortiz, 1946: 82 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Artigas, 1970a: 178, figs. 159, 161–163, 451 (*Scylaticus*); Martin & Papavero, 1970: 18 (cat.; *Scylaticus*); Artigas, 1971: 22; Artigas & Papavero, 1991e: 18. Type, lost. NT DZUC [des. by Artigas, 1976: 178].

philippii Schiner, 1868: 163 (*Scylaticus*). Type-locality: “Chile”. Refs. – Williston, 1891: 73 (cat.; *Scylaticus*); Kertész, 1909: 104 (cat.; *Scylaticus*); Stuardo Ortiz, 1946: 81 (cat.; *Scylaticus*); Hull, 1962 (1): 1945 (*Scylaticus*); Martin & Papavero, 1970: 18 (syn.). Type, NMW.

rubripes Bigot, 1878: 435 (*Scylaticus*). Type-locality: “Chile”. Refs. – Williston, 1891: 73 (cat.; *Scylaticus*); Kertész, 1909: 104 (cat.; *Scylaticus*); Stuardo Ortiz, 1946: 81 (cat.; *Scylaticus*); Hull, 1962 (1): 145 (*Scylaticus*); Martin & Papavero, 1970: 18 (syn.). Type, OXF.

venustus (Philippi), 1865: 689 (*Dasypogon*). Type-locality: Chile, near Santiago. Neotype-locality: Chile, Santiago, El Canelo. Distr. – Chile (Coquimbo, Santiago, Valparaíso). Refs. – Bigot, 1878: 221 (*Scylaticus*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 104 (cat.; *Scylaticus*); Hull, 1962 (1): 145 (*Scylaticus*); Artigas, 1970a: 180, figs. 151, 160, 164, 427 (*Scylaticus*); Martin & Papavero, 1970: 18 (cat.); Artigas, 1971: 22, figs. 65–66 (n. comb.); Artigas & Papavero, 1991e: 18. Type, lost. NT DZUC [des. by Artigas, 1970a: 180].

Genus **Grajahua** Artigas & Papavero, 1991

Grajahua Artigas & Papavero, 1991e: 19. Type-species, *Grajahua lopesi* Artigas & Papavero, 1991 (orig. des.).

lopesi Artigas & Papavero, 1991e: 20, figs. 1–4. Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (Grajaú). Distr. – Brazil (Rio de Janeiro). Type, MZUSP.

Genus **Scylaticina** Artigas & Papavero, 1991

Scylaticina Artigas & Papavero, 1991e: 21. Type-species, *Scylaticina tucumana* Artigas & Papavero, 1991 (orig. des.).

tucumana Artigas & Papavero, 1991e: 23, figs. 9, 17. Type-locality: Argentina, Tucumán, San Pedro Colalao. Distr. – Argentina (Tucumán). Type, IML.

Genus **Scylaticodes** Artigas & Papavero, 1991

Scylaticodes Artigas & Papavero, 1991e: 23. Type-species, *Dasypogon chilensis* Macquart, 1850 (orig. des.).

carrascoi (Artigas), 1974b: 5, figs. 1–10 (*Scylaticus*). Type-locality: Peru, Cuzco, Urubamba. Distr. – Peru (Cuzco, Puno),

- Chile Arica). Ref. – Artigas & Papavero, 1991e: 26 (n. comb.). Type, DZUC.
- chilensis** (Macquart), 1850: 48 (*Dasypogon*). Type-locality: “Chile”. Distr. – Chile (Coquimbo, O’Higgins, Santiago). Refs. – Blanchard, 1852: 367 (*Dasypogon*); Stuardo Ortiz, 1946: 84 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Artigas, 1970a: 172, figs. 148, 152, 154-155, 465 (*Scylaticus*), 1971: 29, figs. 63-64 (*Scylaticus*); Artigas & Papavero, 1991e: 26 (n. comb.). Type, MNHNP.
- tricolor** Philippi, 1865: 688 (*Dasypogon*). Type-locality: Chile, Colchagua. Refs. – Bigot, 1878: 221 (*Dasypogon*); Williston, 1891: 73 (cat., *Scylaticus*); Kertész, 1909: 104 (cat.; *Scylaticus*); Stuardo Ortiz, 1946: 81 (cat.; *Scylaticus*); Hull, 1962 (1): 145 (*Scylaticus*). Type, MNHNS.
- cuneigaster** (Artigas), 1970a: 174, figs. 150, 153, 156, 429 (*Scylaticus*). Type-locality: Chile, Atacama, Carrizal Bajo, Quebrada Algodones. Distr. – Chile (Atacama). Ref. – Artigas & Papavero, 1991e: 26 (n. comb.). Type, DZUC.
- lugens** (Philippi), 1865: 689 (*Dasypogon*). Type-locality: Chile, Colchagua. Neotype-locality: Chile, Valparaíso, Granizo. Distr. – Chile (Santiago, Talca, Valparaíso). Refs. – Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 129 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Artigas, 1970a: 176, figs. 148, 157-158, 450; Artigas & Papavero, 1991e: 26 (n. comb.). Type, lost. NT DZUC [des. by Artigas, 1970a: 176].

Genus *Tillobroma* Hull, 1962

- Clavator* Philippi, 1865: 699 (preocc. Martens, 1860). Type-species, *Clavator punctipennis* Philippi, 1865 (sub. des., Hull, 1962: 152).
- Hypenetes*, subg. **Tillobroma** Hull, 1962 (1): 154. Type-species, *Clavator punctipennis* Philippi, 1865 (orig. des.). Ref. – Artigas & Papavero, 1991: 26 (generic status).
- Hypenetes* Loew of authors, misident.
- Ref. – Artigas, Lewis & Parra, 2005 (rev.).

- aconcagua** Artigas, Lewis & Parra, 2005: 531, figs. 7A-F, 34A. Type-locality: Chile, Aconcagua. Distr. – Chile (Aconcagua). Type, DZUC.
- angulata** Artigas, Lewis & Parra, 2005: 532, figs. 8A-D, 8F, 27F, 34B. Type-locality: Chile, Ñuble, Cardillera de Chillán, Invernada. Distr. – Chile (Ñuble). Type, DZUC.
- asiliformis** (Wulp), 1882b: 101 (*Hypenetes*). Type-locality: “Argentina”. Distr. – Chile (Magallanes), Argentina (Chubut, Río Negro, Neuquén). Refs. – Williston, 1891: 71 (cat.; *Hypenetes*); Kertész, 1909: 74 (cat.; *Hypenetes*); Bromley, 1932: 264 (*Hypenetes*); Hull, 1962 (1): 154 (*Hypenetes* (*Tillobroma*)); Artigas, 1970a: 127, figs. 87-91, 504 (*Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 548, figs. 8E, 8G-9D, 26E, 30D. Type, ZMA. [NT DZUC invalid, des. by Artigas, Lewis & Parra, 1991: 26].
- critesi** (Artigas), 1970a: 128, figs. 92-93, 95, 97, 428 (*Hypenetes*). Type-locality: Chile, Aysén, Balmaceda. Distr. – Chile (Arauco, Aysén, Biobío, Cautín, Curicó, Linares, Llanquihue, Malleco, Ñuble, Osorno, Valdivia), Argentina (Río Negro). Refs. – Artigas, 1971: 24, figs. 46-47 (*Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 549, figs. 10A-G, 28D, 30C. Type, DZUC.
- davidsoni** (Artigas), 1970a: 130, figs. 94, 96, 98-99, 420 (*Hypenetes*). Type-locality: Chile, Maule, Cauquenes. Distr. – Chile (Concepción, Curicó, Linares, Maule, Ñuble, O’Higgins, Santiago, Valparaíso). Refs. – Artigas, 1971: 25, figs. 53-54 (*Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 550, figs. 11A-G, 27D, 34C. Type, DZUC.
- digitata** (Artigas), 1970a: 132, figs. 100-101, 103-105, 484 (*Hypenetes*). Type-locality: Chile, Concepción, Lota-Colcura. Distr. – Chile (Arauco, Cautín, Concepción, Curicó, Malleco, Santiago, Valdivia, Valparaíso). Refs. – Artigas, 1971: 25, figs. 51-52 (*Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 551, figs. 12A-F, 27B, 33A. Type, DZUC.
- fucosa** (Artigas), 1970a: 134, figs. 102, 106-107, 109, 468 (*Hypenetes*). Type-locality: Chile, Arauco, Cordillera de Nahuelbuta, Pichinahuel. Distr. – Chile (Arauco, Malleco, Ñuble, Osorno, Talca). Refs. – Artigas, 1971: 25, figs. 56-57 (*Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 552, figs. 13A-F, 27G, 33C. Type, DZUC.
- fucosoides** Artigas, Lewis & Parra, 2005: 552, figs. 14A-F, 28C, 33D. Type-locality: Chile, Arauco, Nahuelbuta (Parque Cabrera). Distr. – Chile (Arauco, Curicó, Malleco, Valdivia). Type, DZUC.
- fulvicornis** (Macquart), 1846: 195 (1846: 67), pl. 7, fig. 1 (*Dasypogon*). Type-locality: “Brazil”. Distr. – Peru, Bolivia, Brazil (Rio de Janeiro, São Paulo, Santa Catarina), Argentina (Tucumán). Refs. – Walker, 1854: 440 (*Dasypogon*); Schiner, 1868: 159 (*Hypenetes*); Reed, 1888: 188 (*Hypenetes*); Williston, 1891: 71 (cat.; *Hypenetes*); Engel, 1930: 474 (*Hypenetes*); Bromley, 1932: 264 (*Hypenetes*); Carrera, 1945b: 170 (*Hypenetes*); Bromley, 1946: 105 (cat.; *Hypenetes*); Carrera & d’Andretta, 1952: 254 (*Hypenetes*; preys); Hull, 1962 (1): 154 (*Hypenetes*); Carrera & Papavero, 1965: 53 (*Hypenetes* (*Tillobroma*)); Martin & Papavero, 1970: 19 (cat.; *Hypenetes*); Lamas M., 1972: 306 (cat.; *Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 554, figs. 15A-G, 30A. Type, MNHNP.

- huasquina** Artigas, Lewis & Parra, 2005: 555, figs. 16A-F, 33B. Type-locality: Chile, Huasco, 10.3 km north of Puerto Huasco. Distr. – Chile (Atacama, Huasco). Type, DZUC.
- leucoptera** Artigas, Lewis & Parra, 2005: 555, figs. 16E-17D, 28H, 31A. Type-locality: Chile, O'Higgins, C. Poqui. Distr. – Chile (Curicó, O'Higgins, Valparaíso). Type, DZUC.
- leucotrica** Artigas, Lewis & Parra, 2005: 564, figs. 17E-18B, 28A, 31D. Type-locality: Chile, Atacama, 70 km south of Copiapó. Distr. – Chile (Atacama). Type, DZUC.
- magellanica** (Artigas), 1070: 136, figs. 108, 110-112, 418 (*Hypenetes*). Type-locality: Chile, Magallanes, Cerro Guido. Distr. – Chile (Lonquimay, Magallanes, Palena). Refs. – Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 565, figs. 19A-F, 28E, 31C. Type, DZUC.
- nahuelbutae** Artigas, Lewis & Parra, 2005: 566, figs. 20A-D, 28F, 31B. Type-locality: Chile, Angol, Nahuelbuta (Parque Cabrera). Distr. – Chile (Angol). Type, DZUC.
- obtusa** (Engel), 1930: 472 (*Hypenetes*). Type-locality: Bolivia, cordillera de Buenavista, south of Cuevo. Distr. – Peru, Bolivia, Brazil (São Paulo, Paraná), Argentina). Refs. – Bromley, 1932: 264 (*Hypenetes*); Carrera, 1945b: 169 (*Hypenetes*); Hull, 1962 (1): 154 (*Hypenetes (Tillobroma)*); Carrera & Papavero, 1965: 54 (*Hypenetes (Tillobroma)*); Lamas M., 1972: 306 (cat.; *Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 567, figs. 20E-G, 30E. Type, ?
- punctipennis** (Philippi), 1865: 699, pl. 26, figs. 31, 31a-b (*Clavator*). Type-locality: “Chile”. Neotype-locality: Chile, Arauco, Pichinahuel, 1100 m. Distr. – Chile (Arauco, Biobío, Linares, Lonquimay, Malleco, Ñuble, Santiago), Argentina. Refs. – Schiner, 1868: 159 (*Hypenetes*); Williston, 1891: 71 (cat.; *Hypenetes*); Kertész, 1909: 74 (cat.; *Hypenertes*); Bromley, 1932: 264 (*Hypenetes*); Stuardo Ortiz, 1946: 80 (cat.; *Hypenetes*); Hull, 1962 (1): 154 (*Hypenetes (Tillobroma)*); Carrera & Papavero, 1965: 54 (*Hypenetes (Tillobroma)*); Artigas, 1970a: 137, figs. 113-117, 419 (*Hypenetes*); Martin & Papavero, 1970: 19 (cat.; *Hypenetes*); Artigas, 1971: 26, figs. 48-49 (*Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 568, figs. 3, 5A-B, 21A-G, 27A, 31E. Type, lost. NT DZUC (des. by Artigas, 1970a: 137).
- purpurea** Artigas, Lewis & Parra, 2005: 569, figs. 22A-G, 28B, 32A. Type-locality: Chile, Santiago, Quebrada de La Plata. Distr. – Chile (Santiago, Valparaíso). Type, DZUC.
- schineri** (Artigas), 1970a: 139, figs. 118-123, 125, 493 (*Hypenetes*). Type-locality: Chile, O'Higgins, Rancagua. Distr. – Chile (Curicó, Linares, O'Higgins, Santiago, Valparaíso). Refs. – Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 570, figs. 6A-D, 23A-F, 28G, 32C. Type, DZUC.
- spinipes** Artigas, Lewis & Parra, 2005: 571, figs. 24A-E, 30B. Type-locality: Argentina, Córdoba, Copina, 100 m. Distr. – Argentina (Córdoba, Mendoza). Type, DZUC.
- tregualemuensis** Artigas, Lewis & Parra, 2005: 572, figs. 25A-F, 27H, 32B. Type-locality: Chile, Maule, Tregualemu. Distr. – Chile (Maule). Type, DZUC.
- valentinei** (Artigas), 1970a: 141, figs. 124, 126-127, 129, 419 (*Hypenetes*). Type-locality: Chile, Coquimbo. Distr. – Chile (Aconcagua, Angol, Chacabuco, Choapa, Coquimbo, Concepción, Curicó, Limarí, Ñuble, O'Higgins, Talca). Refs. – Artigas, 1971: 26, fig. 50 (*Hypenetes*); Artigas & Papavero, 1991e: 26 (n. comb.); Artigas, Lewis & Parra, 2005: 573, figs. 4, 26A-G, 27C, 32D. Type, DZUC.

Genus **Zabrotica** Hull, 1958

Zabrotica Hull, 1958b: 253. Type-species, *Zabrotica clarkei* Hull, 1958 (orig. des.).

clarkei Hull, 1958b: 254. Type-locality: Peru, Yauli, La Oroya. Distr. – Peru (Arequipa, Lima, Yauli). Refs. – Hull, 1962 (1): 152, figs. 123, 464, 1723, 1769, 1919; Lamas M., 1972: 306 (cat.); Artigas & Papavero, 1991e: 26. Type, USNM.

Unrecognized and unplaced *Tillobromatini*

- cruciger** Hermann, 1921: 119 (*Scylaticus*). Type-locality: “Paraguay”. Ref. – Hull, 1962 (1): 145 (*Scylaticus*). Type, ZSBS.
- nitidigaster** Macquart, 1850: 373 (1850: 69) (*Dasyopogon*). Type-locality: “Chile”. Refs. – Blanchard, 1852: 367 (*Dasyopogon*); Philippi, 1865: 686 (*Dasyopogon*); Williston, 1891: 73 (cat.; *Scylaticus*); Kertész, 1909: 103 (cat.; *Scylatyicus*); Stuardo Ortiz, 1946: 81 (cat.; *Scylaticus*); Hull, 1962 (1): 145 (*Scylaticus*); Martin & Papavero, 1970: 18 (cat.; *Scylaticus*). Type, lost.
- ruficauda** Bigot, 1878: 454 (*Scylaticus*). Type-locality: “Brazil, Amazonia”. Refs. – Osten Sacken, 1891: 418 (*Scylaticus*); Kertész, 1909: 104 (cat.; *Scylaticus*); Bromley, 1946: 106 (cat.; *Scylaticus*); Hull, 1962 (1): 145 (*Scylaticus*); Martin & Papavero, 1970: 18 (cat.; *Scylaticus*). Type, OXF.

Subfamily Stichopogoninae Hardy, 1930

Stichopogonini G. H. Hardy, 1930: 250, 251, 260. Type-genus, *Stichopogon* Loew, 1847.
 Stichopogoninae Artigas & Papavero, 1988a: 97.

Genus **Argyropogon** Artigas & Papavero, 1990

Argyropogon Artigas & Papavero, 1990: 41. Type-species, *Argyropogon argentinus* Artigas & Papavero, 1990 (orig. des.).
argentinus Artigas & Papavero, 1990: 41, figs. 5-18. Type-locality: Argentina, Santa Cruz, 5 km northwest of Piedrabuena, 130 m. Distr. – Argentina (Santa Cruz). Type, DZUC.

Genus **Lasiopogon** Loew, 1847

Dasypogon, subg. **Lasiopogon** Loew, 1847: 508. Type-species, *Dasypogon (Lasiopogon) pilosellus* Loew, 1847 (sub. des., Hull, 1962: 115). [Des. by Rondani, 1856: 156, of *Dasypogon hirtellus* Meigen, 1820, invalid because it was originally included as *species inquirenda*].
Daulopogon Loew, 1874: 377 (unjustified *nom. nov.* for *Lasiopogon* Loew, 1847). Type-species, *Dasypogon (Lasiopogon) pilosellus* Loew, 1847 (aut.).
Alexiopogon Curran, 1934b: 183. Type-species, *Daulopogon terricola* Johnson, 1900 (orig. des.).

Ref. – Cole & Wilcox, 1938 (rev.).

drabicola Cole, 1916: 65. Type-locality: U. S. A., California, Redlands. Distr. – U. S. A. (California), Mexico (Baja California). Refs. – Melander, 1923: 137; Cole & Wilcox, 1938: 46, pl. 1, fig. 8; Martin & Wilcox, 1965: 377 (cat.). Type, USNM.
gabrieli Cole & Wilcox, 1938: 49, pl. 3, fig. 33. Type-locality: U. S. A., California, San Gabriel River Wash, near Whittier. Distr. – U. S. A. (California), Mexico (Baja California). Ref. – Martin & Wilcox, 1965: 377 (cat.). Type, CAS.
zonatus Cole & Wilcox, 1938: 80, pl. 1, fig. 11. Type-locality: U. S. A., California, Orange County, Santa Ana Canyon. Distr. – U. S. A. (California), Mexiuco (Baja California). Ref. – Martin & Wilcox, 1965: 378 (cat.). Type, CAS.

Genus **Lissoteles** Bezzi, 1910

Lissoteles Bezzi, 1910a: 178. Type-species, *Lissoteles hermanni* Bezzi, 1910 (orig. des.).
acapulcae Martin, 1961a: 4 (in key), 7, fig. 2. Type-locality: Mexico, Guerrero, Acapulco. Distr. – Mexico (Guerrero). Refs. – Martin & Papavero, 1970: 4 (cat.); Artigas & Papavero, 1990: 41. Type, CAS.
aquilonius Martin, 1961a: 4 (in key), 8, fig. 3. Type-locality: Mexico, Nayarit, San Blás. Distr. – Mexico (Baja California, Nayarit, Sinaloa, Sonora). Refs. – Lamas M., 1972: 305 (cat.; Peru; ?misident.); Martin & Papavero, 1970: 4 (cat.); Artigas & Papavero, 1990: 41. Type, AMNH.
austrinus Martin, 1961a: 4 (in key), 10. Type-locality: Peru, Tumbes, 10 miles south of Zorritas. Distr. – Peru. Refs. – Lamas M., 1972: 305 (cat.); Martin & Papavero, 1970: 4 (cat.); Artigas & Papavero, 1990: 41. Type, CAS.
autumnalis Martin, 1961a: 4 (in key), 11. Type-locality: Mexico, Jalisco, Tenacatita Bay. Distr. – Mexico. Refs. – Martin & Papavero, 1970: 4 (cat.); Artigas & Papavero, 1990: 41. Type, AMNH.
blantoni Martin, 1961a: 4 (in key), 11. Type-locality: Panama, Río Mar. Distr. – Panama. Ref. – Kalletta, 1976: 66. Type, AMNH.
capronae Martin, 1961a: 4 (in key), 12. Type-locality: Honduras, Tegucigalpa. Distr. – Honduras. Refs. – Martin & Papavero, 1970: 4 (cat.); Artigas & Papavero, 1990: 42. Type, AMNH.
fernandezii Kalletta, 1976: 67. Type-locality: Venezuela, Miranda, Páez, Machurucuto. Distr. – Venezuela (Anzoátegui, Carabobo, Miranda). Ref. – Artigas & Papavero, 1990: 42. Type, FAUVC.
hermanni Bezzi, 1910a: 178. Type-locality: “Panama”. Distr. – Panama. Refs. – Carrera, 1953a: 271; James, 1953: 56; Martin, 1961a: 4, fig. 1; Hull, 1962 (1): 112, fig. 50; Martin & Papavero, 1970: 4 (cat.); Kalletta, 1976: 66; Artigas & Papavero, 1990: 42.
vanduzeei Cole, 1923: 465, fig. 7. Type-locality: Mexico, Baja California, Angeles Bay and San Francisquito Bay. Distr. – Mexico (Baja California). Refs. – Martin, 1961a: 4 (in key), 7, fig. 4; Hull, 1962 (1): 112, figs. 482, 933, 942; Martin & Papavero, 1970: 4 (cat.); Artigas & Papavero, 1990: 42. Type, CAS.

Genus *Stichopogon* Loew, 1847

Dasypogon, subg. ***Stichopogon*** Loew, 1847: 499. Type-species, *Dasypogon elegantulus* Wiedemann in Meigen, 1820: 270 (sub. des., Back, 1909: 332).

Stilopogon Costa, 1883: 62. Type-species, *Dasypogon trifasciatus* Say, 1823 (orig. des.).

Neopogon Bezzi, 1910c: 147. Type-species, *Dasypogon trifasciatus* Say, 1823 (orig. des.).

Stichopogan Wilcox, 1936b: 201, error.

Stictopogon Wilcox, 1937: 37, error.

catulus Osten Sacken, 1887: 170. Type-locality: Mexico, northern Sonora. Distr. – U. S. A. (Arizona), Mexico (Sonora). Refs. – Aldrich, 1905: 262 (cat.); Kertész, 1909: 81 (cat.); Wilcox, 1936d: 208; Hull, 1962 (1): 106; Martin & Wilcox, 1965: 385 (cat.); Martin & Papavero, 1970: 5 (cat.); Weeks & Hespenheide, 1985: 95 (predatory and mating behavior); Artigas & Papavero, 1990: 42; Papavero & Ibáñez Bernal, 2003: 161 (hist. Mexican dipterol). Type, BMNH.

fragilis Back, 1909: 334. Type-locality: U. S. A., New Mexico. Distr. – U. S. A. (Washington to Idaho, south to California and New Mexico), Mexico. Refs. – Hull, 1962 (1): 106; Martin & Wilcox, 1965: 385 (cat.). Type,

schnusei (Bezzi), 1910c: 151 (*Neopogon*). Type-locality: Peru, Cuzco, Rosalina, near Paso de Lares, Urubamba river. Refs. – Hull, 1962 (1): 110 (*Neopogon*); Lamas M., 1972: 305 (cat.); Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, ?

trifasciatus (Say), 1823: 51 (1859: 64) (*Dasypogon*). Type-locality: U. S. A., Pennsylvania. Distr. – U. S. A., Mexico. Refs. – Wiedemann, 1828: 405 (*Dasypogon*); Walker, 1854: 424, 503 (*Dasypogon*); Schiner, 1866: 64; Osten Sacken, 1874: 181, 1878: 70, 97 (cat.); Williston, 1885: 53, 1886: 289; Osten Sacken, 1887: 170; Townsend, 1895: 262; Aldrich, 1905: 262 (cat.); Kertész, 1909: 85 (cat.); Bezzi, 1910c: 148 (*Neopogon*); Bromley, 1935c: 3; Fattig, 1945: 17; Bromley, 1950: 232; Hull, 1962 (1): 110, figs. 450, 932, 941, 1917 (*Neopogon*); Martin & Wilcox, 1965: 385 (cat.); Martin & Papavero, 1970: 5 (cat.); Weeks & Hespenheide, 1985: 95 (predatory and mating behavior); Artigas & Papavero, 1990: 42. Type, lost.

candidus Macquart, 1846: 195 (1846: 67) (*Dasypogon*). Type-locality: U. S. A., Pennsylvania. Refs. – Walker, 1854: 440 (*Dasypogon*); Osten Sacken, 1874: 182; Bellardi, 1861: 78 (*Dasypogon*); Osten Sacken, 1874: 182; Bezzi, 1910c: 150 (*Neopogon*); Papavero & Ibáñez Bernal, 2001: 84 (hist. Mexican dipterol.). Type, lost.

plagiata Walker, 1848: 223 (*Thereva*). Type-locality: U. S. A., Massachusetts. Type, BMNH.

fasciventris Macquart, 1850: 373 (1850: 69), pl. 6, fig. 13 (*Dasypogon*). Type-locality: “Mexico”. Type, OXF.

gelascens Walker, 1860: 277 (*Dasypogon*). Type-locality: U. S. A., Massachusetts. Ref. – Papavero & Ibáñez Bernal, 2001: 95 (hist. Mexican dipterol.). Type, BMNH.

snowii Bezzi, 1910c: 149. Type-locality: U. S. A., Kansas, Wallace Co. Type, MCSNM.

venezuelanus Kaletta, 1976: 69 [*Stichopogon (Neopogon)*]. Type-locality: Venezuela, Miranda, Paz Castillo (Quebrada Soapire). Distr. – Venezuela (Guárico, Miranda). Ref. – Artigas & Papavero, 1990: 42. Type, FAUCV.

Genus *Townsendia* Williston, 1895

Townsendia Williston, 1895: 107. Type-species, *Townsendia minuta* Williston, 1895 (mon.).

albomacula Martin, 1966a: 545, figs. 1, 9-10. Type-locality: Mexico, Nayarit, San Blás. Distr. – Mexico (Colima, Guerrero, Nayarit, Sinaloa). Refs. – Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, KU.

araguensis Kaletta, 1976: 71. Type-locality: Venezuela, Aragua, San Sebastián, orilla aragüense del río Guárico. Distr. – Venezuela (Aragua, Carabobo, Guárico). Ref. – Artigas & Papavero, 1990: 42. Type, FAUCV.

argyrata Curran, 1926: 1. Type-locality: Puerto Rico, Loquillo National Forest. Distr. – Puerto Rico. Refs. – Curran, 1928b: 23; Hull, 1962 (1): 118; Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, AMNH.

dilata Martin, 1966a: 546, figs. 2, 5-6. Type-locality: Mexico, Veracruz, Veracruz. Distr. – Mexico (Veracruz). Refs. – Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, USNM.

fiebrigii Bezzi, 1909: 629, fig. Type-locality: “Paraguay”. Distr. – Brazil (São Paulo), Paraguay. Refs. – Carrera, 1945b: 168; Hull, 1962 (1): 118; Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42.

gracilis Martin, 1966a: 547, figs. 11-12. Type-locality: Mexico, Veracruz, Veracruz. Distr. – Mexico (Veracruz). Refs. – Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, USNM.

minuta Williston, 1895: 108. Type-locality: “Mexico”. Distr. – U. S. A. (Georgia, New Mexico), Mexico. Refs. – Williston, 1901: 307, pl. 5, fig. 19; Aldrich, 1905: 255 (cat.); Kertész, 1909: 96 (cat.); Fattig, 1945: 18; Hull, 1962 (1): 118, figs. 54, 514, 936, 945; Martin & Wilcox, 1965: 385 (cat.); Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Lectotype BMNH.

nemacula Martin, 1966a: 549, figs. 7-8. Type-locality: Mexico, Oaxaca, 17 miles east of Juchitán. Distr. – Mexico (Oaxaca). Refs. – Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, KU.

podexargenteus Enderlein, 1914b: 167 (as *podex-argenteus*). Type-locality: “Costa Rica”. Distr. – Costa Rica. Refs. – Hull,

1962 (1): 118; Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, IZPAN.
triangulata Martin, 1966a: 550, figs. 3-4. Type-locality: Mexico, Puebla, 3 miles west, 1 mile south of Izúcar de Matamoros. Distr. – Mexico (Puebla). Refs. – Martin & Papavero, 1970: 5 (cat.); Artigas & Papavero, 1990: 42. Type, KU.

Subfamily Trigonomiminae Enderlein, 1914

Trigonomimini Enderlein, 1914b: 166. Type-genus, *Trigonomima* Enderlein, 1914.
Xenomyzini G. H. Hardy, 1948: 116. Type-genus, *Xenomyza* Wiedemann, 1817 = *Damalis* Fabricius, 1805.
Damalini Hull, 1962 (1): 51. Type-genus, *Damalis* Fabricius, 1805.
Trigonomiminae Papavero, 1973a: 233, 248.
Trigomiminae Lehr, 1988: 214.

Genus **Haplopogon** Engel, 1930

Haplopogon Engel, 1930. Type-species, *Haplopogon nudus* Engel, 1930 (orig. des.).

Ref. – Martin, 1955 (rev.).

bullatus (Bromley), 1934a: 89 (*Holcocephala*). Type-locality: U. S. A., Texas, Bexar County. Distr. – U. S. A. (Texas), Mexico (Coahuila). Ref. – Pritchard, 1941b: 351, fig. 3 (n. comb.); Hull, 1962 (1): 68; Martin & Wilcox, 1965: 373 (cat.); Wilcox, 1966b: 100 (in key), fig. 1. Type, Texas Agricultural Experiment Station.
latus (Coquillett), 1904c: 33 (*Holopogon*). Type-locality: U. S. A., Texas, Brownsville. Distr. – U. S. A. (Texas), Mexico (Nuevo León). Refs. – Martin, 1955a: 315; Martin & Wilcox, 1965: 373 (cat.); Wilcox, 1966b: 101 (in key), fig. 1. Type, USNM.
lautus Back, 1909: 314 (*Holopogon*), Bromley, 1934: 98 (*Holopogon*), Pritchard, 1941b: 351, fig. 2, Hull, 1962 (1): 68, error.

Genus **Holcocephala** Jaennicke, 1867

Discocephala Macquart, 1838: 50 (1839: 166) (preocc. Jaennicke, 1932). Type-species, *Discocephala rufiventris* Macquart, 1838 (sub. des., Coquillett, 1910: 534) = *Holcocephala abdominalis* (Say, 1823).
Holcocephala Jaennicke, 1867: 359 (nom. nov. for *Discocephala* Macquart, 1838). Type-species, *Discocephala rufiventris* Macquart, 1838 (aut.) = *Holcocephala abdominalis* (Say, 1823).
Helcocephala Loew, 1874: 377, error.
Holocephala Williston, 1891: 72 (cat.), error.
Arthriticopus Enderlein, 1914b: 161. Type-species, *Arthriticopus nodosipes* Enderlein, 1914 (orig. des.). Ref. – Martin & Papavero, 1970: 6 (syn.).
Rhipidocephala Hermann, 1926b: 174. Type-species, *Discocephala analis* Macquart, 1846 (orig. des.).

affinis (Bellardi), 1861: 86, pl. 1, fig. 13 (*Discocephala*). Type-locality: “Mexico”. Distr. – Mexico, Honduras, Costa Rica. Refs. – Osten Sacken, 1874: 182; Aldrich, 1905: 261 (cat.); Kertész, 1909: 90 (cat.); Pritchard, 1938a: 13 (in key), 18; Hull, 1962 (1): 64; Martin & Papavero, 1970: 6 (cat.); Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipterol). Type, TORO.
agalla (Walker), 1849: 361 (*Discocephala*). Type-locality: “Venezuela”. Distr. – Venezuela (Aragua, Miranda). Refs. – Kertész, 1909: 92 (as syn. of *oculata* Fabricius, 1805); Martin & Papavero, 1970: 6 (cat.; as syn. of *oculata* Fabricius, 1805); Ayala & Baez, 1983: 1-4, figs. 1-5 (revalidation of sp.). Type, BMNH.
analis (Macquart), 1846: 197 (1846: 69) (*Discocephala*). Type-locality: “Colombia”. Refs. – Loew, 1860: 72 (*Discocephala*); Williston, 1891: 72 (*Holocephala*); Hermann, 1926b: 176 (*Rhipidocephala*); Hull, 1962 (1): 65 (*Rhipidocephala*); Martin & Papavero, 1970: 6 (cat.). Type, lost.
?apicalis Walker, 1854: 497 (*Discocephala*). Type-locality: “West coast of America”. Refs. – Williston, 1891: 72 (*Holocephala*); Bromley, 1946: 106 (cat.); Hull, 1962 (1): 64. Type, BMNH.
bechyneorum Ayala, 1982: 24, figs. 3-4, 17-18. Type-locality: Venezuela, Bolívar, km 132 de la carretera El Dorado, Santa Elena de Uairé, 1200 m. Distr. – Venezuela (Bolívar). Type, IZAUCV.
?coriacea (Wiedemann), 1821: 230 (*Dasypogon*). Type-locality: “Brazil”. Distr. – Brazil (Rio de Janeiro, São Paulo, Santa Catarina). Refs. – Wiedemann, 1828: 411 (*Dasypogon*); Walker, 1854: 446 (*Dasypogon*); Schiner, 1866: 680 (?*Discocephala*); Williston, 1891: 67 (cat.; ?*Holocephala*); Kertész, 1909: 91 (cat.; ?*Holcocephala*); Bromley, 1946:

- 106 (cat.); Carrera, 1958a: 142; Hull, 1962 (1): 64; Martin & Papavero, 1970: 6 (cat.).
?femorata Hermann, 1926b: 173. Type-locality: Brazil, Rio Grande do Sul, Porto Alegre (Teresópolis). Refs. – Carrera, 1945: 159 (preys); Bromley, 1946: 106 (cat.); Carrera, 1958a: 142 (syn.); Hull, 1962 (1): 64. Type, BUD.
- curvicosta** Carrera, 1958a: 142, figs. 1, 8. Type-locality: Brazil, São Paulo, Salesópolis (Estação Biológica de Boracéia). Distr. – Brazil (São Paulo). Ref. – Martin & Papavero, 1970: 6 (cat.). Type, MZUSP.
- deltoidea** (Bellardi), 1861: 85, pl. 1, fig. 12 (*Discocephala*). Type-locality: “Mexico”. Distr. – Mexico (Veracruz, Zacatecas), Guatemala, Honduras, Costa Rica. Refs. – Osten Sacken, 1874: 182; Aldrich, 1905: 261 (cat.); Kertész, 1909: 91 (cat.); Pritchard, 1938a: 13 (in key), 19; Hull, 1962 (1): 64; Martin & Papavero, 1970: 6 (cat.); Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipterol). Type, TORO.
- dimidiata** Hermann, 1926b: 159 (in key), 170. Type-locality: Peru, mouth of Pachitea river, 150 m. Distr. – Venezuela (Aragua, Táchira), Peru. Refs. – Hull, 1962 (1): 64; Carrera & Machado-Allison, 1963: 246; Martin & Papavero, 1970: 7 (cat.); Lamas M., 1972: 305 (cat.). Syntypes, MUN, TP SMTD [cf. Kallweit, 1992: 195].
- divisa** (Walker), 1860: 279 (*Discocephala*). Type-locality: “Mexico”. Distr. – Mexico (Veracruz). Refs. – Osten Sacken, 1874: 182; Kertész, 1909: 91 (cat.); Pritchard, 1938a: 13 (in key), 18; Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.); Papavero & Ibáñez Bernal, 2001: 95 (hist. Mexican dipterol.). Type, lost.
longipennis Bellardi, 1861: 86, pl. 1, fig. 14 (*Discocephala*). Type-locality: Mexico, Veracruz, Jalapa. Refs. – Osten Sacken, 1897: 182, 1887: 171; Aldrich, 1905: 261 (cat.); Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipterol). Type, TORO.
- fernandezii** Ayala, 1982: 25, figs. 7-8, 13, 19-20. Type-locality: Venezuela, Miranda, San Anonio de los Altos, Los Moralitos, 900 m. Distr. – Venezuela (Aragua, Miranda). Type, IZAUCV.
- fimbriata** Hermann, 1926b: 160 (in key), 167. Type-locality: Brazil, Rio Grande do Sul. Distr. – Brazil, Rio Grande do Sul. Refs. – Bromley, 1946: 106 (cat.); Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.). Syntypes, MUN, NMW.
- flavipes** Hermann, 1926b: 160 (in key), 177 (*Rhipidocephala*). Type-locality: “Mexico”. Ref. – Carrera, 1958a: 141 (misident.; as syn. of *obscuripennis* Enderlein, 1924). Syntypes, ZSBS, SMTD [cf. Kallweit, 1992: 194].
- indigena** Scarbrough & Perez-Gelabert, 2006: 84, figs. 112-117, 123. Type-locality: Dominican Republic, La Vega, road-256 La Comparación-Los Tablones, Parque Nacional Armando Bermúdez. Distr. – Dominican Republic (Barahona, Elias Piña, Independencia, La Vega, Pedernales, San Cristóbal). Type, USNM.
- inornata** (Rondani), 1848: 90 (*Discocephala*). Type-locality: “Brazil” [Pará, Belém; cf. Papavero, 1973d: 343]. Refs. – Williston, 1891: 72 (cat.; *Holocephala*); Kertész, 1909: 91 (cat.); Bromley, 1946: 106 (cat.); Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.). Type, MZUN.
- luteipes** Hermann, 1926b: 164. Type-locality: Peru, Urubamba river, 300 m; Bolivia, La Paz, Mapiri, Sarampiuní, 700 m. Distr. – Peru, Bolivia. Refs. – Carrera, 1958a: 142 (misident; as syn. of *obscuripennis* Enderlein, 1914); Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.); Lamas M., 1972: 305 (cat.); Fisher, 1984: 27. Syntypes, MUN.
- macula** (Rondani), 1848: 90 (*Discocephala*). Type-locality: “Brazil” [Pará, Belém; cf. Papavero, 1973d: 343]. Refs. – Williston, 1891: 72 (cat.; *Holocephala*); Bromley, 1946: 106 (cat.); Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.). Type, MZUN.
- matteii** Ayala, 1982: 26, figs. 9-10, 14, 21-22. Type-locality: Venezuela, Bolívar, El Pao, 500 m. Distr. – Venezuela (Bolívar). Type, IZAUCV.
- minuta** (Bellardi), 1861: 83 (*Discocephala*). Type-locality: Mexico, Tuxpango near Orizaba. Distr. – Mexico (Veracruz). Refs. – Osten Sacken, 1874: 182; Kertész, 1909: 92 (cat.; as syn. of *nitida* Wiedemann, 1830); Martin & Papavero, 1970: 7 (cat.); Papavero & Ibáñez Bernal, 2001: 108 (hist. Mexican dipterol). Syntypes, TORO (part of ST with V-shaped facial marks are either *H. oculata* or *H. nitida*; those without V-shaped marks, *H. minuta*).
- mogiana** Carrera, 1955a: 112, figs. 7-8. Type-locality: Brazil, São Paulo, Mogi das Cruzes. Distr. – Brazil (Rio de Janeiro, São Paulo, Paraná). Refs. – Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.). Type, MZUSP.
- monticola** Ayala, 1982: 27, figs. 5-6, 12, 23-24. Type-locality: Venezuela, Aragua, Vía Tiara, Campamento Rangel, 1200 m. Distr. – Venezuela (Aragua, Miranda). Type, IZAUCV.
- nigrita** (Fabricius), 1805: 152 (*Dioctria*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Brazil (São Paulo). Refs. – Wiedemann, 1821: 180 (*Dioctria*), 1828: 366 (*Dioctria*); Walker, 1854: 497 (*Discocephala*); Schiner, 1868: 69; Williston, 1891: 69 (cat.; *Holocephala*); Kertész, 1909: 111 (cat.); Carrera, 1958a: 144; Hull, 1962 (1): 29 (*Dioctria*); Martin & Papavero, 1970: 7 (cat.). Syntypes, COPN, Kiel [cf. Zimsen, 1964: 457].
?alboatra Walker, 1851: 100, pl. 3, fig. 8 (*Discocephala*). Type-locality: “Brazil”. Refs. – Walker, 1854: 497 (*Discocephala*); Williston, 1891: 72 (*Holocephala*); Kertész, 1909: 91 (cat.); Bromley, 1946: 106 (cat.); Hull, 1962 (1): 64; Carrera, 1958a: 144 (syn). Type, lost.
- nitida** (Wiedemann), 1830: 643 (*Dasypogon*). Type-locality: “Mexico [Deppe; cf. Papavero, 1971e: 103-107; Papavero & Ibáñez Bernal, 2001: 69-71]. Refs. – Walker, 1854: 496, 503 (*Discocephala*); Bellardi, 1861: 84 (*Discocephala*); Schiner, 1866: 681 (*Discocephala*); Osten Sacken, 1874: 182; Williston, 1901: 308; Aldrich, 1905: 261 (cat.); Kertész, 1909: 92 (cat.); Pritchard, 1938a: 13 (in key), 16; Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.); Papavero & Ibáñez Bernal, 2001: 75 (hist. Mexican dipterol.).
- ?interlineata* Walker, 1860: 279 (*Discocephala*). Type-locality: “Mexico”. Refs. – Osten Sacken, 1874: 182; Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.). Type, lost.
- ?minuta* Bellardi, 1861: 83 (*Discocephala*), part (ST with V-shaped facial marks).
- nodosipes** (Enderlein), 1914b: 161, fig. 4 (*Arthriticopus*). Type-locality: Colombia: Cundinamarca, Fusagasugá, and Caldas,

- Victoria (Hacienda Pehlke), 900 m [cf. Papavero, 1973d: 299]. Distr. – Colombia. Refs. – Hull, 1962 (1): 64 [*Holcocephala (Arthriticopus)*]; Martin & Papavero, 1970: 7 (n. comb.). Syntypes, IZPAN.
- obscuripennis** Enderlein, 1914b: 162, fig. 5. Type-locality: Brazil, Santa Catarina. Distr. – Brazil (Santa Catarina). Refs. – Bromley, 1946: 106 (cat.); Carrera, 1958a: 142; Hull, 1962 (1): 64; Martin & Papavero, 1970: 7 (cat.). Syntypes, IZPAN.
- oculata** (Fabricius), 1805: 151 (*Diocetria*). Type-locality: “South America (Smidt)” [i. e., Guyana, between rivers Demerara and Essequibo; cf. Papavero, 1971e: 21]. Distr. – Venezuela, Guyana, Peru, Brazil, Bolivia, Argentina. Refs. – Wiedemann, 1821: 230 (*Dasypogon*), 1828: 412 (*Dasypogon*); Macquart, 1838: 50 (*Discocephala*); Walker, 1854: 497 (*Discocephala*); Schiner, 1866: 681 (*Discocephala*); Williston, 1891: 72 (cat.; *Holocephala*); Kertész, 1909: 92 (cat.); Hermann, 1926b: 159 (in key), 161; Engel, 1930: 472; Curran, 1934c: 332; Pritchard, 1938a: 13 (in key), 17; Carrera, 1958a: 141; Carrera & d’Andretta, 1961: 68 (prey); Hull, 1962 (1): 64; Carrera & Machado-Allison, 1963: 245; Martin & Papavero, 1970: 7 (cat.); Lamas M., 1972: 305 (cat.); Fisher, 1984: 27; LaPierre, 2000: 643 (prey selection and diurnal activity; Costa Rica). Syntypes, COPN, Kiel [cf. Zimsen, 1964: 467].
- ?*minuta* Bellardi, 1861: 83 (*Discocephala*), part (ST with V-shaped facial marks).
- fasciventris* Hermann, 1926b: 159 (in key), 163 (as *oculata* var.). Type-locality: Peru, rivers Ucayali and Urubamba; Junín, Chanchamayo; and Bolivia, La Paz, Sarampíuní. Syntypes, MUN.
- flaviventris* Engel, 1930: 472 (as *oculata* var.; error for *fasciventris* Hermann).
- pardalina** Hermann, 1926b: 166. Type-locality: “Peru”. Refs. – Hull, 1962 (1): 64; Martin & Papavero, 1970: 8 (cat.); Lamas M., 1972: 305.
- pectinata** Carrera, 1955a: 115, figs. 9-10. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. – Brazil (São Paulo, Santa Catarina). Refs. – Hull, 1962 (1): 64; Martin & Papavero, 1970: 8 (cat.). Type, BMNH.
- pennipes** Hermann, 1926b: 159 (in key), 168. Type-locality: Peru, Callanga. Refs. – Hull, 1962 (1): 64; Martin & Papavero, 1970: 8 (cat.); Lamas M., 1972: 305 (cat.). Type, BUD.
- peruviana** Hermann, 1926b: 160 (in key), 172. Type-locality: Peru, Lima, Callanga. Refs. – Hull, 1962 (1): 64; Martin & Papavero, 1970: 8 (cat.). Type, BUD.
- rufithorax** (Wiedemann), 1828: 410 (*Dasypogon*). Type-locality: “Brazil” [Freyreiss; cf. Papavero, 1971e: 57-58; and Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-102]. Distr. – Peru, Brazil (Pará), Bolivia. Refs. – Macquart, 1838: 50 (*Discocephala*); Walker, 1854: 496 (*Discocephala*); Schiner, 1866: 680 (*Discocephala*); Williston, 1891: 72 (cat.; *Holocephala*); Kertész, 1909: 92 (cat.); Hermann, 159 (in key), 1924b: 160. Bromley, 1946: 106 (cat.); Carrera, 1960a: 150; Hull, 1962 (1): 64; Martin & Papavero, 1970: 8 (cat.); Lamas M., 1972: 305 (cat.). Syntypes, BZM (no. 2244; cf. Schumann, 1973: 153), SENCK [cf. Geller-Grimm, 1999: 214].
- scopifera** (Schiner), 1868: 161 (*Discocephala*). Type-locality: “South America”. Refs. – Williston, 1891: 72 (cat.; *Holocephala*); Kertész, 1909: 92 (cat.); Hermann, 1926b: 159 (in key), 171; Hull, 1962 (1): 64, fig. 44; Martin & Papavero, 1970: 8 (cat.); Lamas M., 1972: 305 (cat.). Type, NMW.
- spinipes** Hermann, 1926b: 158 (in key), 165. Type-locality: “Venezuela”. Distr. – Mexico (Veracruz), Venezuela. Refs. – Pritchard, 1938a: 13; Hull, 1962 (1): 66; Martin & Papavero, 1970: 8 (cat.). Type, NMW.
- stylata** Pritchard, 1938a: 13 (in key), 14. Type-locality: Mexico, Oaxaca, Oaxaca. Distr. – Mexico (Hidalgo, Oaxaca, San Luís Potosí). Refs. – Hull, 1962 (1): 64; Martin & Papavero, 1970: 8 (cat.). Type, MIN.
- uruguayensis** Lynch Arribálzaga, 1882: 133. Type-locality: Uruguay, Montevideo. Refs. – Lynch Arribálzaga, 1883a: 143 (cat.); Williston, 1891: 72 (cat.; *Holocephala*); Kertész, 1909: 93 (cat.); Hermann, 1926b: 159 (in key), 163 (as *oculata* var.); Hull, 1962 (1): 64; Martin & Papavero, 1970: 8 (cat.).
- vallestris** Ayala, 1982: 22, figs. 1-2, 11, 15-16. Type-locality: Venezuela, Guárico, La Puerta, río Guárico. Distr. – Venezuela ((Aragua, Barinas, Carabobo, Guárico, Miranda, Portuguesa, Yaracuy, Zulia)). Type, IZAUCV.
- ?**vicina** (Macquart), 1838: 1838: 26 (1839: 142) (*Diocetria*). Type-locality: “Brazil and Guadeloupe”. Refs. – Bromley, 1946: 106 (cat.; *Diocetria*); Hull, 1962 (1): 29 (*Diocetria*); Martin & Papavero, 1970: 8 (cat.). Syntypes, lost.
- vittata** (Walker), 1837: 338 (*Xylophagus*). Type-locality: Brazil, São Paulo. Ref. – Martin & Papavero, 1970: 8 (n. comb.). Type, BMNH.

Genus **Meliponomima** Artigas & Papavero, 1989

- Meliponomima** Artigas & Papavero, 1989: 37. Type-species, *Meliponomima martensis* Artigas & Papavero, 1989 (orig. des.).
- martensis** Artigas & Papavero, 1989: 38, figs. 3-5. Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (Dona Marta). Distr. – Brazil (Rio de Janeiro). Type, MZUSP.

Genus **Orrhodops** Hull, 1958

- Orrhodops** Hull, 1958c: 324. Type-species, *Damalis americana* Curran, 1930 (orig. des.).

americanus (Curran), 1930b: 5 (*Damalis*). Type-locality: U. S. A., Arizona, Sierras, Back Dyke Prospect. Distr. – U. S. A. (Arizona), Mexico. Refs. – Hull, 1958c: 324 (n. comb.), 1962 (1): 58, figs. 27, 838, 847, 1656, 1657, 1922, 2058; Martin & Wilcox, 1965: 381 (cat.). Type, AMNH.

occidentalis (Williston), 1901: 309, pl. 5, fig. 21 (*Damalis*). Type-locality: Mexico, Guerrero, Tepetlapa, and Jalisco, Santiago Escuintla. Distr. – México. Refs. – Hull, 1962 (1): 55 (*Damalis*), 58 (probably belongs to *Orrhodops*); Martin & Papavero, 1970: 8 (n. comb.); Papavero & Ibáñez Bernal, 2003: 175 (hist. Mexican dipterol). Syntypes, BMNH.

Genus **Seabramyia** Carrera, 1960

Seabramyia Carrera, 1960a: 147. Type-species, *Seabramyia tijucana* Carrera, 1958 (orig. des.).

tijucana Carrera, 1960a: 148, figs. 1-4. Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (Alto da Boa Vista and Floresta da Tijuca). Distr. – Brazil (Rio de Janeiro). Refs. – Hull, 1962 (2): 596; Martin & Papavero, 1970: 8 (cat.; *Holcocephala* (*Seabramyia*)). Type, MNRJ.

Nomina nuda (Trigonomiminae)

pullata Burmeister, 1861: 170 (*Discocephala*).

Unplaced genus and species of Asilidae

Genus †**Arariopogon** Grimaldi, 1990

†**Arariopogon** Grimaldi, 1990: 173. Type-species, *Arariopogon axelrodi* Grimaldi, 1990 (orig. des.).

†**axelrodi** Grimaldi, 1990: 173, figs. 21-24, 26-28. Type-locality: Brazil, Chapada do Araripe (Santa Formation, Lower Cretaceous). Type, AMNH.

†sp. Grimaldi, 1990: 174, figs. 25, 29-30. Brazil, Chapada do Araripe (Santa Formation, Lower Cretaceous).

Unrecognized and unplaced species of Asilidae (asterisk (*) indicates that the type is lost)

***albibilosus** Macquart, 1846: 220 (1846: 92) (*Asilus*). Type-locality: “Brazil” and Colombia, Bogotá. Refs. – Walker, 1855: 701 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 250 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 88 (cat.; *Asilus*).

antiphus Walker, 1849: 443 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 704 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 250 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 88 (cat.; *Asilus*).

atripes Fabricius, 1805: 170 (*Dasypogon*). Type-locality: “West Indies”. Ref. – Wiedemann, 1821: 195 (*Asilus*), 1828: 455 (*Asilus*); Walker, 1855: 695 (*Asilus*); Schiner, 1866: 685 (?*Asilus*); Aldrich, 1905: 282 (cat.; *Asilus*); Kertész, 1909: 251 (cat.; ?*Asilus*). Type, COPN [cf. Zimsen, 1964: 470].

audouinii Macquart, 1838: 155 (1839: 271) (*Gonytes*). Type-locality: unknown. Refs. – Walker, 1855: 770 (*Leptogaster*); Schiner, 1867: 357 (*Leptogaster*); Osten Sacken, 1878: 230 (cat.; *Leptogaster*); Williston, 1891: 68 (*Leptogaster*); Kertész, 1909: 208 (cat.; *Leptogaster*).

aurarius Wiedemann, 1821: 288 (*Dasypogon*). Type-locality: “Brazil”. Refs. – Wiedemann, 1828: 410 (*Dasypogon*); Walker, 1854: 442 (*Dasypogon*); Schiner, 1866: 680 (*Dasypogon*); Williston, 1891: 67 (*Dasypogon*); Kertész, 1909: 125 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 88 (cat.; *Dasypogon*). Type, BZM [no. 543; cf. Schumann, 1973: 89].

baletus Walker, 1849: 427 (*Asilus*). Type-locality: “Venezuela”. Refs. – Walker, 1855: 704 (*Asilus*); Williston, 1891: 90 (cat.;

- Asilus*); Kertész, 1909: 251 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 88 (cat.; *Asilus*).
***caeruleiventris** Macquart, 1846: 221 (1846: 93) (*Asilus*). Type-locality: "Colombia". Refs. – Walker, 1855: 701 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 253 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 88 (cat.; *Asilus*).
cepphicus Say, 1829: 158 (1852: 354) (*Dasypogon*). Type-locality: "Mexico". Refs. – Walker, 1854: 503 (*Dasypogon*); Osten Sacken, 1874: 185; Aldrich, 1905: 267 (cat.; *Dasypogon*); Kertész, 1909: 126 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 88 (cat.; *Dasypogon*).
chilensis Macquart, 1850: 372 (1850: 68) (*Dasypogon*; preocc. Macquart, 1850). Type-locality: "Chile". Refs. – Blanchard, 1852: 367 (*Dasypogon*); Philippi, 1865: 685 (*Dasypogon*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 127 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 88 (cat.; *Dasypogon*).
cinereum Bigot, 1878: 432 (*Bathypogon*). Type-locality: "Colombia". Refs. – Hull, 1962 (1): 150 (*Bathypogon*); Martin & Papavero, 1970: 18 (cat.; *?Creolestes*). Type, OXF.
colombiae Macquart, 1838: 148 (1848: 264) (*Asilus*). Type-locality: "Colombia". Refs. – Walker, 1855: 707 (*Asilus*); Kertész, 1909: 253 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 88 (cat.; *Asilus*). Type, MHNKP.
columbiae Williston, 1891: 90 (cat.; *Asilus*), error.
costalis Lynch Arribálzaga, 1880: 29 (*Dasypogon*). Type-locality: Argentina, Buenos Aires. Refs. – Lynch Arribálzaga, 1883a: 143 (cat.; *Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 127 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 88 (cat.; *Dasypogon*).
cuyanus Lynch Arribálzaga, 1880: 175 (*Asilus*). Type-locality: Argentina, Mendoza. Refs. – Lynch Arribálzaga, 1883a: 151 (cat.; *?Eutolmus*), 1883: 80, 85; Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 255 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 88 (cat.; *Asilus*).
***dorsalis** Macquart, 1848: 180 (1848: 20), pl. 1, fig. 12 (*Dasypogon*; preocc. Wiedemann, 1824). Type-locality: "Rio Negro". Refs. – Walker, 1854: 439 (*Dasypogon*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 127 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 88 (cat.; *Dasypogon*).
eburnus Walker, 1849: 332 (*Dasypogon*). Type-locality: Brazil, Pará [i. e., Pará, Belém]. Refs. – Walker, 1854: 432 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 65 (cat.; *Heteropogon*); Bromley, 1946: 105 (cat.; *Heteropogon*); Martin & Papavero, 1970: 20 (cat.; *?Heteropogon*).
eburneus Hull, 1962 (1): 194 (*Heteropogon*), error.
flavipennis Macquart, 1846: 192 (1846: 64), pl. 7, fig. 4 (*Dasypogon*; preocc. Wiedemann, 1828). Type-locality: "Colombia" (as Nouvelle Grenade). Refs. – Walker, 1854: 435 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 128 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 89 (cat.; *Dasypogon*).
flavipennis Wiedemann, 1828: 376 (*Dasypogon*). Type-locality: "Brazil" [Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-102]. Refs. – Walker, 1854: 444 (*Dasypogon*); Schiner, 1866: 677 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 128 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 89 (cat.; *Dasypogon*). Syntypes, BZM [no. 568; cf. Schumann, 1973: 111].
fraternus Macquart, 1846: 191 (1846: 63) (*Dasypogon*). Type-locality: Brazil, Minas Gerais. Ref. – Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 89 (cat.; *Dasypogon*).
gamaxus Walker, 1851: 142 (*Asilus*). Type-locality: "Colombia". Refs. – Walker, 1855: 703 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 256 (cat.; *Asilus*); Martin & Papavero, 1970: 89 (cat.; *Asilus*).
gammaeus Hull, 1962 (2): 545 (*Asilus*), error.
herdonius Walker, 1851: 140 (*Asilus*). Type-locality: "South America". Refs. – Walker, 1855: 702 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 257 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 89 (cat.; *Asilus*).
imitator Lynch Arribálzaga, 1883b: 79 (*Asilus*). Type-locality: Argentina, Río Quinto. Refs. – Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 257 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 89 (cat.; *Asilus*).
inamatus Walker, 1860: 283 (*Asilus*). Type-locality: "Mexico". Refs. – Aldrich, 1905: 282 (cat.; *Asilus*); Kertész, 1909: 257 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 89 (cat.; *Asilus*); Papavero & Ibáñez Bernal, 2001: 96 (hist. Mexican dipterol.).
inauratus Aldrich, 1905: 282 (*Asilus*), error.
***lebasii** Macquart, 1838: 148 (1848: 264) (*Asilus*). Type-locality: "Colombia". Refs. – Macquart, 1846: 93 (*Asilus*); Walker, 1855: 703 (*Asilus*); Williston, 1891: 90 (cat.; *Asilus*); Kertész, 1909: 258 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 89 (cat.; *Asilus*).
***lebasii** Macquart, 1838: 38 (1839: 154) (*Dasypogon*). Type-locality: "Colombia". Refs. – Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 89 (cat.; *Asilus*).
***luctuosus** Macquart, 1838: 39 (1839: 155) (*Dasypogon*). Type-locality: unknown. Refs. – Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 89 (cat.; *Dasypogon*).
***megastylus** Philippi, 1865: 697 (*Asilus*). Type-locality: Chile, Santiago and Choapa, Illapel. Refs. – Kertész, 1909: 260 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*).
meastylus Williston, 1891: 91 (cat.; *Asilus*), error.
melanacrus Wiedemann, 1828: 72 (*Asilus*). Type-locality: "Brazil" [Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-102]. Refs. – Walker, 1855: 638 (*Eanax*); Schiner, 1866: 687 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 260 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*).

Type, BZM [no. 2015; cf. Schumann, 1973: 132].

- mellipes** Wiedemann, 1828: 452 (*Asilus*). Type-locality: Brazil [Sello; cf. Papavero, 1971e: 56-60, 69-80, 99-102]. Refs. – Walker, 1855: 706 (*Asilus*); Schiner, 1866: 685 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 260 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*). Type, BZM [no. 2044; cf. Schumann, 1973: 132].
- ***mexicanus** Macquart, 1846: 196 (1846: 68), pl. 6, fig. 10 (*Dasypogon*). Type-locality: “Mexico”. Refs. – Walker, 1854: 428 (*Dasypogon*); Osten Sacken, 1874: 185; Aldrich, 1905: 76 (cat.; *Dasypogon*); Kertész, 1909: 130 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 90 (cat.; *Dasypogon*); Papavero & Ibáñez Bernal, 2001: 84 (hist. Mexican dipterol.).
- nigrinus** Macquart, 1848: 190 (1848: 30), pl. 3, fig. 5 (*Asilus*; preocc. Wiedemann, 1821). Type-locality: “Brazil”. Refs. – Walker, 1855: 702 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 261 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 478 (*Nerax*), 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*). Type, OXF.
- nigrostriata** Engel, 1930: 465, figs. 6-7 (*Heligmoneura*). Type-locality: Argentina, Formosa, Misión Tacaagle and San José. Refs. – Hull, 1962 (2): 590 (*Neomochtherus*); Martin & Papavero, 1970: 90 (cat.).
- ocrealis** Rondani, 1863: 45 (*Philammosius*). Type-locality: “Chile”. Refs. – Williston, 1891: 69 (cat.; *Philammosius*); Hull, 1962 (1): 106 (*Stichopogon*); Martin & Papavero, 1970: 5 (cat.; ?*Stichopogon*). Type, MZUN.
- ochrealis* Kertész, 1909: 84 (cat.; *Stichopogon*), emend.
- parvus** Rondani, 1850b: 369 (789) (*Dasipogon*). Type-locality: “Equatorial America”. Refs. – Osten Sacken, 1891: 419 (*Dasypogon*); Kertész, 1909: 130 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 90 (cat.; *Dasypogon*). Type, MZUN.
- phalma** Walker, 1849: 331 (*Dasypogon*). Type-locality: unknown. Refs. – Walker, 1854: 486 (*Dasypogon*); Williston, 1891: 68 (*Dasypogon*); Kertész, 1909: 65 (cat.; *Heteropogon*); Hull, 1962 (1): 164 (*Heteropogon*).
- phalna* Martin & Papavero, 1970: 20 (?*Heteropogon*), error.
- ***poecilopus** Philippi, 1865: 697 (*Asilus*). Type-locality: Chile: Valparaíso, and Choapa, Illapel. Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 262 (cat.; *Asilus*); Bromley, 1932: 281 (*Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*).
- pumilus** Macquart, 1838: 37 (1839: 153) (*Dasypogon*). Type-locality: Brazil, northern São Paulo. Refs. – Walker, 1854: 434 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 130 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 90 (cat.; *Dasypogon*). Type, MNHNP.
- ***punctipennis** Macquart, 1838: 46 (1839: 166) (*Dasypogon*). Type-locality: “Chile”. Refs. – Blanchard, 1852: 365 (*Dasypogon*); Walker, 1854: 441 (*Dasypogon*); Philippi, 1865: 685 (*Dasypogon*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 130 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 90 (cat.; *Dasypogon*).
- rubiginipennis** Macquart, 1838: 45 (1839: 161) (*Dasypogon*). Type-locality: Brazil, northern São Paulo. Refs. – Walker, 1854: 439 (*Dasypogon*); Williston, 1891: 68 (cat.; *Dasypogon*); Kertész, 1909: 131 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 90 (cat.; *Dasypogon*). Type, MNHNP.
- rufipalpis** Macquart, 1838: 146 (1839: 262) (*Asilus*). Type-locality: Western part of Uruguay, along Uruguay river (as “depuis de l’embouchure de l’Uruguay jusqu’aux Missions”). Refs. – Walker, 1855: 705 (*Asilus*); Lynch Arribálzaga, 1880: 174 (*Mochtherus*), 1883a: 151 (cat.; *Neomochtherus*), 1883b: 16 (*Neomochtherus*); Williston, 1891: 89 (cat.; *Heligmoneura*); Kertész, 1909: 288 (cat.; *Heligmoneura*); Hull, 1962 (2): 590 (*Neomochtherus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*).
- rufiventris** Walker, 1854: 503 (*Dasypogon*; preocc. Wiedemann, 1821). Type-locality: “South America”. Ref. – Martin & Papavero, 1970: 90 (cat.; *Asilus*).
- rufiventris** Wiedemann, 1821: 223 (*Dasypogon*). Type-locality: “South America”. Refs. – Wiedemann, 1828: 388 (*Dasypogon*); Walker, 1854: 432 (*Dasypogon*); Schiner, 1866: 678 (*Dasypogon*); Williston, 1891: 76 (cat.; *Dasypogon*); Kertész, 1909: 131 (cat.; *Dasypogon*); Bromley, 1946: 107 (cat.; *Dasypogon*); Hull, 1962 (1): 228 (*Dasypogon*); Martin & Papavero, 1970: 90 (cat.; *Dasypogon*).
- ***sericans** Walker, 1857: 130 (*Asilus*). Type-locality: Brazil, Amazonia. Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 263 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*).
- ***servillei** Macquart, 1834: 313 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 703 (*Asilus*); Kertész, 1909: 263 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 90 (cat.; *Asilus*). *servillea* Williston, 1891: 91 (cat.; *Asilus*), error.
- ***tatius** Walker, 1851: 148 (*Asilus*). Type-locality: “Colombia”. Refs. – Walker, 1855: 702 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 264 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 91 (cat.; *Asilus*).
- ***tenuiventris** Macquart, 1855: 86 (1855: 66), pl. 2, fig. 11 (*Asilus*). Type-locality: “South America”. Refs. – Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 264 (cat.; *Asilus*); Hull, 1962 (2): 545 (*Asilus*); Martin & Papavero, 1970: 91 (cat.; *Asilus*).
- therevinus** Rondani, 1850b: 369 (*Asilus*). Type-locality: “Equatorial America”. Refs. – Osten Sacken, 1891: 421 (*Asilus*); Kertész, 1909: 264 (cat.; *Asilus*); Hull, 1962 (2): 546 (*Asilus*); Martin & Papavero, 1970: 91 (cat.; *Asilus*). Type, MZUN.
- tristis** Wiedemann, 1828: 470 (*Asilus*). Type-locality: “Brazil”. Refs. – Walker, 1855: 635 (*Erax*); Schiner, 1866: 687 (*Asilus*); Williston, 1891: 91 (cat.; *Asilus*); Kertész, 1909: 265 (cat.; *Asilus*); Bromley, 1946: 116 (cat.; *Asilus*); Hull, 1962 (2): 546 (*Asilus*); Martin & Papavero, 1970: 91 (cat.; *Asilus*). Type, BZM [no. 2042; cf. Schumann, 1973: 166].

References

- Adisoemarto, S. & D. M. Wood, 1975. The Nearctic species of *Dioctria* and six related genera (Diptera, Asilidae). *Quest. ent.* 11: 505-576, 139 figs.
- Agassiz, L., 1846. *Nomenclator zoologicus, nomina systematica generum animalium tam viventium quam fossilium, secundum ordinem alphabeticum disposita, adjectis auctoribus, libris in quibus reperiuntur, anno editionis, etymologia et familiis ad quas pertinent in variis classibus. Index.* Soloduri [= Solothurn].
- Alayo Dalmau, P. & I. García Ávila, 1983. *Lista anotada de los dipteros de Cuba*, Editorial Científico-Técnica, La Habana.
- Alcock, J., 1974. Observations on the behavior of *Mallophora faunrix* Osten Sacken (Diptera, Asilidae). *Pan-Pacific Entomologist* 50 (1): 68-72, 3 figs.
- Aldrich, J. M., 1905. A catalogue of North American Diptera. *Smithson. misc. Collns* 46 [2 (= publ. 1444)]: 1-680.
- Aldrich, J. M., 1923. New genera of two-winged flies of the subfamily Leptogastrinae of the family Asilidae. *Proc. U. S. natn. Mus.* 62: 1-6, 3 figs.
- Almeida, D. S., I. Castro & F. Bravo, 2006. Nova ocorrência de espécies de *Mallophora* Macquart (Asilidae, Diptera) no estado da Bahia, Brasil. *Sitientibus, Ciênc. biol.* 6 (4): 257-260.
- Arnaud, P. H., Jr., 1974. *Myolestes* Brèthes, 1904 (Diptera, Asilidae), a valid generic name and a senior homonym of *Myolestes* Matthew, 1909 (Mammalia: Nyctitheriidae). *Wasmann J. Biol.* 32 (2): 203-205.
- Arnett, R. H., Jr., 1985. *American insects. A handbook of the insects of America north of Mexico*, xiv + 850 pp. Van Nostrand Reinhold, New York.
- Artigas C., J. N., 1961. Diferenciación del género chileno "Deromyia" Philippi y el norteamericano [sic] "Diogmites" Loew (Diptera: Asilidae). *Gayana, Concepción* 2: 5-11.
- Artigas C., J. N., 1964. Revisión del género *Pronomopsis* Hermann con la descripción de una nueva especie (Diptera –Asilidae). *Gayana, Concepción* 12: 3-9, 14 figs.
- Artigas C., J. N., 1970a. Los asilidos de Chile (Diptera – Asilidae). *Gayana, Concepción* 17: 1-472, 504 figs.
- Artigas C., J. N., 1970b. *Lochmorhynchus leoninus* n. sp., una nueva especie de Asilidae para Chile y Perú (Diptera – Asilidae). *Boln Soc. Biol. Concepción* 42: 67-70, 4 figs.
- Artigas C., J. N., 1971. Las estructuras quitinizadas de la spermatheca [sic] y funda del pene de los asilidos y su valor sistemático a través del estudio por taxonomía numérica (Diptera, Asilidae). *Gayana, Concepción* 18: 1-105.
- Artigas C., J. N., 1973. *Alyssomyia quinquemaculata* y *Alyssomyia misera*, dos nuevas especies de Asilidae chilenos y clave para las especies del género (Diptera –Asilidae). *Revta chilena Ent.* 7: 123-129, 15 figs.
- Artigas C., J. N., 1974a. *Aymarasilus inti* n. sp., nuevo género y especie de asilido de Chile (Diptera – Asilidae). *Boln Soc. Biol. Concepción* 47: 227-232, 10 figs.
- Artigas C., J. N., 1974b. *Scylaticus carrascoi* n. sp. del altiplano chileno-peruano (Diptera – Asilidae). *Boln Soc. Biol. Concepción* 48: 5-9, 10 figs.
- Artigas C., J. N., 1974c. *Zoticus fitzroyi*, una nueva especie de asilido de la Patagonia argentina. *Revta chilena Ent.* 8: 1198-121, 7 figs.
- Artigas C., J. N., 1980. *Myaptex acutus* n. sp., asilido nuevo de Chile (Diptera – Asilidae). *Brenesia* 17: 365-370, 13 figs.
- Artigas, C., J. N., 1981. *Lochmorhynchus albinigrus* n. sp., nueva especie de asilido argentino (Diptera – Asilidae). *Boln Soc. Biol. Concepción* 52: 187-190, 7 figs.
- Artigas C., J. N., 1982. *Menexenus schlingeri* sp. n., nuevo asilido de Argentina y consideraciones sobre el género *Menexenus* Artigas, 1970 (Diptera – Asilidae). *Boln Soc. Biol. Concepción* 53: 19-23.
- Artigas C., J. N., 1983. *Apoxyria cymbafer* sp. n., nueva especie y primer registro del género para Chile (Diptera, Asilidae, Laphystiinae). *Boln Soc. Biol. Concepción* 54: 27-33.
- Artigas C., J. N., 1996. *Martintella*, a new name for *Martinella* Artigas & Papavero, 1995 (Diptera, Asilidae). *Boln Soc. Biol. Concepción* 67: 75.
- Artigas C., J. N. & A. O. Angulo, 1980. Revisión del género *Mallophora* Macquart por sistemática alfa y numérica (Diptera –Asilidae). *Gayana, Zool.* 43: 1-182.
- Artigas C., J. N. & M. Hengst, 1998. Asilidae, pp. 353-364, in Morrone, J. J. & S. Coscarón, eds., *Biodiversidad de artrópodos argentinos: Una perspectiva biotaxonómica*, 599 pp. Ediciones Sur, La Plata.
- Artigas C., J. N. & M. Hengst, 1999. Clave ilustrada para los géneros de asilidos argentinos. *Revta chilena Hist. nat.* 72 (1): 107-150.
- Artigas C., J. N., P. D. Lewis & L. E. Parra, 2005. Review and phylogeny of the genus *Tillobroma* Hull with the description of ten new species and its relation with the genus *Hypenetes* Loew (Diptera, Asilidae, Stenopogoninae). *Revta chilena Hist. nat.* 78 (3): 519-588.
- Artigas, J. N. & N. Papavero, 1988a. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. I. Key to subfamilies and subfamily Leptogastrinae Schiner. *Gayana, Zool.* 52(1-2): 95-114, 59 figs.
- Artigas, J. N. & N. Papavero, 1988b. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. II. Key to the genera of Dasypogoninae Macquart, with descriptions of new genera and species and new synonymies. *Gayana, Zool.* 52(3-4): 199-260, 216 figs.
- Artigas, J. N. & N. Papavero, 1989a. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. III. Key to the genera of Trigonomiminae Enderlein, with descriptions of a new genus and species. *Boln Soc. Biol. Concepción* 60: 35-41, 11 figs.
- Artigas, J. N. & N. Papavero, 1990. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. V. Subfamily Stichopogoninae G. H. Hardy. *Boln Soc. Biol. Concepción* 61: 1-10, 11 figs.

- Soc. Biol. Concepción* 61: 39-47.
- Artigas, J. N. & N. Papavero, 1991a. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VII.1. Subfamily Stenopogoninae Hull. A preliminary classification into tribes. *Gayana, Zool.* 55(2): 139-144.
- Artigas, J. N. & N. Papavero, 1991b. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VII.2. Subfamily Stenopogoninae Hull – Tribes Acronychini, Bathypogonini (with description of a new genus) and Ceraturgini, and a catalogue of the Neotropical species. *Gayana, Zool.* 55(3): 247-255, 12 figs.
- Artigas, J. N. & N. Papavero, 1991c. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VII.3. Subfamily Stenopogoninae Hull – Tribes Dioctriini and Echthodopini. *Gayana, Zool.* 55 (4): 261-266, 8 figs.
- Artigas, J. N. & N. Papavero, 1991d. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VII.4. Subfamily Stenopogoninae Hull – Tribe Enigmomorphini, with descriptions of three new genera and species and a catalogue of the Neotropical species. *Boln Soc. Biol. Concepción* 62: 27-53, 35 figs.
- Artigas, J. N. & N. Papavero, 1991e. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VII.5. Subfamily Stenopogoninae Hull – Tribe Tillobromini, with descriptions of three new genera and two new species and a catalogue of the Neotropical species. *Revta chilena Ent.* 19: 17-27, 17 figs.
- Artigas, J. N. & N. Papavero, 1991f. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VII.7. Subfamily Stenopogoninae Hull – Tribe Cyrtopogonini, with descriptions of four new genera and one new species and a catalogue of the Neotropical species. *Boln Soc. Biol. Concepción* 62: 55-81, 51 figs.
- Artigas, J. N. & N. Papavero, 1993. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VII.6. Subfamily Stenopogoninae Hull – Tribes Phellini, Plesiommatini, Stenopogonini and Willistoninini. *Gayana, Zool.* 57(2): 309-321, 12 figs.
- Artigas, J. N. & N. Papavero, 1995a. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.3. Subfamily Asilinae Leach – *Eichoichemus*-group, with the proposal of two new genera and a catalogue of the Neotropical species. *Gayana, Zool.* 59(1): 97-102, 8 figs.
- Artigas, J. N. & N. Papavero, 1995b. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.4. Subfamily Asilinae Leach, *Glaphyropyga*-group, with the proposal of two new genera and a catalogue of the Neotropical species. *Boln Soc. Biol. Concepción* 66: 11-33, 99 figs.
- Artigas, J. N. & N. Papavero, 1995c. The American genera of Asilidae (Diptera). Keys for identification with an atlas of female spermathecae and other morphological details. IX.5. Subfamily Asilinae Leach, *Lochmorhynchus*-group, with a catalogue of the Neotropical species. *Gayana, Zool.* 59(2): 131-144, 48 figs.
- Artigas, J. N. & N. Papavero, 1995d. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.7. Subfamily Asilinae Leach, *Proctacanthus*-group, with the proposal of a new genus and a catalogue of the Neotropical species. *Gayana, Zool.* 59(2): 145-160, 59 figs.
- Artigas, J. N. & N. Papavero, 1995e. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.8. Subfamily Asilinae Leach, *Eicherax*-group, with a catalogue of the Neotropical species. *Boln Soc. Biol. Concepción* 66: 35-42, 20 figs.
- Artigas, J. N. & N. Papavero, 1995f. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.9. Subfamily Asilinae Leach, *Myaptex*-group, with the proposal of two new genera and a catalogue of the Neotropical species. *Revta chilena Ent.* 22: 55-73, 73 figs.
- Artigas, J. N. & N. Papavero, 1995g. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.10. Subfamily Asilinae Leach, *Lecania*-group, with a catalogue of the Neotropical species. *Theoria. Ciencia, Arte y Humanidades*, Concepción 4: 33-56.
- Artigas, J. N. & N. Papavero, 1997a. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.1. Subfamily Asilinae Leach (including Apocleinae Lehr). Key to generic groups. *Argos Zool.*, São Paulo 34(2): 57-63.
- Artigas, J. N. & N. Papavero, 1997b. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.2. Subfamily Asilinae Leach – *Efferia*-group, with the proposal of five new genera and a catalogue of the Neotropical species. *Argos Zool.*, São Paulo 34(3): 65-95, 64 figs.
- Artigas, J. N. & N. Papavero, 1997c. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. IX.6. Subfamily Asilinae Leach, *Mallophora*-group, with a catalogue of the Neotropical species. *Argos Zool.*, São Paulo 34(4): 97-120, 67 figs.
- Artigas, J. N., N. Papavero & N. C. A da Costa, 1997. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VIII. Subfamily Laphystiinae G. H. Hardy, with descriptions of five new genera and species and a catalogue of the Neotropical species. *Argos Zool.*, São Paulo 34(1): 1-55, 134 figs.
- Artigas, J. N., N. Papavero & T. Pimentel, 1988. The American genera of Asilidae (Diptera): Keys for identification with

- an atlas of female spermathecae and other morphological details. IV. Key to the genera of Laphriinae Macquart (except tribe Atomosiini Hermann), with the description of three new tribes and five new species. *Bolm Mus. paraense Emílio Goeldi (Zool.)* 4(2): 211-256, 72 figs.
- Artigas, J. N., N. Papavero & A. L. Serra, 1991d. The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. VI. Tribe Atomosiini Hermann (Laphriinae), with descriptions of two new genera and three new species, and a catalogue of the Neotropical species. *Gayana, Zool.* 55 (1): 53-85, 104 figs.
- Artigas C., J. N. & L. E. Parra, 2006. Phylogeny of the genus *Alyssomyia* Hull, with descriptions of two new species (Diptera, Asilidae, Stenopogoninae, Enigmormorphini). *Stud. neotr. Fauna Environm.* 41 (2): 97-116.
- Artigas C., J. N. & E. Reyes, 1995. Aparato reproductor de las hembras de tres especies de asilidos chilenos, con énfasis especial en la estructura de las spermathecae [sic] (Diptera - Asilidae). *Boln Soc. Biol. Concepción* 44: 195-205, 15 figs.
- Ayala L., J. M., 1978. Contribución al estudio de la tribu Asilini (Diptera: Asilidae). *Revta Fac. Agron. Maracay* 26: 31-46, 8 figs.
- Ayala L., J. M., 1981. Contribución al estudio de los asilidos de Venezuela (Diptera- Asilidae). *Acta Macarao* (n. s.) 2: 51-59, 13 figs.
- Ayala L., J. M., 1982. El género *Holcocephala* Jaennicke (Diptera, Asilidae) en Venezuela. I. *Boln Ent. venezolana* (n. s.) 2 (3): 21-32.
- Ayala L., J. M., 1983. El género *Glaphyropyga* Schiner (Diptera, Asilidae), en Venezuela. *Boln Ent. venezolana* (n. s.) 3 (2): 5-20, 23 figs.
- Ayala L. & M. Baez, 1983. El género *Holcocephala* Jaennicke (Diptera, Asilidae) en Venezuela. II. *Boln Ent. venezolana* (n. s.) 3 (1): 1-4.
- Back, E. A., 1904. New species of North American Asilidae. *Canadian Entomologist* 36: 289-293.
- Back, E. A., 1909. The robber-flies of America north of Mexico, belonging to the subfamilies Leptogastrinae and Dasypogoninae. *Trans. American ent. Soc.* 35: 137-400, 11 pls.
- Baker, C. F., 1904. Reports on Californian and Nevadan Diptera. I. *Invertebrata pacifica* 1: 17-40.
- Baker, R. H., 1939. A list of robber flies from Coahuila, Mexico (Diptera: Asilidae). *Bull. Brooklyn ent. Soc.* 34: 167-170.
- Bambara, S. B., 1983. But bumble bees don't eat bees. *Gleanings in Bee Culture* 111 (3): 243.
- Banks, N., 1911. Four new species of Asilidae. *Canadian Entomologist* 43: 128-130.
- Banks, N., 1920. Descriptions of a few new Diptera. *Canadian Entomologist* 52: 65-67.
- Becker, T., 1919. Diptères. Brachycères, in *Mission du Service Géographique de l'Armée pour la mesure d'un arc méridien équatorial en Amérique du Sud*: 143-233. Paris.
- Becker, T., 1925. H. Sauter's Formosa-Ausbeute: Asilinae (III) (Dipt.). *Ent. Mitt.* 14: 62-85, 123-129, 240-250, 9 figs.
- Bellardi, L., 1861. *Saggio di ditterologia messicana. Parte II* 2: 99 pp., 2 pls. Torino. (Also publ. in *Mem. r. Accad. Sci. Torino* 21: 103-199, 2 pls., 1864).
- Bellardi, L., 1862. *Saggio di ditterologia messicana. Appendice*, 28 pp., 1 pl. Torino. [Also publ. in *Mem. R. Accad. Sci. Torino* 21: 200-225, 1 pl., 1864].
- Bezark, L. G., 1984. Description of the first Central American species of *Stenopogon* Loew (Diptera, Asilidae). *Pan-Pacific Entomologist* 60: 53-54, 4 figs.
- Bezzi, M., 1902. Neue Namen für einige Dipteren-Gattungen. *Ztschr. Hymenopterol. Dipterol.* 2: 190-192.
- Bezzi, M., 1908. Diagnoses d'espèces nouvelles de diptères d'Afrique. *Ann. Soc. ent. Belgique* 54: 347-388.
- Bezzi, M., 1909. De specie altera dipterorum generis *Townsendia* Willist. *Ann. Mus. natn. hung.* 7: 628-630, 1 fig.
- Bezzi, M., 1910a. Un nuovo genere di asilidi dell'America Centrale. *Boll. Labor. Zool. Portici* 4: 175-197.
- Bezzi, M., 1910b. Synonymie und systematischen Stellung einiger Dipteren. *Societas entomologica* 25 (17): 65-67.
- Bezzi, M., 1910c. Revisio systematica generis dipterorum *Stichopogon*. *Ann. Mus. natn. hung.* 8: 128-159.
- Bigot, J. M. F., 1852. Essai d'une classification générale et synoptique de l'ordre des insectes diptères. *Ann. Soc. ent. France* (2) 10: 471-489.
- Bigot, J. M. F., 1853. Note pour servir de complément et de correction à l'Essai d'une classification générale et synoptique de l'ordre des insectes diptères. *Ann. Soc. ent. France* (3) 1: 295-317.
- Bigot, J. M. F., 1857a. Diptères, pp. 328-349, in Sagra, R. de la, *Historia física, política y natural de la isla de Cuba* 7: 371 pp. Paris, "1856". (Also publ. in French, pp. 783-829, Paris, 829 pp., 1857).
- Bigot, J. M. F., 1857b. Essai d'une classification générale et particulière de l'ordre des insectes diptères. Ve. *Ann. Soc. ent. France* (3) 5: 517-564.
- Bigot, J. M. F., 1857c. Diptères nouveaux provenant du Chili. *Ann. Soc. ent. France* (3) 5: 277-308, pl. 6.
- Bigot, J. M. F., 1875. Diptères nouveaux ou peu connus. 4e. partie. V. Asilides exotiques nouveaux. *Ann. Soc. ent. France* (5) 5: 237-248.
- Bigot, J. M. F., 1878. Diptères nouveaux ou peu connus. 10e. partie. XV. Tribu des Asilidi. Curies des Laphridae et Dasypogonidae. *Ann. Soc. ent. France* (5) 8: 213-240, 401-446.
- Bigot, J. M. F., 1879. [Note: On synonymy of Asilidae]. *Ann. Soc. ent. France* 5 (9) (Bull.): lxvii-lxviii.
- Bigot, J. M. F., 1880. (Diagnoses de trois genres nouveaux de diptères). *Ann. Soc. ent. France* (5) 10 (Bull.): xlvi-xlviii.
- Bigot, J. M. F., 1889. [Note: On relationships of *Archilestes* and synonymy: *Pseudarchilestes* proposed]. *Ann. Soc. ent. France* (6) 9 (Bull.): clxxxiii.
- Bigot, J. M. F., 1890. [Note: Proposes the name *Pseudoarchilestes*]. *Wien. ent. Ztg* 9: 96.
- Blanchard, E., 1852. Orden IX. Dípteros, pp. 327-468, in Gay, C., *Historia física y política de Chile. Zoología* 7: 411 pp. Paris.
- Brauer, F. M., 1885. Systematisch-zoologische Studien. *Sitzber. k. Akad. Wiss. Wien, math.-naturwiss. Cl. (Abt. 1)* 21: 237-413, 1 pl.
- Brauer, F. M. & J. Bischof, 1903. Von Ihrer Königl. Hoheit der Prinzessin Therese von Bayern auf seiner Reise in Südamerika gesammelte Insekten. (Fortsetzung und Schluss). V. Dipteren. *Berlin. ent. Ztschr.* 47 (1902): 243-245, pl. 4.

- Brèthes, J., 1904. Insectos de Tucumán. *An. Mus. nac. Hist. nat. Buenos Aires* 4: 17-14, 329-347.
- Brèthes, J., 1908. Catálogo de los dípteros de las repúblicas del Plata. *An. Mus. nac. Hist. nat. Buenos Aires* 16: 277-305.
- Brèthes, J., 1925. Sur quelques diptères chiliens. *Revta chilena Hist. nat.* 28: 104-111.
- Bristowe, W. S., 1925. Notes on the habits of insects and spiders in Brazil. *Trans. ent. Soc. London* 72: 475-504. [Asilidae, pp. 483-484].
- Bromley, S. W., 1925. The *Bremus* resembling mallophorae of the southeastern United States (Diptera Asilidae). *Psyche* 32: 190-194.
- Bromley, S. W., 1928. New neotropical *Erax* in the American Museum of Natural History. *American Mus. Novitates* 334: 1-5.
- Bromley, S. W., 1929a. The Asilidae of Cuba (Diptera). *Ann. ent. Soc. America* 22 (2): 272-295, 14 figs., 1 pl.
- Bromley, S. W., 1929b. New Asilidae from Mexico (Diptera). *Psyche* 36 (1): 45-47.
- Bromley, S. W., 1931a. New neotropical *Andrenosoma* (Diptera: Asilidae). *Trans. American ent. Soc.* 57: 129-134.
- Bromley, S. W., 1931b. New Asilidae with revised key to the genus *Stenopogon* Loew (Diptera). *Ann. ent. Soc. America* 24 (2): 427-435.
- Bromley, S. W., 1931c. A preliminary annotated list of the robber flies of Ohio (Diptera, Asilidae). *Ohio State Mus. Sci. Bull.* 1 (2): 1-18, pls. I-IV.
- Bromley, S. W., 1932. Asilidae, pp. 2-1-282, in British Museum (Natural History), *Diptera of Patagonia and South Chile* 5 (3): 199-293. London.
- Bromley, S. W., 1933. Corrections in nomenclature (Diptera: Asilidae). *Ent. News* 44: 15.
- Bromley, S. W., 1934a. The robber-flies of Texas (Diptera: Asilidae). *Ann. ent. Soc. America* 27: 74-113, 2 pls.
- Bromley, S. W., 1934b. Asilidae, pp. 327-360, figs. 20-24, in Curran, C. H., The Diptera of Kartabo, Bartica District, British Guiana, with descriptions of new species from other British Guiana localities. *Bull. American Mus. nat. Hist.* 66: 287-523, 55 figs., 1 map.
- Bromley, S. W., 1934c. Addition to the Ohio list of robber flies. II. (Diptera: Asilidae). *Ohio J. Sci.* 34 (3): 163-164.
- Bromley, S. W., 1934d. Two new dasypogonine robber flies from the Southwest (Asilidae: Diptera). *J. New York ent. Soc.* 42: 225-226.
- Bromley, S. W., 1935a. The Laphriinae robber flies of North America (Diptera: Asilidae). *Ohio State Univ. Abstr. Doctor's Diss.* (1933-1934) 14: 125-134.
- Bromley, S. W., 1935b. Two new South American Asilidae (Diptera). *Arb. Morphol. taxon. Ent.*, Berlin-Dahlem 2 (2): 109-111, 2 figs.
- Bromley, S. W., 1935c. Notes on Texas robber flies with the description of a new species of *Proctacanthella* (Asilidae: Diptera). *Occas. Papers Mus. Zool. Univ. Michigan* 304: 1-7.
- Bromley, S. W., 1936. The genus *Diogmites* in the United States of America with descriptions of new species (Diptera: Asilidae). *J. New York ent. Soc.* 44: 225-237.
- Bromley, S. W., 1937a. The genus *Stenopogon* Loew in the United Atates of America (Asilidae: Diptera). *J. New York ent. Soc.* 45: 291-309, 4 figs.
- Bromley, S. W., 1937b. New and little known Utah Diptera with notes on the taxonomy of Diptera. *Proc. Utah Acad. Sci. Arts Letters* 14: 99-109, 4 figs.
- Bromley, S. W., 1940. New U. S. A. robber flies (Diptera: Asilidae). *Bull. Brooklyn ent. Soc.* 35 (1): 13-21.
- Bromley, S. W., 1946a. The robber flies of Brazil (Asilidae, Diptera). [Sociedade Brasileira de Entomologia] *Livro de homenagem a R. F. d'Almeida*: 103-117, pl. 5. São Paulo.
- Bromley, S. W., 1947. Ohio robber flies. IV. (Diptera, Asilidae). *Ohio J. Sci.* 47 (2): 67-68.
- Bromley, S. W., 1950. Florida Asilidae (Diptera) with description of one new species. *Ann. ent. Soc. America* 43 (2): 227-239, 1 pl.
- Bromley, S. W., 1951. Asilid notes (Diptera), with descriptions of thirty-two new species. *American Mus. Novitates* 1532: 1-36, 8 figs.
- Brower, L. P., J. v. Z. Brower & P. W. Westcott, 1960. Experimental studies of mimicry. 5. The reaction of toads (*Bufo terrestris*) to bumblebees (*Bombus americanorum*) and their robber fly mimics (*Mallophora bombooides*), with a discussion of aggressive mimicry. *American Naturalist* 94: 343-355.
- Brown, C. J. D., 1929. A morphological and systematical study of Utah Asilidae. *Trans. American ent. Soc.* 54: 295-320, pl. 28-30.
- Brown, F. M., 1941. A gazetteer of entomological station in Ecuador. *Ann. ent. Soc. America* 34: 809-851.
- Bueno, V. H. P., 1986. Aspectos etológicos de *Porasilius barbiellinii* Curran, 1934 (Diptera, Asilidae) em pastagem de *Brachiaria decumbens* Stapf. *Revta brasileira Ent.* 30 (1): 17-26.
- Bueno, V. H. P., 1987. Aspectos biológicos e ritmo diário das atividades de *Porasilius barbiellinii*, predador da cigarrinha-das-pastagens *Deois flavopicta* (Stål, 1854) (Homoptera: Cercopidae). *Pesq. agropec. brasileira*, Brasília, D. F. 22 (9/10): 903-915.
- Bueno, V. H. P. & E. Berti Filho, 1987. Ocorrência de asilídeos em pastagens de *Brachiaria decumbens* Stapf, no município de Três Corações, MG, e flutuação populacional de *Porasilius barbiellinii* Curran, 1934 (Diptera, Asilidae). *Revta Agric.*, Piracicaba 63 (2): 141-158.
- Burmeister, H. C. C., 1829. *De insectorum systemate naturali*, 40 pp. Grunert, Halle.
- Burmeister, H. C. C., 1861. *Reise durch die La Plata Staaten, mit besonders Rücksicht, auf die physische Beschaffenheit und der Culturzustand der argentinischen Republik* 1: vi + 503 pp.; 2: iv + 538 pp. Halle.
- Byers, G. W., 1962. Catalogue of the types in the Snow Entomological Museum. Part III (Diptera). *Univ. Kansas Sci. Bull.* 43 (5): 131-181.
- Carrera, M., 1943. Nova espécie de *Aphamartania* Schiner, 1866, de Curitiba (Dipt., Asilidae). *Argos Mus. paranaense* 3: 119-122, pl. 10.
- Carrera, M., 1944. Chave sinóptica da subfamília Leptogastrinae (Diptera, Asilidae), com a descrição de um novo gênero e uma nova espécie. *Papéis avulsos Zool.*, São Paulo 4: 85-94, 3 figs.
- Carrera, M., 1945a. Relação de alguns Asilidae (Diptera) com suas presas. *Papéis avulsos Zool.*, São Paulo 5 (17): 159-166, 1 pl.
- Carrera, M., 1945b. Pequenas notas sobre Asilidae (Diptera). I. Sobre algumas espécies dos gêneros *Townsendia*,

- Hypenates e Aphantartania*. II. Descrição de uma nova espécie de *Hybozelodes* (Laphriinae). *Papéis avulsos Zool.*, São Paulo 5 (18): 167-174.
- Carrera, M., 1945c. Estudo sobre os gêneros *Glaphyropyga* e *Senoprosopis* com descrição de novo gênero e novas espécies. *Papéis avulsos Zool.*, São Paulo 5 (19): 175-192, 38 figs.
- Carrera, M., 1946a. Pequenas notas sobre Asilidae (Diptera). III. Redescrição de *Dissmeryngodes anticus* (Wied.) e descrição do alótipo de *Othonomyia triangularis* Hermann (Laphriinae, Atomosiini). *Papéis avulsos Zool.*, São Paulo 7 (7): 93-100.
- Carrera, M., 1946b. Sobre algumas espécies do gênero *Atoniomyia* Hermann, 1912 (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 6 (9): 113-128, 20 figs.
- Carrera, M., 1946c. Nova espécie de *Senobasis* Macq., 1838 (Diptera, Asilidae). [Sociedade Brasileira de Entomologia] *Livro de homenagem a R. F. d'Almeida*: 121-124. São Paulo.
- Carrera, M., 1946d. Sobre o gênero *Cerozodus* Bigot, 1857 (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 7 (22): 247-256.
- Carrera, M., 1947a. Asilídeos coligidos no Paraguai pela Missão Científica Brasileira (Diptera). *Papéis avulsos Zool.*, São Paulo 8 (3): 39-48.
- Carrera, M., 1947b. Sobre o gênero *Leptopteromyia* Williston, 1908 (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 8 (): 89-96, 7 figs.
- Carrera, M., 1947c. Novo gênero e nova espécie de Asilidae (Diptera) do nordeste brasileiro. *Papéis avulsos Zool.*, São Paulo 8 (7): 203-208, 9 figs.
- Carrera, M., 1947d. Segunda relação de alguns Asilidae (Diptera) e suas presas, com descrição prévia de duas novas espécies. *Papéis avulsos Zool.*, São Paulo 8 (23): 265-271.
- Carrera, M., 1948a. Sobre as espécies de *Diogmites* da fauna amazônica (Diptera, Asilidae). *Bolm Mus. paraense Hist. nat. E. Goeldi* 10: 115-122, 3 figs.
- Carrera, M., 1948b. Sobre o gênero *Lycomyia* Bigot, 1857 (Diptera, Asilidae). *Revta Ent.*, Rio de Janeiro 19: 423-428, 2 figs.
- Carrera, M., 1949. Contribuição ao conhecimento dos Asilidae neotropicais (Diptera). I. Sobre as espécies brasileiras com esporão na tibia. *Arqos Zool.*, São Paulo 7: 1-148, 10 pls., 210 figs.
- Carrera, M., 1950a. Uma nova espécie de *Glaphyropyga* da Zona do Canal no Panamá (Diptera, Asilidae). *Dusenia*, Curitiba 1: 27-32, 1 fig.
- Carrera, M., 1950b. Asilídeos da Argentina (Diptera). I. Sobre o gênero *Prolepsis* Walker, 1851. *Dusenia*, Curitiba 1: 83-95, 4 figs.
- Carrera, M., 1950c. Synoptical keys for the genera of Brazilian "Asilidae" (Diptera). *Revta brasileira Biol.*, Rio de Janeiro 10(1): 99-111.
- Carrera, M., 1951. Sobre alguns gêneros da tribo Laphriini (Diptera, Asilidae). *Arqos Mus. nac.*, Rio de Janeiro 41: 113-126, 2 pls., 14 figs.
- Carrera, M., 1952a. Pequenas notas sobre Asilidae (Diptera). IV. Descrição de duas novas espécies de *Atomosia* e *Rhopalogaster*. *Papéis avulsos Zool.*, São Paulo 10 (10): 209-212.
- Carrera, M., 1952b. Sobre o gênero *Threnia* Schiner, 1866 (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 10 (12): 235-253, 15 figs.
- Carrera, M., 1952c. Sobre a tribo Megapodini (Diptera, Asilidae, Dasypogoninae). *Arqos Zool.*, São Paulo 8 (2): 53-88, 29 figs.
- Carrera, M., 1953a. Pequenas notas sobre Asilidae (Diptera). V. Sobre alguns Dasypogoninae das coleções do Museu Britânico e do Instituto Miguel Lillo. *Papéis avulsos Zool.*, São Paulo 8 (16): 271-277.
- Carrera, M., 1953b. As espécies neotropicais do gênero *Diogmites* (Asilidae). *Arqos Zool.*, São Paulo 8: 169-208, 26 figs.
- Carrera, M., 1955a. Novos gêneros e novas espécies de Dasypogoninae neotropicais (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 12 (2): 99-118, 10 figs.
- Carrera, M., 1955b. Sobre o gênero *Dicranus* Loew, 1851 (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 12 (10): 235-246.
- Carrera, M., 1955c. Asilídeos da Argentina (Diptera). II. *Aczelia*, novo gênero para *Laparus argentinus* Wulp, 1882. *Papéis avulsos Zool.*, São Paulo 12 (14): 297-302, 3 figs.
- Carrera, M., 1958a. Dípteros de Boracéia. I. Asilidae. *Papéis avulsos Zool.*, São Paulo 13 (12): 141-154, 9 figs.
- Carrera, M., 1958b. Sobre o gênero *Dasyllis* Loew, 1851 (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 13 (16): 213-220, 4 figs.
- Carrera, M., 1959. Sobre alguns asilídeos neotropicais (Diptera) do "Zoologische Sammlung des Bayerischen Staates". *Opuscula entomologica* 30: 1-13, 6 figs.
- Carrera, M., 1960a. Asilidae (Diptera) da coleção Seabra. *Arqos Zool.*, São Paulo 11 (17): 147-170, 10 figs.
- Carrera, M., 1960b. Asilidae (Diptera) do Uruguai e regiões circunvizinhas. *Revta Soc. uruguaya Ent.* 4: 45-53.
- Carrera, M. & M. A. V. d'Andretta, 1950. Asilídeos do México (Diptera). *Papéis avulsos Zool.*, São Paulo 9 (12): 159-191, 61 figs.
- Carrera, M. & M. A. V. d'Andretta, 1952. Relação de alguns Asilidae e suas presas (III). *Papéis avulsos Zool.*, São Paulo 10 (13): 253-259.
- Carrera, M. & M. A. V. d'Andretta, 1953. Asilídeos do Peru (Diptera). *Papéis avulsos Zool.*, São Paulo 11 (9): 63-78, 11 figs.
- Carrera, M. & C. E. Machado-Allison, 1963. Contribución al conocimiento de los Asilidae (Diptera) de Venezuela. *Acta biol. venezolana* 3 (15): 233-267, 5 figs.
- Carrera, M. & C. E. Machado-Allison, (1968). Sobre el genero *Lastaurus* Loew, 1851 (Dipt., Asilidae). *Boln Soc. venezolana Cienc. nat.* 110 (tomo 26): 484-503, "1966".
- Carrera, M. & N. Papavero, 1962. Saropogonini neotropicais (Diptera, Asilidae, Dasypogoninae). *Studia ent.*, Rio de Janeiro 5: 39-64, 6 figs.
- Carrera, M. & N. Papavero, 1965. Sobre os gêneros *Pritchardia* Stuardo, *Alyssomyia* Hull e *Hypenates* Loew (Diptera, Asilidae). *Papéis avulsos Zool.*, S. Paulo 18(6): 47-55, 2 figs.
- Carrera, M. & M. A. Vulcano, 1961. Relação de alguns Asilidae (Diptera) e suas presas. (IV). *Revta brasileira Ent.*, São Paulo 10: 67-80.
- Castelo, M. K., 2001a. Investigaciones básicas para resolver

- um viejo problema: el moscardón cazador de abejas. *Campo & Abejas* 5 (22): 4-5.
- Castelo, M. K., 2001b. El moscardón cazador de abejas perjudicó la cosecha de miel. *Revista de Apicultura Los Lirios* 5 (55): 4-7.
- Castelo, M. K., 2002a. Moscardón cazador de abejas, *Mallophora ruficauda* (Diptera: Asilidae). Algunas consideraciones sobre su presencia em los apiarios. *Ciencia apícola* 1 ((1): 10-18.
- Castelo, M. K., 2002b. Moscardón cazador de abejas. El mayor depredador de Argentina (1-2). *El Colmenar* 16 (67): 53-55, (68): 32-36.
- Castelo, M. K., 2004a. Host-seeking behavior in larvae of the robber fly *Mallophora ruficauda* (Diptera: Asilidae). *J. Insect Physiol.* 50 (4): 331-336.
- Castelo, M. K. & A. F. Capurro, 2000. Especificidad y densidad dependencia inversa em parasitoides con oviposición fuera del hospedador: El caso de *Mallophora ruficauda* (Diptera: Asilidae) en la pampa argentina. [IPEF] *Ecología austral*, Buenos Aires 13 (33): 31-40.
- Castelo, M. K. & A. F. Capurro, 2001. Moscardón cazador de abejas: Un importante enemigo de la actividad apícola em Argentina. *Gestión apícola* 5 (23): 13-18.
- Castelo, M. K. & J. C. Corley, 2004a. Evaluación de la capacidad reguladora del moscardón cazador de abejas *Mallophora ruficauda* (Diptera: Asilidae) sobre los gusanos del suelo (Coleoptera: Scarabaeidae). [Instituto Nacional de Tecnología Agropecuaria] *Revta Invest. Agropecuarias*, Buenos Aires 33 (1): 61-80.
- Castelo, M. K. & J. C. Corley, 2004b. Oviposition behaviour in the robber fly *Mallophora ruficauda* (Diptera: Asilidae). *Ann. ent. Soc. America* 97 (5): 1050-1054.
- Castelo, M. K. & C. R. Lazzari, 2004. Host-seeking behavior in larvae of the robber-fly *Mallophora ruficauda* (Diptera: Asilidae). *J. Insect Physiol.* 50 (4): 331-336.
- Castelo, M. K., M. Ney-Nifle, J. Corley & C. Bernstein, 2006. Oviposition height increases parasitism success by the robber fly *Mallophora ruficauda* (Diptera: Asilidae). *Behav. Ecol. Sociobiol.* 61 (2): 231-243.
- Castillo, E. E., R. V. Jerez & J. N. Artigas C., 1994. Microescultura coriónica en huevos de Asilidae (Diptera – Asilinae, Dasypogoninae, Laphriinae y Stenopogoninae). *Boln Soc. Biol.. Concepción* 65 (1): 107-116.
- Cekalovic K., T. & J. N. Artigas, 1969. Catálogo de los tipos de Insecta depositados en la colección del Departamento de Zoología de la Universidad de Concepción (INCO). Parte I. Dic. 1968. *Boln Soc. Biol. Concepción* 41: 111-133.
- Cekalovic K., T. & J. N. Artigas C., 1974. Catálogo de los tipos de Insecta depositados en la colección del Departamento de Zoología de la Universidad de Concepción (INCO) (Enero, 1969 a Enero, 1974). II. *Boln Soc. Biol. Concepción* 47: 233-246. [Asilidae, pp. 242-243].
- Cekalovic K., T. & J. N. Artigas C., 1981. Catálogo de los tipos depositados en la colección del Departamento de Zoología de la Universidad de Concepción, Chile. *Boln Soc. Biol.. Concepción* 51: 75-107.
- Clements, A. N. & F. D. Bennett, 1969. The structure and biology of a new species of *Mallophora* Macq. (Diptera, Asilidae) from Trinidad, W. I. *Bull. ent. Res* 58 (3): 455-463, 11 figs.
- Cockerell, T. D. A., 1894. On the habits of some Asilidae. *Ent. News* 5: 173-174.
- Cockerell, T. D. A., 1908. Fossil insects from Florissant, Colorado. *Bull. American Mus. nat. Hist.* 24: 59-69.
- Cockerell, T. D. A., 1909. New fossil insects from Florissant, Colorado. *Ann. ent. Soc. America* 2: 251-256.
- Cockerell, T. D. A., 1910. Fossil insects and a crustacean from Florissant, Colorado. *Bull. American Mus. nat. Hist.* 28: 275-288.
- Cockerell, T. D. A., 1911. Fossil insects from Florissant, Colorado. *Bull. American Mus. nat. Hist.* 30: 71-82.
- Cockerell, T. D. A., 1913. A fossil asilid fly from Colorado. *Entomologist* 46: 213-214.
- Cockerell, T. D. A., 1920. Eocene insects from the Rocky Mountains. *Proc. U. S. natl Mus.* 57 (2313): 233-260, pls. 32-36.
- Cole, F. R., 1916. New species of Asilidae from southern California. *Psyche* 23: 63-69, 3 pls.
- Cole, F. R., 1923. Diptera from the islands and adjacent shores of the Gulf of California. II. *Proc. California Acad. Sci.* (4) 12: 457-481, 16 figs.
- Cole, F. R., 1924. Notes on the dipterous family Asilidae, with descriptions of new species. *Pan-Pacific Entomologist* 1: 7-13, 6 figs.
- Cole, F. R., 1969. *Carreraomyia*, new name for *Carreraia*. *J. Kansas ent. Soc.* 41 (4) (1968): 543.
- Cole, F. R. & A. E. Pritchard, 1964. The genus *Mallophora* and related asilid genera in North America (Diptera: Asilidae). *Univ. California Publs Ent.* 36: 43-100, 4 pls., 11 figs.
- Cole, F. R. & J. Wilcox, 1938. The genera *Lasiopogon* Loew and *Alexiopogon* Curran in North America (Diptera – Asilidae). *Entomologica americana* (n. s.) 18 (1-2): 1-91, pls. 1-3.
- Copello, A., 1922. Biología de “*Mallophora ruficauda*” Wied. *Physis*, Buenos Aires 6: 30-42, 2 pls.
- Copello, A., 1927. Biología del moscardón cazador de abejas (*Mallophora ruficauda* Wied.). *Minist. Agric., Sección Propaganda e Informes*, Buenos Aires 699: 1-19, 10 figs.
- Copello, A., 1942. Moscardón cazador de abejas “*Mallophora ruficauda*”. *Revta Apicultura*, Buenos Aires 19 (218): 4-5, (219): 10-11, (220): 13-14, (221): 14, (222): 12-13.
- Coquillett, D. W., 1893a. Synopsis of the asilid genus *Anisopogon*. *Canadian Entomologist* 25: 20-22.
- Coquillett, D. W., 1893b. Synopsis of the asilid genus *Blacodes*. *Canadian Entomologist* 25: 33-35.
- Coquillett, D. W., 1893c. Synopsis of the asilid genera *Mallophora* and *Nicocles*. *Canadian Entomologist* 25: 118-120.
- Coquillett, D. W., 1893d. A new asilid genus related to *Erax*. *Canadian Entomologist* 25: 175-177.
- Coquillett, D. W., 1898. Synopsis of the asilid genus *Ospriocerus*. *Ent. News* 9: 37.
- Coquillett, D. W., 1902. New orthorrhaphous Diptera from Mexico and Texas. *J. New York ent. Soc.* 10: 136-141.
- Coquillett, D. W., 1904a. New North American Diptera. *Proc. ent. Soc. Washington* 6: 90-98.
- Coquillett, D. W., 1904b. New North American Diptera. *Proc. ent. Soc. Washington* 6: 166-192.

- Coquillett, D. W., 1904c. Diptera from southern Texas with descriptions of new species. *J. New York ent. Soc.* 12: 31-35.
- Coquillett, D. W., 1910. The type-species of North American genera of Diptera. *Proc. U. S. natl Mus.* 37 (no. 1719): 499-647.
- Costa, A., 1863. Nuovi studii sulla entomologia della Calabria Ulteriore. *Atti Accad. Sci. fis. matem. Napoli* 1 (2): 1-80, 4 pls.
- Costa, A., 1886. Acquiste fatti durante l'anno 1863. *Annuario Mus. Zool. Univ. Napoli* 3 (1863): 13-41.
- Costa, A., 1883. Notizie ed osservazioni sulla geo-fauna sarda. *Atti Accad. Sci. fis. mat. Napoli* (2) 1 (2): 1-109.
- Coronado Blanco, J. M. & E. Ruiz Cancino, 1999. Primer registro de *Atomosia macquarti* Bellardi (Diptera: Asilidae) como depredador de *Unaspis citri* (Comstock) (Homoptera, Diaspididae). *Folia ent. mexicana* 105: 81-82.
- Couri, M. S., C. J. E. Lamas M., C. C. de C. Aires, C. A. de Mello-Patiu, V. C. Maia, D. M. Pamplona & P. Magno, 2000. Diptera da Serra do Navio (Amapá, Brasil): Asilidae, Bombyliidae, Calliphoridae, Micropezidae, Muscidae, Sarcophagidae, Stratiomyidae [sic], Syrphidae, Tabanidae e Tachinidae. *Revta brasileira Zoociências*, Juiz de Fora 2 (1): 91-101.
- Cresson, E. T., Jr., 1926. Concerning the types of *Mallophora rex* and *cheysomela* [sic] Bromley. *Psyche* 33: 91.
- Crouzel, I. S. de, 1965. Parasitismo em "gusano branco" en la República Argentina. *Revta Soc. ent. argentina* 27: 83-87. [Asilidae, pp. 85-87].
- Curran, C. H., 1925a. Descriptions of four new neotropical Diptera. *Trans. American ent. Soc.* 51 (3): 259-261.
- Curran, C. H., 1925b. *Buckellia*, a new genus of Asilidae (Diptera). *Canadian Entomologist* 57: 156.
- Curran, C. H., 1926. New Diptera from the West Indies. *American Mus. Novitates* 220: 1-14.
- Curran, C. H., 1928a. New species of *Ommatius* from America, with key (Asilidae, Diptera). *American Mus. Novitates* 327: 1-6.
- Curran, C. H., 1928b. Insects of Porto Rico and the Virgin Islands. Diptera or two-winged flies. Pt. 1, pp. 1-118, 39 figs., in New York Academy of Sciences, *Scientific survey of Puerto Rico and the Virgin Islands*. Vol. 11. New York.
- Curran, C. H., 1930a. New Diptera from North and Central America. *American Mus. Novitates* 415: 1-16, 1 fig.
- Curran, C. H., 1930b. New American Asilidae (Diptera). *American Mus. Novitates* 425: 1-21, 3 figs.
- Curran, C. H., 1931a. First supplement to the "Diptera of Porto Rico and the Virgin Islands". *American Mus. Novitates* 456: 1-23, 4 figs.
- Curran, C. H., 1931b. New American Asilidae (Diptera). II. *American Mus. Novitates* 487: 1-25.
- Curran, C. H., 1934a. New American Asilidae (Diptera). III. *American Mus. Novitates* 752: 1-18, 1 fig.
- Curran, C. H., 1934b. *The families and genera of North American Diptera*, 512 pp., figs. New York.
- Curran, C. H., 1934c. The Diptera of Kartabo, Bartica District, British Guiana, with descriptions of new species from other British Guiana localities. *Bull. American Mus. nat. Hist.* 66: 287-523, 55 figs., 1 map.
- Curran, C. H., 1935. New American Asilidae (Diptera). IV. *American Mus. Novitates* 806: 1-12.
- Curran, C. H., 1936. Three new neotropical Diptera. *Revta chilena Hist. nat.* 40: 331-335.
- Curran, C. H., 1941. Some new species of *Mallophora* Macquart (Asilidae, Diptera). *J. New York ent. Soc.* 49: 269-284.
- Curran, C. H., 1942. American Diptera. *Bull. American Mus. nat. Hist.* 80: 51-84.
- Curran, C. H., 1951. The West Indian species of *Mydas* and *Proctacanthus* (Diptera: Mydidae and Asilidae). *American Mus. Novitates* 1507: 1-9.
- Curran, C. H., 1953. The Asilidae and Mydidae of the Bimini Islands, Bahamas, British West Indies (Diptera). *American Mus. Novitates* 1644: 1-6.
- Dennis, D. S., J. K. Barnes & L. Knutson, 2008. The pupal case of *Mallophora atra* Macquart (Diptera: Asilidae) from South America. *Proc. ent. Soc. Washington* 110 (1): 234-241.
- Dennis, D. S. & J. A. Gowen, 1978. A "nocturnal" foraging record for *Diogmites neoternatus* (Diptera: Asilidae). *Proc. ent. Soc. Washington* 80: 313-314.
- Dennis, D. S. & L. Knutson, 1988. Descriptions of pupae of South American robber flies (Diptera: Asilidae). *Ann. ent. Soc. America* 81: 851-864.
- Dennis, D. S. & R. J. Lavigne, 1976. Descriptions and notes on the pupae and pupal cases of ten species of Wyoming robber flies (Diptera: Asilidae). *Proc. ent. Soc. Washington* 78: 277-303, 43 figs.
- Dennis, D. S., R. J. Lavigne & S. W. Bullington, 1986. Ethology of *Efferia cressoni* with a review of the comparative ethology of the genus (Diptera: Asilidae). *Proc. ent. Soc. Washington* 88: 42-55.
- De Santis, L., 1989. El moscardón cazador de abejas. [Cooperativa Agrícola Ltda. del Oeste Mones Cazón] *Revta Industria apícola*, Buenos Aires 1 (1): 16-24.
- De Santis, L. & L. G. Cornejo, 1990. *El moscardón cazador de abejas "Mallophora ruficauda"*, 17 pp. Conferencia de la Cooperativa Apícola del Oeste Ltda. de Mones Cazón.
- Dikow, T., 2004. Family-group names in the Asilidae (Diptera). *Studia dipterologica* 10 (2) (2003): 459-471.
- Dikow, T., 2009. Phylogeny of Asilidae inferred from morphological characters of imagines (Insecta, Diptera, Brachycera, Asiloidea). *Bull. Amer. Mus. nat. Hist.* 319: 1-175.
- Dozier, H. L., 1920. An ecological study of hammock and pine woods insects in Florida. *Ann. ent. Soc. America* 13: 325-380.
- Enderlein, G., 1914a. Dipterologische Studien. XI. Zur Kenntnis tropischer Asiliden. *Zool. Anz.* 44 (6): 241-263, 8 figs.
- Enderlein, G., 1914b. Dipterologischen Studien. XII. Zur Kenntnis der Asilidensubfamilien Dasypogoninae und Archilaphriinae. *Wien. ent. Ztg* 33: 151-174, 8 figs.
- Enderlein, G., 1936. 22. Ordnung: Zweiflügler, Diptera. Abt. 16, pp. 1-259 [=Lfg. 2, part], in Brohmer, P., P. Ehrmann & G. Ulner, eds., *Die Tierwelt Mitteleuropas* 6 (Insekten. III. Teil), Leipzig.
- Engel, E. O., 1885. Eine neue *Emphysomera* (Diptera, Fam. Asilidae). *Ent. Nachrichten* 11: 146-147.
- Engel, E. O., 1929. New or little known Asilidae from South Africa. *Ann. Transvaal Mus.* 13: 154-171.

- Engel, E. O., 1930. Die Ausbeute der deuschen Chaco-Expedition 1925/1926. Asilidae (Diptera). *Konowia* 8 (1929): 457-474, 8 figs.
- Engel, E. O., 1925-1930. Asilidae. (Fam.) 25, in Lindner, E., ed., *Die Fliegen der palaearktischen Region* 4 (2): 1-491, figs.
- Fabricius, J. C., 1775. *Systema entomologiae, sistens insectorum classes, ordines, genera, species adiectis synonymis, locis, descriptionibus, observationibus*, 832 pp. Flensburgi et Lipsiae [= Flensburg and Leipzig].
- Fabricius, J. C., 1777. *Genera insectorum eorumque characteres naturales secundum numerum, figuram, situm et proportionem omnium partium oris adiecta mantissa specierum nuper detectarum*, 310 pp. Chilonii [= Kiel].
- Fabricius, J. C., 1781. *Species insectorum exhibentes eorum differentias specificas, synonyma, auctorum, loca natalia, metamorphosin* 2: 517 pp. Hamburgi et Kilonii [= Hamburg and Kiel].
- Fabricius, J. C., 1787. *Mantissa insectorum sistens species nuper detectarum* 2: 382 pp. Hafniae [= Copenhagen].
- Fabricius, J. C., 1794. *Entomologia systematica emendata et aucta* 4: 472 pp. Hafniae [= Copenhagen].
- Fabricius, J. C., 1805. *Systema Antliatorum secundum ordines, genera, species*, xiv + 372 + 30 pp. Brunsvigae [= Brunswick].
- Farr, T. H., 1962. A new species of *Schildia* from Jamaica (Diptera: Asilidae). *Florida Entomologist* 45: 191-194, 8 figs.
- Farr, T. H., 1963. The robber-flies of Jamaica. The subfamily Leptogastrinae. *Bull. Inst. Jamaica, Sci. Ser.* 13 (1): 5-22.
- Farr, T. H., 1965. The robber-flies of Jamaica (Diptera, Asilidae). Pt. 2. *Bull. Inst. Jamaica, Sci. Ser.* 13: 5-36.
- Farr, T. H., 1973. A new species of *Eurhabdus* from Jamaica (Diptera: Leptogastridae). *Florida Entomologist* 56 (4): 324-326, 4 figs.
- Fattig, F. W., 1933. The food of the robber fly *Mallophora orcina* (Wied.) (Diptera). *Canadian Entomologist* 65: 119-120.
- Fattig, F. W., 1945. The Asilidae or robber flies of Georgia. *Emory Univ. Mus. Bull.* 3: 3-33.
- Fernández Velázquez, A. M. & P. Rodríguez Velázquez, 1996. Registros nuevos del género *Ommatius* (Diptera: Asilidae) para Cuba. *Cocuyo* 5: 10-11.
- Fisher, E. M., 1977. A review of the North American genera of Laphystiini, with a revision of the genus *Zabrops* Hull (Insecta, Diptera: Asilidae). *Proc. California Acad. Sci.* (4) 41 (5): 183-213, 74 figs., 1 table.
- Fisher, E. M., 1983. *Pilicaformidolosa* (Mosca asesina, robber fly), pp. 755-758, in Janzen, D. H., ed., *Costa Rican natural history*. University of Chicago Press, Chicago.
- Fisher, E. M., 1985. A preliminary list of the robber flies (Diptera: Asilidae) of the Tambopata Reserved Zone, Madre de Dios, Peru. *Revta peruana Ent.* 27: 25-36.
- Fisher, E. M., 1991. *Pilicaformidolosa* (Mosca asesina, robber fly), pp. 769-772, in Janzen, D., M. Chavarria A. & L. Diego Gómez P., eds., *Historia natural de Costa Rica*. University of Chicago Press, Chicago & London.
- Fisher, E. M. & H. A. Hespenheide, 1982. Taxonomy and ethology of a new Central American species of the robber fly genus *Glaphyropyga* (Diptera: Asilidae). *Proc. ent. Soc. Washington* 84 (4): 716-725, figs.
- Fisher, E. M. & H. A. Hespenheide, 1992. Taxonomy and biology of Central American robber flies with an illustrated key to genera (Diptera, Asilidae), pp. 611-632, in Quintero Arias, D. & A. Aiello, eds., *Insects of Panama and Mesoamerica. Selected studies*. Oxford University Press, London.
- Foldi, I., 2005. Ground pearls: A generic revision of the Margarodidae *sensu stricto* (Hemiptera: Sternorrhyncha: Coccoidea). *Ann. Soc. ent. France* (n.s.) 41 (1): 81-125.
- Forbes, G. S., 1987. The status of *Efferia similis* (Williston), with descriptions of three new Nearctic *Efferia* species in the *albibarbis* group (Diptera, Asilidae). *Pan-Pacific Entomologist* 63: 292-300.
- Forbes, G. S., 1988. Three new species of *Efferia* from southern California and Arizona (Diptera: Asilidae). *Ann. ent. Soc. America* 81: 554-559.
- Geller-Grimm, F., 1997. A new species of *Ichneumolaphria* Carrera from Brazil and the description of the male of *Pseudorus dandrettae* Carrera, 1949 (Diptera: Asilidae). *Studia dipterologica* 4 (2): 457-462.
- Geller-Grimm, F., 1999. Raubfliegen-Typen des Senckenberg-Museums in Frankfurt am Main, das überwiegend von Wiedemann und Jaennicke bearbeitet wurden (Insecta, Diptera, Asilidae). *Senckenbergiana biologica* 78: 205-217.
- Geller-Grimm, F., 2004. A world catalogue of the genera of the family Asilidae (Diptera). *Studia dipterologica* 10 (2) (2003): 473-526.
- Gemignani, E. V., 1930. Las especies argentinas del género *Mallophora* (Diptera) y descripción de cuatro nuevas especies. *Revta Soc. ent. argentina* 3: 133-144, 2 figs.
- Gemignani, E. V., 1931. El alotipo de *Mallophora vegeta* E. Lynch Arribálzaga (Diptera: Asilidae). *Revta Soc. ent. argentina* 3: 265-266.
- Gemignani, E. V., 1936. Uma nueva especie del género *Tolmerolestes* E. Lynch Arribálzaga (Diptera: Asilidae). *Revta chilena Hist. nat.* 39: 42-47, 1 pl.
- Ghahari, H., R. J. Lavigne & F. Geller-Grimm, 2007. Bibliography of Asilidae (Insecta: Diptera) 1996-2006. *Far Easter Entomologist (Dal'nevostochnyi Entomolog)*, Vladivostock 176: 1-39.
- Gistel, J. von N. F. X., 1848. *Naturgeschichte des Thierreiches für höheren Schulen*, xvi + 216 + [4] pp. R. Hoffmann, Stuttgart.
- Gmelin, J. F., 1792. *Carolus Linnaeus Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, synonymis, locis. Editio decima tertia, aucta, reformata. Vol. I. Regnum animalium*, Pt. 5: 2225-3020. G. E. Beer, Lipsiae [= Leipzig].
- Goslin, R. M., 1950. Some robber flies from Campbell County, Tennessee. *J. Tennessee Acad. Sci.* 25 (4): 303-306.
- Gowdey, C. C., 1926. Catalogus insectorum jamaicensis. [Department of Agriculture of Jamaica] *Ent. Bull.* 4 (1-2): 1-114.
- Grimaldi, D. A., 1990. Diptera, pp. 164-183, figs. 1-34, in Grimaldi, D. A., ed., Insects from the Santana Formation, Lower Cretaceous, of Brazil. *Bull. American Mus. nat. Hist.* 195: 1-191.
- Guérin-Méneville, F. E., (1831). Pls. 20-21 (= livrs. 23-24, part), in Duperrey, L. I., ed., *Voyage autour du monde sur la corvette de sa majesté La Coquille. Zoologie. Atlas*.

- Insectes*, 21 pls. Paris.
- Guérin-Méneville, F. E., (1838). Première division. Crustacés, arachnides et insectes, pp. 1-216, 217-318 (= livrs. 25, 26), in Duperrey, L. I., ed., *Voyage autour du monde sur la corvette de sa majesté La Coquille. Zoologie 2* (2), Paris.
- Hansen, H. J., 1883. *Fabrica oris dipterorum. Dipterenes mund i anatomisk systematisk Henseende. Første afdeling (Tabanidae, Bombyliidae, Asilidae, Thereva, Mydas, Apiocera)*, 220 pp., 5 pls. C. A. Reitzel, Kjøbenhavn.
- Hardy, D. E., 1943. New Therevidae and Asilidae in the Snow Entomological Collection. *J. Kansas ent. Soc.* 16: 24-29, 4 figs.
- Hardy, D. E., 1947. The genus *Leptopteromyia* (Diptera – Asilidae). *J. Kansas ent. Soc.* 20: 72-75.
- Hardy, G. H., 1926. A new classification of the Australian robberflies belonging to the subfamily Dasypogoninae (Diptera, Asilidae). *Proc. Linn. Soc. New South Wales* 51: 305-312.
- Hardy, G. H., 1927a. On the phylogeny of some Diptera Brachycera. *Proc. Linn. Soc. New South Wales* 52: 380-386.
- Hardy, G. H., 1927b. Further notes on the classification of Australian robberflies (Diptera, Asilidae). *Proc. Linn. Soc. New South Wales* 52: 387-398.
- Hardy, G. H., 1928. Third contribution towards a new classification of Australian Asilidae. *Proc. Linn. Soc. New South Wales* 53: 468-473.
- Hardy, G. H., 1930. Fifth contribution towards a new classification of Australian Asilidae (Diptera). *Proc. Linn. Soc. New South Wales* 55 (3): 249-260.
- Hardy, G. H., 1948. On classifying Asilidae. *Entomologist's monthly Mag.* 84: 116-119, 7 figs.
- Harris, T. W., 1833. Insects, pp. 566-595, in Hitchcock, E., *Report on the geology, mineralogy, botany and zoology of Massachusetts*, 12 + 700 pp., 3 maps. J. S. & C. Adams, Amherst.
- Helsdingen, P. J. van, 1996. An annotated list of the Wiedemann types in the Leiden Museum. *Studia dipterologica* 3 (2): 171-193.
- Hendel, F., 1936-1937. 16. Ordnung der Pterygogenea: Diptera – Fliegen, pp. 1729-1756, 1757-1884, 1885-1980 (1936), 1981-2140 (1937), in Kükenthal's *Handbuch der Zoologie* 4 (2). Walter de Gruyter, Berlin & Leipzig.
- Hengst, M. & J. N. Artigas C., 2003. Revisión del género *Philonerax* Bromley, 1932 (Diptera: Asilidae: Asilinae) y dos nuevas especies. *Revta chilena Hist. nat.* 76: 287-296, 7 figs.
- Hermann, F., 1905. Beitrag zur Kenntnis der Asiliden (Dipt.). *Berlin. ent. Ztschr.* 50: 14-42, 29 figs.
- Hermann, F., 1908. Beitrag zur Kenntnis der Asiliden (IV). Dipt. Nachträge zu meiner Mitteilung über die um das Genus *Laphystia* sich gruppierenden Asilidenformen. *Berlin. ent. Ztschr.* 53: 152-170, pl. 4.
- Hermann, F., 1912. Beiträge zur Kenntnis der südamerikanischen Dipteren Fauna auf Chile, Peru, und Bolivia, ausgeführt von W. Schnuse. Familie Asilidae. *Nova Acta Acad. caes.-leopold.-carol.* 96: 1-275, 5 pls., 85 figs.
- Hermann, F., 1920. Beitrag zur allgemeinen Systematik der Asiliden. *Zool. Jahrb. (Abt. Syst. Geogr. Biol. Tiere)* 43: 161-194.
- Hermann, F., 1921. in Zürcher, L., Dipteren-Ausbeute aus Paraguay: Mydaiden und Asiliden. *Arch. Naturgesch.* 87 (A, 3): 118-122.
- Hermann, F., 1926a. Die Gattungen der Leptogastrinen (Dipt.). *Verh. zool.-bot. Ges. Wien* 74-75 (1924-1925): 140-152.
- Hermann, F., 1926b. Beitrag zur Kenntnis der Asiliden. Der Verwandtschaftskreis des gen. *Holcocephala* Jännicke. *Verh. zool.-bot. Ges. Wien* 74-75 (1924-1925): 153-191, 9 figs.
- Hine, J. S., 1906. A preliminary report on the horseflies of Louisiana, with a discussion of remedies and natural enemies. *Circular of the State Crop Pest Commission of Louisiana* 6: 1-37, 20 figs.
- Hine, J. S., 1907. Robber flies of the genus *Philonicus*. *Ohio Naturalist* 7 (6): 115-118.
- Hine, J. S., 1909. Robberflies of the genus *Asilus*. *Ann. ent. Soc. America* 2: 136-170, 2 pls. [= Contributions from the Ohio State University Department of Zoology and Entomology no. 32].
- Hine, J. S., 1911a. Robberflies of the genera *Promachus* and *Proctacanthus*. *Ann. ent. Soc. America* 4: 153-172. [= Contributions from the Ohio State University Department of Zoology and Entomology no. 33].
- Hine, J. S., 1911b. New species of Diptera of the genus *Erax*. *Ohio J. Sci.* 11 (6): 307-311.
- Hine, J. S., 1916. Descriptions of robber flies of the genus *Erax*. *Ohio J. Sci.* 17 (1): 21-22.
- Hine, J. S., 1917. Costa Rican Diptera collected by Philip P. Calvert, 1909, 1910. 2. Tabanidae and Asilidae. *Trans. American ent. Soc.* 43: 291-299.
- Hine, J. S., 1918. Notes on robber flies from southwest Texas, collected by the Bryant Walker Expedition, with a description of a new species of *Erax*. *Occas. Papers Michigan State Univ. Mus. Zool.* 61: 1-7.
- Hine, J. S., 1919. Robberflies of the genus *Erax*. *Ann. ent. Soc. America* 12: 103-157, 3 pls. [= Contributions from the Ohio State University Department of Zoology and Entomology no. 57].
- Hine, J. S., 1922. Some robberflies in the University of Michigan Museum of Zoology, and the description of a new species. *Occas. Papers Mus. Zool. Univ. Michigan* 121: 1-7.
- Hogue, C. L., 1974. The insects of the Los Angeles Basin. *Nat. Hist. Mus. (Los Angeles Co.) Sci. Ser.* 27: 1-173.
- Hradský, M., 1968. A new Brazilian species of *Phellopteron Hull* (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 21 (17): 165-167, 2 figs.
- Hull, F. M., 1956. Some new genera of Asilidae (Diptera). *Bull. Brooklyn ent. Soc.* 51: 69-72.
- Hull, F. M., 1957. More flies of the family Asilidae (Diptera). *Ann. Mag. nat. Hist.* 12 (10): 884-895.
- Hull, F. M., 1958a. Some robber flies (Diptera: Asilidae). *Ent. News* 69: 99-108.
- Hull, F. M., 1958b. Some species and genera of the family Asilidae (Diptera). *Proc. ent. Soc. Washington* 60 (6): 251-257.
- Hull, F. M., 1958c. New genera of robber flies (Diptera: Asilidae). *Revta brasileira Biol.*, Rio de Janeiro 18 (3): 317-324.
- Hull, F. M., 1958d. Some flies of the family Asilidae (Diptera)

- from Australia and Brazil. *Proc. ent. Soc. London* (B) 27: 159-164.
- Hull, F. M., 1958e. Some new species of the genus *Bathypogon* Loew (Diptera, Asilidae). *Bull. Brooklyn ent. Soc.* 53 (3): 62-65.
- Hull, F. M., 1958f. A new genus and two new species of Asilidae (Diptera). *Bull. Brooklyn ent. Soc.* 53: 94-99.
- Hull, F. M., 1958g. Some flies of the family Asilidae (Diptera). *Psyche* 64: 90-96.
- Hull, F. M., 1960. New species of Syrphidae and Asilidae. *Pan-Pacific Entomologist* 36: 69-74.
- Hull, F. M., 1962. Robber flies of the world. The genera of the family Asilidae. *Smithson. Inst. Bull.* 294 (1): 1-432, (2): 433-907, 1536 figs.
- Hull, F. M., 1963. A new species of *Laphystia* Loew (Diptera: Asilidae). *Ent. News* 64: 202-203.
- Hunter, W. D., 1900-1901. A catalogue of the Diptera of South America. Part II. *Trans. American ent. Soc.* 27: 121-155.
- ICZN (International Commission on Zoological Nomenclature), 1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. *Bull. zool. Nomencl.* 20: 339-342.
- Jaennicke, J. F., 1867. Neue exotische Dipteren. *Abh. senckenb. naturf. Ges.* 6: 311-405, 2 pls. [Also publ. sep., as "Neue exotische Dipteren aus den Museen zu Frankfurt a. M. und Darmstadt", 100 pp., 2 pls., Frankfurt, 1868].
- James, M. T., 1933. New Asilidae from Colorado. *American Mus. Novitates* 596: 1-3.
- James, M. T., 1937. New Colorado Asilidae (Diptera). *Ent. News* 48: 12-15.
- James, M. T., 1939. A preliminary review of certain families of Diptera from the Florissant Miocene beds. II. *J. Paleontol.* 132: 42-48.
- James, M. T., 1941. The robber flies of Colorado (Diptera, Asilidae). *J. Kansas ent. Soc.* 14: 27-53.
- James, M. T., 1953. The Diptera collected on the Cockerell and Hubbell Expedition to Honduras. Pt. 2. Asilidae. *J. Washington Acad. Sci.* 43: 46-57.
- Johnson, C. W., 1894. List of the Diptera of Jamaica with descriptions of new species. *Proc. Acad. Nat. Sci. Philadelphia* 1894 (2): 271-281.
- Johnson, C. W., 1903. A new genus and four new species of Asilidae. *Psyche* 10: 111-114.
- Johnson, C. W., 1919. A revised list of the Diptera of Jamaica. *Bull. American Mus. nat. Hist.* 41 (8): 421-449.
- Kaletta, F., 1974. Las especies del género *Eumecosoma* Schiner (Diptera, Asilidae) conocidas en Venezuela, con descripción de dos especies nuevas. *Revta Fac. Agron. Maracay* 8 (2): 99-106, 1 fig.
- Kaletta, F., 1976. Algunas especies nuevas de la tribu Stichopogonini de Venezuela (Diptera, Asilidae). *Revta Fac. Agron. Maracay* 9 (1): 65-74.
- Kaletta, F., 1978. Nuevos Atomosiini (Diptera, Asilidae) de Venezuela. *Revta Fac. Agron. Maracay* 26: 47-63.
- Kaletta, F., 1983. Zwei neue *Cophura*-Arten aus Venezuela (Diptera: Asilidae). *Ent. Ztschr.*, Frankfurt a. Main 93 (20): 300-303.
- Kaletta, F., 1986. Nuevos Atomosiini de Venezuela (Diptera Asilidae). *Boln Soc. venezolana Cienc. nat.* 144 (Tomo 41): 257-265.
- Kallweit, U., 1992. Das Typenmaterial der Zweiflügler des Staatlichen Museum für Tierkunde Dresden (Insecta). Teil 1: Diptera-Asilidae. *Ent. Abhandl. Ber. Staatl. Mus. Tierkunde Dresden* 54 (11): 193-198.
- Karl, O., 1935. Die Dipterenfauna Pommerns. Diptera Brachycera. *Stettin. ent. Ztg* 96: 106-130.
- Kertész, K., 1909. *Catalogus dipterorum hucusque descriptorum. (IV). Oncodidae, Nemestrinidae, Mydidae, Asilidae* 4: 348 pp. Budapest.
- Kirby, W. & W. Spence, 1817. *Introduction to entomology, or elements of the natural history of insects* 2: 1-259, 2 pls. Longman, London.
- Kirby, W. F., 1882. Diptera, in *Zoological Record. Insecta* 1881: 233-254. London.
- Knutson, L. V., 1971. *Trachysphyrus nigricornis* (Brullé), prey of *Araiopgon gayi* (Macquart) (Hymenoptera: Ichneumonidae – Diptera: Asilidae). *Proc. ent. Soc. Washington* 73: 248.
- Knutson, L. V., 1972. Pupa of *Neomochtherus augustipennis* (Hine), with notes on feeding habits of robber flies and a review of publications on morphology of immature stages (Diptera: Asilidae). *Proc. biol. Soc. Washington* 85: 163-178.
- Knutson, L. V., 1976. Key to subfamilies of robberflies based on pupal cases with a description of the pupal case of *Doryclus distendens* (Asilidae: Megapodinae). *Proc. biol. Soc. Washington* 88: 509-514.
- Lamas M., C. J. E. & G. B. Mellinger, 2008. A new species of *Cnodalomyia* Hull, 1962 (Diptera, Asilidae, Asilinae) from Brazil. *Zootaxa*, Auckland 1676: 37-43, 11 figs.
- Lamas M., G., 1971a. Description of the first South American species of *Stenopogon* Loew (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 25 (2): 15-18, 5 figs.
- Lamas M., G., 1971b. The genus *Porasilius* Curran in South America (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 25 (7): 45-55, 16 figs.
- Lamas M., G., 1972. A catalogue of Peruvian Asilidae (Diptera), with keys to the identification and descriptions of two new species. *An. Ier. Congr. latino-amer. Ent. (Revta peruana Ent.)* 15 (2): 304-316, figs.
- Lamas M., G., 1973a. Taxonomy and evolution of the "Prolepsis-complex" in the Americas (Diptera, Asilidae). *Arqos Zool.*, São Paulo 24 (1): 1-71, 100 figs., 7 figs.
- Lamas M., G., 1973b. Some comments on the genus *Cteodontina* Enderlein (Diptera, Asilidae), with the description of a new species from coastal Peru. *Papéis avulsos Zool.*, São Paulo 26 (23): 275-280.
- Lamas M., G., 1973c. A revision of *Eccritosia* Schiner, 1866 (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 27 (4): 45-68, 16 figs.
- LaPierre, L. M., 2000. Prey selection and diurnal activity of *Holcocephala oculata* (F.) (Diptera: Asilidae) in Costa Rica. *Proc. ent. Soc. Washington* 102 (3): 643-651.
- Latreille, P. A., 1802. *Histoire naturelle, générale et particulière, des crustacés et des insectes. Tome troisième*, xii + 13-467 + 1 pp. Dufart, Paris.
- Latreille, P. A., 1805. *Histoire naturelle, générale et particulière, des crustacés et des insectes* 14: 1-432, pls. 104-112. Paris.
- Lavigne, R. J., 1977. Notes on the ethology of *Eccritosia*

- zamon* (Townsend) (Diptera, Asuiliidae). *Proc. ent. Soc. Washington* 79 (4): 626-630.
- Lavigne, R. J., 1978. A case of homonymy in the genus *Machimus* (Diptera: Asilidae) in Mexico. *Proc. ent. Soc. Washington* 80: 55.
- Lavigne, R. J., 1979. Notes on the ethology of *Efferia aryrogaster* (Diptera: Asilidae) in Mexico. *Proc. ent. Soc. Washington* 81: 544-551.
- Lavigne, R. J., 1999. Bibliography update 1977-1996 for the Asilidae (Insecta: Diptera). *Univ. Wyoming agric. Exptl. Sta. Sci. Monogr.* 55: 1-91.
- Lavigne, R. J., 2003. Evolution of courtship behavior among the Asilidae (Diptera), with a review of courtship and mating. *Studia dipterologica* 9 (2) (2002): 703-712.
- Lavigne, R. J. & S. Bollington, 1999. Ethology of *Heteropogon paurosomus* Pritchard (Diptera: Asilidae) in Mexico. *J. ent. Res. Soc.*, Ankara 1 (3): 9-19.
- Lavigne, R. J. & D. S. Dennis, 1979. Notes on the ethology of *Proctacanthus nearno* (Diptera, Asilidae) in Mexico. *Proc. ent. Soc. Washington* 81: 438-442.
- Lavigne, R. J. & D. S. Dennis, 1980. Ethology of *Proctacanthella leucopogon* in Mexico (Diptera: Asilidae). *Proc. ent. Soc. Washington* 82: 160-168.
- Lavigne, R. J. & D. S. Dennis, 1985. Ethology of three coexisting species of *Efferia* (Diptera: Asilidae) in Mexico. *Proc. ent. Soc. Washington* 87: 146-160.
- Lavigne, R. J., D. S. Dennis & J. A. Gowen, 1978. Asilid literature update 1956-1976 including a brief review of robber fly biology. *Univ. Wyoming agric. exptl Sta. Sci. Monogr.* 36: 1-134.
- Lavigne, R. J., C. R. Nelson & E. T. Schreiber, 1994. New prey records for *Proctacanthus* (Diptera: Asilidae) with comments on prey choice. *Ent. News* 105: 85-97.
- Leach, W. E., 1815. Entomology, pp. 57-172, in Brewster, D., ed., *The Edinburgh Encyclopedia* 9 (1) (ENG-FLU): 1-384. Baldwin, Edinburgh.
- Lehr, P. A., 1969. Ekologo-morfologicheskii analiz ktyrei (Diptera, Asilidae). II. *Ent. Obozren.* 48 (3): 532-560, 11 figs. [Transl.: Ecologo-morphological analysis of robber flies (Diptera, Asilidae). II. *Ent. Rev.* 48: 341-357].
- Lehr, P. A., 1977. [On Laphriinae (Diptera, Asilidae) of Middle Asia], pp. 45-51, in Gorodkov, K. B., ed., *Sistematiка I evolutsya dvukrylykh nasekomykh*, 128 pp. Sbornik Inst. Zool. Akad. Nauk SSSR, Leningrad.
- Lehr, P. A., 1988. Family Asilidae, pp. 197-326, in Soós, A. & L. Papp, eds., *Catalogue of Palaearctic Diptera* 5. Akadémiai Kiadó, Budapest & Elsevier, Amstterdam.
- Lindner, E., 1929. Zur Ökologie südamerikanischer Asiliden (Dipt.). *Ztschr. wiss. Insektenbiol.* 24: 167-173.
- Linnaeus, C., 1758. *Systema naturae per regna tria naturae. Ed. X1*: 824 pp. Holmiae [=Stockholm].
- Linsley, E. G., 1960. Ethology of some bee- and wasp-killing robber flies of southeastern Arizona and western New Mexico (Diptera: Asuiliidae). *Univ. California Publs Ent.* 16: 357-382, 8 pls., 9 tables.
- Linsley, E. G. & M. A. Cazier, 1963. Attraction of insects to exsudates of *Verbesina encelioides* and *Iva ambrosiaeefolia*. *Bull. South California Acad. Sci.* 62: 109-129.
- Linsley, E. G. & R. L. Usinger, 1966. Insects of the Galapagos Islands. *Proc. California Acad. Sci.* 33: 166 [*Ommatius marginellus*].
- Llano, R. J., 1959. Observaciones biológicas de insectos bonaerenses. *Supl. Revta Educación*, La Plata 12: 1-136.
- Loew, H., 1847. Über die europäischen Raubfliegen (Diptera -Asilica). *Linnaea ent.*, Stettin 2: 384-568, 587-591 [cont.].
- Loew, H., 1848. Über die europäischen Raubfliegen (Diptera -Asilica) [cont.]. *Linnaea ent.*, Stettin 3: 386-495 [cont.].
- Loew, H., 1849. Über die europäischen Raubfliegen (Diptera -Asilica) [concl.]. *Linnaea ent.*, Stettin 4: 1-155.
- Loew, H., 1851. Bemerkungen über die Familie Asiliden. *Programm Realschule zu Meseritz* 1851: 1-22.
- Loew, 1858a. Bidrag till kännedommen om Afrikas Diptera [part]. *Öfvers. k. svenska Vetenskapsakad. Förhandl.* (1857) 14: 337-383.
- Loew, H., 1858. Bericht über die neueren Erscheinungen auf dem Gebiete der Dipterologie. *Berlin. ent. Ztschr.* 2: 325-349.
- Loew, H., 1859. Bidrag till kännedommen om Afrikas Diptera [part]. *Öfvers. k. svenska Vetenskapsakad. Förhandl.* 15: 335-341.
- Loew, H., 1860. Die Dipteren-Fauna Südafrika's. Erste Abtheilung. *Abh. Naturwiss. Ver. Sachsen Thüringen*, Halle (1858-1861) 2: 57-402, 2 pls. [Also sep. publ., xi + 330 pp., 2 pls., Berlin, 1860].
- Loew, H., 1861. Diptera aliquot in insula Cuba collecta. *Wien. ent. Monatschr.* 5: 33-43.
- Loew, H., 1866. Diptera Americae septentrionalis indigena. Centuria septima. *Berlin. ent. Ztschr.* 10: 1-54. [Also sep. publ., pp. 61-114, 1872b].
- Loew, H., 1872a. Diptera Americae septentrionalis indigena. Centuria decima. *Berlin. ent. Ztschr.* 16: 49-115. [Also sep. publ., pp. 225-291, 1872b].
- Loew, H., 1872b. *Diptera Americae septentrionalis indigena*. 2 (Centuriae 6-10), 300 pp., Belin.
- Loew, H., 1874. Neue nordamerikanische Dasypogonina. *Berlin. ent. Ztschr.* 18: 353-377.
- Lynch Arribálzaga, E., 1879. Asilides argentinos. *An. Soc. cient. argentina* 8: 145-153 [cont.].
- Lynch Arribálzaga, E., 1880. Asilides argentinos [cont.]. *An. Soc. cient. argentina* 9: 26-33, 49-57, 224-230, 252-265 [cont.].
- Lynch Arribálzaga, E., 1881. Asilides argentinos [cont.]. *An. Soc. cient. argentina* 11: 17-32, 112-128 [cont.].
- Lynch Arribálzaga, E., 1882a. Asilides argentinos [cont.]. *An. Soc. cient. argentina* 13: 186-192 [cont.].
- Lynch Arribálzaga, E., 1882b. Asilides argentinos [cont.]. *An. Soc. cient. argentina* 14: 132-143 [cont.].
- Lynch Arribálzaga, E., 1883a. Catálogo de los dípteros hasta ahora descritos que se encuentran en las repúblicas del río de la Plata [cont.]. *Boln Acad. nac. Cienc. Córdoba* 4: 109-152.
- Lynch Arribálzaga, E., 1883b. Asilides argentinos [concl.]. *An. Soc. cient. Argentina* 15: 5-18, 79-90. [Note: although this is not the conclusion of Lynch Arribálzaga's paper, the last part was never published; his "Asilides argentinos" was left incomplete].
- Macquart, J., 1834. *Histoire naturelle des insectes. Diptères* 1: 578 pp., 12 pls. [in Roret, N. E., ed., Collection des suites à Buffon], Paris.
- Macquart, J., 1838. *Diptères nouveaux ou peu connus* 1 (2): 5-207, 14 pls. Paris. [Also publ. in *Mém. Soc. r. Sci.*

- Agric. Arts Lille* 1838 (3): 121-323, 14 pls., 1839].
- Macquart, J., 1846. Diptères exotiques nouveaux ou peu connus. 1er. Supplément. *Mém. Soc. r. Sci. Agric. Arts Lille* (1845) 1844: 133-364, 20 pls. [Also publ. sep., as Supplément I, pp. 5-238, 20 pls., Paris, 1846].
- Macquart, J., 1848. Diptères exotiques nouveaux ou peu connus. Suite de 2me. supplément [i. e., 3e. supplément]. *Mém. Soc. r. Sci. Agric. Arts Lille* 1847 (2): 161-237, 7 pls. (Also sep. publ., as Supplément III; 1-77, 7 pls., Paris, 1848).
- Macquart, J., 1850. Diptères nouveaux ou peu connus. 4e. supplément (part). *Mém. Soc. r. Sci. Agric. Arts Lille* 1849: 307-465 (text), 466-479 (expl. of figs., index), pls. 1-14. [Also publ. sep., pp. 5-161, Paris].
- Macquart, J., 1855. Diptères exotiques nouveaux ou peu connus. 5e. supplement. *Mém. Soc. imp. Sci. Agric. Arts Lille* 1854: 25-156, 7 pls. [Also sep. publ., pp. 5-136, 7 pls., Paris, 1855].
- Maia, V. C., M. S. Couri, C. J. E. Lamas M., D. Pamplona & C. C. de C. Aires, 2002. A survey of the Diptera fauna of Serra do Cipó (Minas Gerais, Brazil). *Entomologist's monthly Mag.* 138: 109-117.
- Maldonado-Capriles, J. & C. A. Navarro, 1967. Additions and corrections to Wolcott's "Insects of Puerto Rico". *Caribbean J. Sci.* 7 (1-2): 45-64.
- Malloch, J. R., 1915. Some additional records of Chironomidae from Illinois and notes on other Illinois Diptera. *Bull. Illinois St. Lab. Nat. Hist.* 11: 305-363, pls. 80-84.
- Malloch, J. R., 1917. A preliminary classification of Diptera, exclusive of Pupipara, based upon larval and pupal characters, with keys to imagines in certain families. Part 1. *Bull. Illinois State Lab. Nat. Hist.* 12 (3): v + pp. 161-409, pls. 28-57. [Asilidae, pp. 373-389, pls. 53-55].
- Marcangeli, J., 1998. Notes on the behaviour of the robber-fly *Mallophora ruficauda*, preying on honey bees in Argentina. *Bee World*, London 79 (2): 69-70.
- Martin, C. H., 1955a. Notes on the genus *Haplopogon* and a new species (Diptera: Asilidae). *Ohio J. Sci.* 55 (5): 315-316.
- Martin, C. H., 1955b. New species in the genus *Parataracticus* Cole from Southern California (Diptera: Asilidae). *J. Kansas ent. Soc.* 28: 116-120.
- Martin, C. H., 1957a. A revision of the Leptogastrinae in the United States (Diptera: Asilidae). *Bull. American Mus. nat. Hist.* 3: 347-385.
- Martin, C. H., 1957b. The Asilidae of the Bahama Islands, with the description of two new species (Diptera). *American Mus. Novitates* 1847: 1-7, 5 figs.
- Martin, C. H., 1959. The *Holopogon* complex of North America, excluding Mexico, with descriptions of a new genus and subgenus (Diptera, Asilidae). *American Mus. Novitates* 1980: 1-40, 21 figs.
- Martin, C. H., 1960. A new species of *Microstylum* (Diptera: Asilidae) from Mexico. *J. Kansas ent. Soc.* 33: 44-45.
- Martin, C. H., 1961a. A revision of the genus *Lissoteles* (Diptera, Asilidae). *American Mus. Novitates* 2027: 1-13, 4 figs.
- Martin, C. H., 1961b. The misidentification of *Erax* Scopoli in the Americas (Diptera: Asilidae). *J. Kansas ent. Soc.* 34: 1-4.
- Martin, C. H., 1962a. The mistaken identity of *Proctacanthus arno* Townsend, and a new species (Diptera, Asilidae). *J. Kansas ent. Soc.* 35: 185-188.
- Martin, C. H., 1962b. Changes in the status of *Efferia barbata* (Fab.) and *Efferia albobarbis* (Macq.) (Diptera: Asilidae). *J. Kansas ent. Soc.* 35 (2): 247-253.
- Martin, C. H., 1962c. The plumose hairs of *Heteropogon* and two new species from Mexico (Diptera: Asilidae). *J. Kansas ent. Soc.* 35: 317-377.
- Martin, C. H., 1963. The designation of lectotypes, neotypes, and redescriptions of Bellardi's species of *Diogmites* from Mexico (Diptera: Asilidae). *Boll. Ist. Mus. Zool. Univ. Torino* 6 (1958-1962): 3-31.
- Martin, C. H., 1965a. Distribution patterns and corrected identification of asilid species reported as common to North and South America (Diptera: Asilidae). *Trans. American ent. Soc.* 91: 1-37, 3 maps.
- Martin, C. H., 1965b. Generic and subfamily changes, new synonymy, a new species, and notes on Asilidae (Diptera). *J. Kansas ent. Soc.* 38: 110-134, 23 figs.
- Martin, C. H., 1966a. The genus *Townsendia* Williston in Mexico (Diptera: Asilidae). *J. Kansas ent. Soc.* 39 (3): 542-551.
- Martin, C. H., 1966b. New Asilidae from Mexico in the genera *Itolia* and *Sphageus* (Diptera). *Pan-Pacific Entomologist* 42: 212-218, 8 figs.
- Martin, C. H., 1966c. The status of *Andrenosoma* Rondani, *Pilica* Curran, and *Pogonosoma* Rondan, with two new species (Diptera: Asilidae). *Pan-Pacific Entomologist* 42: 323-328.
- Martin, C. H., 1967a. The Mexican species of *Promachus* in the subgenus *Amblyonychus* (= *Promachina*) (Diptera: Asilidae). *J. Kansas ent. Soc.* 40: 156-163.
- Martin, C. H., 1967b. New Asilidae from Mexico and Arizona (Diptera). *Pan-Pacific Entomologist* 43: 193-201, 4 figs.
- Martin, C. H., 1968a. The new family Leptogastridae (the grass flies) compared with the Asilidae (robber flies) (Diptera). *J. Kansas ent. Soc.* 41 (1): 70-100, 137 figs.
- Martin, C. H., 1968b. Revision of *Stenopogon* Loew (= *Scleropogon* Loew) as represented in Mexico (Diptera: Asilidae). *Proc. California Acad. Sci.* 35 (17): 371-400, 35 figs.
- Martin, C. H., 1968c. A revision of *Ospriocerus* (Diptera: Asilidae). *Proc. California Acad. Sci.* 35 (18): 401-424, 66 figs.
- Martin, C. H., 1968d. New Mexican *Acronyches* and *Parataracticus* (Diptera: Asilidae). *Pan-Pacific Entomologist* 44 (4): 179-183.
- Martin, C. H., 1968e. Family Leptogastridae, in Papavero, N., ed., *A catalogue of the Diptera of the Americas south of the United States* 25b: 1-11. Departamento de Zoologia, Secretaria da Agricultura, São Paulo.
- Martin, C. H., 1971. A review of the genus *Leptopteromyia* of the Western Hemisphere (Diptera, Leptogastridae). *Pan-Pacific Entomologist* 47: 264-270.
- Martin, C. H., 1972. A new species of *Leptopteromyia* (Diptera: Leptogastridae). *Pan-Pacific Entomologist* 48 (4): 270.
- Martin, C. H., 1975a. Revision of *Systellogaster* (Diptera: Leptogastridae). *Opuscula zoologica* 134: 1-5.
- Martin, C. H., 1975b. Review of the genus *Schildia* Aldrich (Diptera: Leptogastridae). *Proc. ent. Soc. Washington*

- 77: 189-193.
- Martin, C. H., 1975c. The generic and specific characters of four old and six new Asilini genera in the western United States, Mexico and Central America (Diptera: Asilidae). *Occas. Papers California Acad. Sci.* 119: 1-107, 78 figs., 1 table, 1 map.
- Martin, C. H. & N. Papavero, 1970. Family Asilidae, in Papavero, N., ed., *A catalogue of the Diptera of the Americas south of the United States* 35b: 1-139. Museu de Zoologia, Universidade de São Paulo, São Paulo.
- Martin, C. H. & J. Wilcox, 1965. Asilidae, pp. 360-401, in Stone, A., C. W. Sabrosky, W. W. Wirth & J. R. Coulson, eds., *A catalog of the Diptera of America North of Mexico. [U. S. Dept. Agric., Agric. Res. Serv.] Handbook 276*: 1696 pp. Washington, D. C.
- McAtee, W. L., 1919. Key to the Nearctic species of the genus *Laphria* (Diptera, Asilidae). *Ohio J. Sci.* 19 ("1918"): 143-170, pls. 10-11.
- Meigen, J. W., 1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*, 40 pp. Paris.
- Meigen, J. W., 1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkunde* 2: 259-281.
- Meigen, J. W., 1820. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten* 2: x + 365 pp., pls. 12-21. Aachen.
- Melander, A. L., 1923. The genus *Lasiopogon* (Diptera: Asilidae). *Psyche* 30: 135-145.
- Melander, A. L., 1947. Some fossil Diptera from Florissant, Colorado. *Psyche* 53: 43-48.
- Morgan, K. R. & T. E. Shelly, 1988. Body temperature regulation in desert robber flies (Diptera: Asilidae). *Ecol. Ent.* 14: 419-428.
- Morgan, K. R., T. E. Shelly & L. S. Kinsey, 1985. Body temperature regulation, energy metabolism, and foraging in light-seeking and shade-seeking robber flies. *J. comp. Physiol. (B)* 155: 561-570.
- Naveiro, J., 1975. Factores que inciden en la pérdida de reinas. *Gaceta del Colmenar [Revista de la Sociedad Argentina de Apicultores]* 37 (4).
- Nelson, C. R., 1987. Robber flies of Utah (Diptera: Asilidae). *Great Basin Naturalist* 47 (1): 38-90.
- Neuhaus, G. H., 1886. *Diptera marchica. Systematisches Verzeichniss der Zweiflügler der Mark Brandenburg mit kurzer Beschreibung und analytischen Bestimmungs-Tabellen*, 371 pp. Berlin.
- Newman, E., 1834. Attempted division of British insects into natural orders. *Ent. Mag.* 2: 379-431.
- Notario, A., J. F. Michela, D. C. Fiorentino & L. Castresana, 2000. Contribución al conocimiento de *Schinopsis quebracho-colorado* Bark y Meyer y de sus predadores asociados. *Boln San. Vegetal. Plagas* 26 (1): 15-20.
- Oldroyd, H., 1964. The genus *Pseudorus* (Diptera: Asilidae). *Ann. Mag. nat. Hist.* (13) 7: 5-11, 6 figs.
- Oldroyd, H., 1965. Rediscovery of *Dasypecus* Philippi (Diptera: Asilidae). *Ann. Mag. nat. Hist.* 13 (8): 351-355.
- Olivier, C. A., 1789. *Encyclopédie méthodique. Histoire naturelle. Insectes* 4: 1-331. Paris.
- O'Neill, K. M., 1992. Body size assymmetries in predatory interactions among robber flies (Diptera: Asilidae). *Ann. ent. Soc. America* 85: 34-38.
- O'Neill, K. M. & W. P. Kemp, 1992. Behavioral thermoregulation in two species of robber flies occupying different grasslands microhabitats. *J. thermal Biol.* 17 (6): 323-331.
- Osten Sacken, C. R., 1874. A list of the Leptidae, Mydaidae and Dasypogonina of North America. *Bull. Buffalo Soc. nat. Hist.* 2: 169-187.
- Osten Sacken, C. R., 1877. Art. XIII. Western Diptera: Descriptions of new genera and species of Diptera from the region west of the Mississippi and especially from California. *Bull. U. S. geol. geogr. Surv. Terr.* 3: 180-354.
- Osten Sacken, C. R., 1878. Catalogue of the described species of Diptera of North America. Ed. 2. *Smithson. misc. Collns* 16: x + 276 pp.
- Osten Sacken, C. R., 1887. Diptera, pp. 129-160, 161-176, 177-208, 209-216, pl. 3, in Godman, F. D. & O. Salvin, eds., *Biologia centrali-americana. Zoologia-Insecta-Diptera* 1: 378 pp., 6 pls. London.
- Osten Sacken, C. R., 1891. Additions and corrections to the Catalogue of the described species of South American Asilidae by S. W. Williston. *Berlin. ent. Ztschr.* 36: 417-428.
- Osterberger, H. A., 1930. *Erax interruptus* Macq. as a predator. *J. econ. Ent.* 23: 709-711.
- Özdikmen, H., 2006. Replacement names for some Asilidae genera (Diptera). *Munis Entomology & Zoology*, Ankara 1 (1): 93-96.
- Pamplona, D. M. & C. C. de C. Aires, 1999. Revision of *Triorla* Parks and description of new species, emphasizing the systematic value of the spermathecae (Diptera, Asilidae). *Revta brasileira Zool.* 16 (4): 1127-1134.
- Papavero, N., 1971a. Neotropical *Acronyches* (Diptera, Asilidae), new or otherwise. *Papéis Avulsos Zool.*, São Paulo 23 (16): 145-151.
- Papavero, N., 1971b. A new Mexican *Saropogon* (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 23 (17): 153-156.
- Papavero, N., 1971c. Uma nova espécie de *Aphamartania* Schiner, do Panamá (Diptera, Asilidae). *Papéis avulsos Zool.*, São Paulo 24 (2): 53-56, 3 figs.
- Papavero, N., 1971d. Notes on some types of Neotropical Asilidae (Diptera), with descriptions of three new species. *Papéis avulsos Zool.*, São Paulo 25 (3): 19-29, 6 figs.
- Papavero, N., 1971e. *Essays on the history of Neotropical Dipterology, with special reference to collectors (1750-1905)* 1: vii + 216 pp., 10 pls., 12 maps. Museu de Zoologia, Universidade de São Paulo, São Paulo.
- Papavero, N., 1973a. Studies of Asilidae (Diptera) systematics and evolution. I. A preliminary classification in subfamilies. *Argos Zool.*, São Paulo 23 (3): 217-274, 3 figs.
- Papavero, N., 1973b. Studies of Asilidae (Diptera) systematics and evolution. II. The tribes of Dasypogoninae. *Argos Zool.*, São Paulo 23 (4): 275-294, 3 figs.
- Papavero, N., 1973d. *Essays on the history of neotropical Dipterology, with special reference to collectors (1750-1905)*, 2: iii + pp. 217-446. Museu de Zoologia, Universidade de São Paulo, São Paulo.

- Papavero, N., 1975. Studies of Asilidae (Diptera) systematics and evolution. IV. Tribe Megapodini Carrera (Dasypogoninae), with a review of the Neotropical species. *Argos Zool.*, São Paulo 26 (3): 191-318, 175 figs.
- Papavero, N. & N. Bernardi, 1973. Studies of Asilidae (Diptera) systematics and evolution. III. Tribe Blepharepiini (Dasypogoninae). *Argos Zool.*, São Paulo 24 (3): 163-209, 1 pl., 41 figs.
- Papavero, N. & N. Bernardi, 1974. *Archilestris* Loew in the Americas, with the description of two new neotropical species (Diptera, Asilidae, Stenopogoninae). *Papéis avulsos Zool.*, São Paulo 27 (20): 263-278, 20 figs.
- Papavero, N. & S. Ibáñez-Bernal, 2001. Contributions to a history of Mexican Dipterology. –Part I. Entomologists and their work before the *Biología Centrali-Americana*. *Acta zool. mex.* (n. s.) 84: 65-173, 10 figs.
- Papavero, N. & S. Ibáñez-Bernal, 2003. Contributions to a history of Mexican Dipterology. Part II. The *Biología Centrali-Americana*. *Acta zool. mex.* (n. s.) 88: 143-232, 4 figs.
- Paramonov, S. J., 1964. Notes on Australian Diptera (XXXVIII-XL). Notes on Australian Asilidae, with the description of two new genera. XL. The presence of the American genus *Eccritosia* Schin. (Asilidae) in Australia, and the absence of the genus *Philodicus* Lw. *Ann. Mag. nat. Hist.* (13) 7: 157-159.
- Paramonov, S. J., 1966. Notes on Australian Diptera (XL- XLIII). Notes on three genera of Australian Asilidae, with the description of seven new species. *Ann. Mag. nat. Hist.* (13) 9: 137-144.
- Parks, L., 1968. Synopsis of robberfly genera allied to *Efferia* and *Eicherax*, including a new genus. *Pan-Pacific Entomologist* 44 (3): 171-179.
- Perty, M., 1833. Insecta brasiliensia, pp. 125-224, pls. 25-40, in his *Delectus animalium articulatorum quae in itinere per Brasiliam annis MDCCCXVII-MDCCCXX jussu et auspiciis Maximiliani Josephi I. Bavariae Regis augustissimi peracto collegerunt Dr. J. B. de Spix et Dr. C. F. Ph. de Martius*, 44 + 224 pp., 40 pls. Monachii [= München], "1830-1834" [pp. 1-60, pls. 1-2, 1830; pp. 61-124, pls. 13-24, 1832].
- Philippi, R. A., 1865. Aufzählung der chilenischen Dipteren. *Verh. zool.-bot. Ges. Wien* 15: 595-782.
- Poinar, G. O., Jr., 1992. *Life in amber*, xiii + 350 pp. Stanford University Press, Stanfrod, California.
- Pritchard, A. E., 1935. New Asilidae from the southwestern United States (Diptera). *American Mus. Novitates* 813: 1-13.
- Pritchard, A. E., 1938a. Synopsis of North and Central American *Holcocephala* with a description of a new species (Diptera: Asilidae). *J. New York ent. Soc.* 46: 11-21.
- Pritchard, A. E., 1938b. Revision of the robberfly genus *Taracticus* Loew, with descriptions of three new species (Diptera: Asilidae). *J. New York ent. Soc.* 46: 179-190.
- Pritchard, A. E., 1938c. The genus *Hodophylax* James, with a description of *basingeri*, new species (Diptera, Asilidae). *Pan-Pacific Entomologist* 14: 129-131, 1 fig.
- Pritchard, A. E., 1941a. *Annamyia*, a new genus of Asilidae, with a revision of the genus *Aphamartania* Schiner (Diptera). *Proc. ent. Soc. Washington* 43 (6): 131-140, 1 fig., pl. 16.
- Pritchard, A. E., 1941b. The genus *Haplopogon* in the New World, with the description of *erinus* n. sp. *Ann. ent. Soc. America* 34: 350-354.
- Pritchard, A. E., 1942. A revision of the genus *Cerotainiops* Cresson (Diptera, Asilidae). *J. Kansas ent. Soc.* 15: 19-24.
- Pritchard, A. E., 1943. Revision of the genus *Cophura* Osten Sacken (Diptera, Asilidae). *Ann. ent. Soc. America* 36: 281-309, 1 pl.
- Rabinovich, M. & J. C. Corley, 1997. An important new predator of honey bees. The robber fly *Mallophora ruficauda* Wiedemann (Diptera – Asilidae) in Argentina. *American Bee J.* 137 (4): 303-306.
- Rabinovich, R., R. Quiroga & M. K. Castelo, 1997. Moscardón y chaqueta amarilla. Importantes avances e investigaciones en Luján. *Espacio apícola* 7 (27): 18-21.
- Remedi de Gavotto, A. L., 1964. Ciclo biológico de *Cyclocephala signaticollis* Burm. (Coleoptera, Scarabaeidae) y caracteres específicos de su larva. [Instituto Nacional de Teconología Agropecuaria] *Revta Invest. Agropecuarias*, Buenos Aires (5) 1 (10): 151-161.
- Ricardo, G., 1927. Notes on the two genera *Nusa* and *Pogonosoma* (Laphriinae). *Ann. Mag. nat. Hist.* 20: 205-212.
- Ricardo, G., 1929. *Insects of Samoa* 6 (3) (Stratiomyidae, Tabanidae and Asilidae): 298-364. British Museum (Natural History), London.
- Rinaldi, A. J. M., L. A. Pailhé & E. R. Popolizio, 1971. Un díptero cazador de abejas en Trucumán: El *Eicherax ricnotes* Engel (Asilidae). *Revta agron. Noroeste argentino*, San Miguel de Tucumán 8 (3-4): 451-455.
- Roeder, V. von, 1885. Dipteren der Insel Portorico. *Stettin. ent. Ztg* 46: 337-349.
- Roeder, V. von, 1887. Ueber die Gattungen *Doryclus* Jaen. und *Megapoda* Mcq. *Berlin. ent. Ztschr.* 31: 76-78.
- Roeder, V. von, 1900. *Tricloscelis*, nov. gen. dasypogoninorum (Diptera). *Stettin. ent. Ztg* 61: 337-340.
- Rondani, C., 1848. Esame di varie specie d'insetti ditteri brasiliani. [Truqui's] *Studi entomologici*, Torino 1: 63-112.
- Rondani, C., 1850a. Osservazioni sopra alquante specie di esapodi ditteri del Museo Torinese. *Nuovi Ann. Sci. nat. Bologna* 2 (3): 165-197, 1 pl.
- Rondani, C., 1850b. Dipterorum species aliquae in America Aequatoriali collectae a Cajetano Osculati, observatae et distinctae novis breviter descriptis. *Nuovi Ann. Sci. nat. Bologna* (3) 2: 357-372.
- Rondani, C., 1856. *Dipterologiae italicae prodromus*. Vol. 1. *Genera italica ordinis dipterorum ordinatim disposita et in familias et stirpes aggregata*, 228 pp. Parmae [= Parma].
- Rondani, C., 1861. *Dipterologiae italicae prodromus* 4: 1-174. Parmae [= Parma].
- Rondani, C., 1863. *Diptera exotica revisa et annotata*, 99 pp., 1 pl. Modena. [Also publ. under the title "Dipterorum species et genera aliqua exotica", in *Arch. Zool. Anat. Fisiol. Modena* (1863) 3 (1): 1-99, pl. 5, 1864].
- Rondani, C., 1868. Diptera aliqua in America Meridionali laeta a Prof. A. Strobel, annis 1866 et 1867. *Ann. Soc. Naturalisti Modena* 3: 24-40, pl. 4.
- Rondani, C., 1873. *Muscaria exotica Musei Civivi Januensis observata et distincta*. *Ann. Mus. civ. Stor. nat. Genova* 4: 282-294, 295-300.

- Rueda, M. C., 1998. Contribución al conocimiento de la tribu Atomosiini (Diptera: Asilidae: Laphriinae): Redescripción de *Cerotainia leonina* Hermann, 1912 y descripción de una especie nueva. *Acta zool. illoana* 44 (1): 167-175, 3 figs..
- Rueda, M. C., 2003a. Análisis de la diversidad de Atomosiini (Diptera, Asilidae, Laphriinae) en el noroeste de la República Argentina. *Revista Unlar Ciencia* 4 (2): 7-12.
- Rueda, M. C., 2003b. Asilidae (Diptera) neotropicales. Tribu Atomosiini en Argentina. *Revista Unlar Ciencia* 4 (2): 13-19.
- Rueda, M. C., 2004. Género *Cerotainia* Schiner, 1866 (Diptera, Asilidae, Laphriinae, Atomosiini) en América del Sur: Redescripción de *Cerotainia jolyi* Kaletta y nuevos datos sobre *Cerotainia sarae* Rueda. *Revta cient. Univ. nac. La Rioja* 5 (1): 2-7.
- Sabrosky, C. W., 1999. Family-group names in Diptera. An annotated catalog. Bibliography by F. Christian Thomson, Neal L. Evenhuis & Curtis W. Sabrosky. *Myia* 10: 1-576. [North American Dipterists' Society & Backhuys Publishers, Leiden].
- Say, T., 1823. Descriptions of dipterous insects in the United States. *J. Acad. nat. Sci. Philadelphia* 3: 9-54, 73-104.
- Say, T., 1824. Appendix. Part I. – Natural history. 1. Zoology. E. Class Insecta, pp. 268-378, n Keating, W. H., *Major Long's second expedition* 2: 459 pp., pls. 6-15; 3: 47-53, 73-74. Philadelphia.
- Say, T., 1829. Descriptions of North American dipterous insects. *J. Acad. nat. Sci. Philadelphia* (1829-1830) 6: 149-178.
- Say, T., 1859. In Le Conte, J. L., ed., *The complete writings of Thomas Say on the entomology of North America* 1: 412 pp., 2: 814 pp.
- Scarborough, A. G., 1984a. Four species of *Ommatius* Wiedemann (Diptera: Asilidae) from Puerto Rico and the Virgin Islands. *Proc. ent. Soc. Washington* 86: 619-629.
- Scarborough, A. G., 1984b. Synopsis of *Ommatius* Wiedemann (Diptera: Asilidae) from Hispaniola. *J. New York ent. Soc.* 92: 131-149.
- Scarborough, A. G., 1985a. New *Ommatius* Wiedemann (Diptera: Asilidae) from Cuba and the Bahamas. *J. New York ent. Soc.* 93: 1226-1239.
- Scarborough, A. G., 1985b. *Ommatius* (Diptera: Asilidae) in the Lesser Antilles. *Proc. ent. Soc. Washington* 87: 641-655.
- Scarborough, A. G., 1988a. A new species of *Efferia* Coquillett (Diptera: Asilidae), *staminea* species group, from Grand Cayman Island, West Indies. *Florida Entomologist* 71 (2): 150-153.
- Scarborough, A. G., 1988b. New robber flies (Diptera: Asilidae) from Mayaguana Island, Bahamas and Cuba. *Ent. News* 99 (2): 90-94.
- Scarborough, A. G., 1990. Revision of the New World *Ommatius* Wiedemann (Diptera: Asilidae). I. The *pumilus* species group. *Trans. American ent. Soc.* 116: 65-102.
- Scarborough, A. G., 1993. Revision of the New World species of *Ommatius* Wiedemann (Diptera: Asilidae): The neotropical *costatus* species group. *Revta Biol. trop.*, San José 41 (3): 729-753.
- Scarborough, A. G., 1996. The genus *Leptogaster* (Diptera: Asilidae) in the West Indies. *Ent. News* 107 (4): 193-206.
- Scarborough, A. G., 1997a. New and old species of *Ommatius* Wiedemann (Diptera: Asilidae) from Hispaniola. *Insecta mundi* 11 (1): 9-24.
- Scarborough, A. G., 1997b. West Indian species of *Beameromyia* Martin (Diptera: Asilidae). *Insecta Mundis* 11 (3-4): 237-246.
- Scarborough, A. G., 2000. Two additional species of robber flies of the genus *Ommatius* Wiedemann (Diptera: Asilidae) from the Bahamas with replacement names for two other species. *Proc. ent. Soc. Washington* 102 (4): 912-918.
- Scarborough, A. G., 2002. Synopsis of the *holosericeus* complex of the genus *Ommatius* Wiedemann (Diptera: Asilidae): *ampliatus* and *holosericeus* species groups. *Trans. American ent. Soc.* 128 (2-3): 133-222.
- Scarborough, A. G., 2007. A new species of *Ommatius* Wiedemann from Brazil (Diptera: Asilidae) with notes on the *Ommatius costatus* species group. *Ann. ent. Soc. America* 133 (3-4): 465-472, 10 figs.
- Scarborough, A. G., 2008. New *Ommatius* Wiedemann from the Americas with two new species groups, keys, and taxonomic notes (Diptera: Asilidae). *Insecta Mundis* 32: 1-14, 48 figs.
- Scarborough, A. G. & L. V. Knutson, 1989. Asilidae, Bombyliidae, Conopidae, and Pipunculidae (Diptera) of Dominica (West Indies). *Florida Entomologist* 72: 519-537.
- Scarborough, A. G. & D. E. Perez-Gelabert, 2003. *Ommatius* Wiedemann (Diptera: Asilidae) from Hispaniola: Five new species and a modified key, phenology, and distribution records of species. *Boln Soc. ent. aragonesa* 33: 41-58, 51 figs.
- Scarborough, A. G. & D. E. Perez-Gelabert, 2005. Review of *Wilcoxius* Martin (Diptera: Asilidae) with five new species. *Boln Soc. ent. aragonesa* 36: 41-49.
- Scarborough, A. G. & D. E. Perez-Gelabert, 2006. A review of the asilid (Diptera) fauna from Hispaniola with six genera new to the island, fifteen new species, and checklist. *Zootaxa*, Auckland 1381: 1-91.
- Scarborough, A. G., D. E. Perez-Gelabert & S. A. Page, 2005. Synopsis of leptogastrines (Diptera: Asilidae) from Hispaniola. *Trans. American ent. Soc.* 131: 29-67.
- Scarborough, A. G. & D. E. Perez-Gelabert, 2009. Review of the West Indian species of *Efferia* Coquillett (Diptera: Asilidae) with 13 new species and checklist: Part II. Hispaniola, Puerto Rico, and Lesser Antilles including Tobago and Trinidad. *Zootaxa*, Auckland 1994: 1-66.
- Scarborough, A. G. & G. O. Poinar, Jr., 1992. Upper Eocene robber flies of the genus *Ommatius* (Diptera: Asilidae) in Dominican amber. *Insecta Mundis* 6: 13-18.
- Scarborough, A. G. & P. R. Rutkauskas, 1983. A new species of *Ommatius* (Diptera: Asilidae) from San Salvador, the Bahamas. *Proc. ent. Soc. Washington* 85: 144-151.
- Scarborough, A. G. & B. E. Sraver, 1979. Predatory behavior and prey of *Atomosia puelia* (Diptera: Asilidae). *Proc. ent. Soc. Washington* 81: 630-639.
- Schaeffer, C., 1916. New Diptera of the family Asilidae with notes on known species. *Trans. ent. Soc. America* 35: 65-69.
- Schiner, I. R., 1860. Vorläufiger Commentar zum dipterologischen Theile der "Fauna austriaca", mit einer näheren Begründung der in derselben

- aufgenommenen neuern Dipteren-Gattungen. *Wien. ent. Monatschr.* 4: 208-216.
- Schiner, I. R., 1862. *Fauna austriaca*. Theil I, Hefl 8 [part], pp. 65-674 + I-lxxx. Wien.
- Schiner, I. R., 1866. Die Wiedemann'sche Asiliden, interpretiert und in die seither errichteten neuen Gattungen eingereiht. *Verh. zool.-bot. Ges. Wien* 16 (Abhandl.): 649-722, pl. 12, pp. 845-848 (Nachtrag).
- Schiner, I. R., 1867. Neue oder wenige bekannte Asiliden der k. zoologische Hofskabinetes in Wien. Ein Beitrag zur Kenntniss der Asiliden. *Verh. zool.-bot. Ges. Wien* 17: 355-412.
- Schiner, I. R., 1868. Diptera, pp. 1-388, 4 pls., in [Wüllersdorf Urbair, B. von, in charge], *Reise der österreichischen Fregatte Novara um die Erde. Zoologie* 2 (1, B). Wien.
- Schinz, H. R., 1823. *Das Thierreich eingetheilt nach den Bau der Thiere als Grundlage ihrer Naturgeschichte und der vergleichenden Anatomie*. 3. Krebsen, Spinnen, Insekten, xviii + 932 pp. J. G. Cotta, Stuttgart & Tübingen.
- Schumann, H., 1973. Die Asiliden-Typen der Dipteren-Sammlung des Zoologisches Museums in Berlin. *Mitt. zool. Mus. Berlin* 49 (1): 81-174.
- Scudder, S. H., 1884. Nomenclator zoologicus. Part I. Supplemental list of genera in zoology. *Bull. U. S. natl. Mus.* 19 (1): 1-367.
- Shelly, T. E., 1984a. Prey selection by the Neotropical robber fly, *Atractia marginata* (Diptera: Asilidae). *Proc. ent. Soc. Washington* 86: 120-126.
- Shelly, T. E., 1984b. Comparative foraging behavior of Neotropical robber flies (Diptera: Asilidae). *Oecologia*, Berlin 62: 188-195.
- Shelly, T. E., 1985a. Observations on the reproductive behavior of two neotropical robber fly species (Diptera: Asilidae). *Proc. ent. Soc. Washington* 87: 253-254.
- Shelly, T. E., 1985b. Ecological comparisons of robber fly species (Diptera: Asilidae) coexisting in a Neotropical forest. *Oecologia*, Berlin 67: 57-70.
- Shelly, T. E., 1986a. Rates of prey consumption by Neotropical robber flies (Diptera: Asilidae). *Biotropica* 18: 166-170.
- Shelly, T. E., 1986b. Foraging success of Neotropical robber flies: Variation with attack. distance and angle. *Pan-Pacific Entomologist* 62: 124-127.
- Shelly, T. E., 1987. Natural history of three riparian species of robber flies in a Panamanian forest (Diptera: Asilidae). *Biotropica* 19: 180-184.
- Shelly, T. E. & D. L. Pearson, 1980. Predatory behavior of *Proctacanthella leucopogon* (Diptera: Asilidae): Prey recognition and prey records. *Environm. Ent.* 9: 7-9.
- Shelly, T. E. & D. L. Pearson, 1983. Diurnal variation in the predatory behavior of the grassland robber fly, *Proctacanthella leucopogon* (Williston) (Diptera: Asilidae). *Pan-Pacific Entomologist* 57: 380-384.
- Shelly, T. E. & D. A. Weinberger, 1981. Courtship and diet of the neotropical robber fly, *Mallophora schwarzi* Curran (Diptera: Asilidae). *Pan-Pacific Entomologist* 57: 380-384.
- Snow, W. A., 1896. List of Asilidae, supplementary to Osten Sacken's catalogue of North American Diptera, 1878-1895. *Kansas Univ. Quart.* 4: 173-190.
- Soria, S. de J. & A. F. Dal Conte, 2005. Ecologia e controle das pragas da videira. *EMRAPA* [Empresa Brasileira de Pesquisa Agropecuária] *Circ. técn.* 63: 1-20, 14 figs.
- Soria, S. de J. & R. P. de Mello, 1998. Ocorrência de *Prolepsis lucifer* (Wiedemann, 1828) (Diptera, Asilidae) no sul do Brasil, com anotações morfológicas sobre larvas e pupas. *Entomología y Vectores* 5 (6): 279-294.
- Soria, S. de J., R. P. de Mello & A. M. de Oliveira, 2004. Novos registros de *Prolepsis lucifer* (Wiedemann, 1828) (Diptera, Asilidae) como predador de *Eurhizococcus brasiliensis* (Hempel in Wille, 1922) (Hemiptera, Margarodidae) em diferentes regiões vitícolas do Rio Grande do Sul. *Entomología y Vectores*, Rio de Janeiro 11 (2): 323-331.
- Sousa Sánchez, A., 1969. *Colecciones botánicas de C. A. Purpus en México, periodo 1898-1925*. University of California Press, Berkeley.
- Speiser, P., 1920. *Hexameritia*, nom. nov. for *Eutrichodes* Hermann. *Zool. Jahrb.* 43: 447.
- Strand, E., 1928. Miscellanea nomenclatorica zoologica et paleontologica. *Arch. Naturgesch.* 92 (1926): 30-75.
- Stuardo Ortiz, C., 1946. *Catálogo de los dipteros de Chile*, 1253 pp. Ministerio de Agricultura, Santiago.
- Thomson, C. G., 1869. 6. Diptera. Species novas descripsit..., pp. 443-614, pl. 9 [= h. 12, no. 2], in K. Vetenskaps-Akademien, *Kongliga svenska fregatten Eugenies resa omkring jorden* 2 (Zoologie), [Sec.] 1 (Insekter): 617 pp., 9 pls., Stockholm, "1868".
- Titus, E. S. G. & F. C. Pratt, 1905. Catalogue of the exhibit of economic entomology at the Louisiana Purchase Exposition, 1904. *Bull. U. S. Dept. Agric., Div. Ent.* 47: 1-155.
- Tomasovic, G., 2002a. Résumé de la connaissance des Asilidae (Diptera Brachycera) de Guyane Française et description de trois espèces nouvelles. *Bull. Soc. belge Ent.* 138: 85-90.
- Tomasovic, G., 2002b. Etude sur les types d'*Erax* Scopoli, 1763 de Macquart et description de 4 nouvelles espèces (Diptera: Asilidae). *Notes faun. Gembloux* 49: 51-83, 20 figs.
- Tomasovic, G. & Y. Braet, 2001. Description d'une nouvelle espèce de *Pseudorus* Walker, appartenant à un complexe mimétique néotropical (Diptera, Asilidae). *Lambillonea* 101 (3): 379-385.
- Townsend, C. H. T., 1895. On the Diptera of Baja California, including some species from adjacent regions. *Proc. California Acad. Sci.* (2) 4: 593-620.
- Tricca, G. & O. Trujillo-Cenóz, 1980. The sexual behavior of the robber-fly *Mallophora ruficauda* Wied. (Asilidae, Asilinae) in normal and experimental conditions. *Physiology and Behavior* 25 (5): 627-632.
- Valdés-Raguès, P., 1908. Dípteros cubanos. *An. Acad. Cienc. Cuba*, Habana 45: 302-319.
- Vieira, R., I. Castro, D. Almeida, E. Alvim & F. Bravo, 2006. Asilidae (Diptera) da Bahia, Brasil: Sinopse das espécies e chave de identificação. *Sitientibus, Ser. Ciênc. Biol.*, Feira de Santana 6 (4): 243-260, 14 figs.
- Vieira, R., I. Castro & F. Bravo, 2005a. Two new species of *Ommatius* Wiedemann (Diptera: Asilidae) from Brazil. *Zootaxa*, Auckland 764: 1-7, 13 figs.
- Vieira, R., I. Castro & F. Bravo, 2005b. A new species of *Ommatius* Wiedemann (Diptera: Asilidae) from Brazil. *Zootaxa*, Auckland 1017: 19-24, 7 figs.

- Walker, F., 1837. Descriptions of Diptera collected by Capt. King in the survey of Straits of Magellan. *Trans. Linn. Soc. London* 17: 331-359.
- Walker, F., 1848. *List of the specimens of dipterous insects in the collection of the British Museum* 1: 1-229. London.
- Walker, F., 1849. *List of the specimens of dipterous insects in the collection of the British Museum* 2: 231-484. London.
- Walker, F., 1850. Characters of undescribed Diptera in the British Museum. *Zoologist* 8 (Appendix): lxv, xcv-xcix, cxxl.
- Walker, F., 1851. *Insecta Saundersiana, or characters of undescribed insects in the collection of William Wilson Saunders, Esq.* 1: 76-156, 2 pls. London.
- Walker, F., 1854. *List of the specimens of dipterous insects in the collection of the British Museum, Suppl.* 3: 507-775, 5 pls. London.
- Walker, F., 1855. *List of the specimens of dipterous insects in the collection of the British Museum* 6. (Suppl. II): 331-506. British Museum, London.
- Walker, F., 1856. *Insecta Saundersiana, or characters of undescribed insects in the collection of William Wilson Saunders, Esq.* 1: 415-474. London.
- Walker, F., 1857. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. ent. Soc. London* (2) (= ser. 2) 4: 119-158.
- Walker, F., 1858. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. ent. Soc. London* (2) (= ser. 2) 4: 190-235.
- Walker, F., 1860. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. ent. Soc. London* (2) (= ser. 2) 5: 268-296.
- Walker, F., 1861. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. ent. Soc. London* (2) (= ser. 2) 5: 297-334.
- Weeks, L. & A. H. Hespenheide, 1985. Predatory and mating behavior of *Stichopogon* (Diptera: Asilidae) in Arizona. *Pan-Pacific Entomologist* 61: 95-104.
- Westwood, J. O., 1835. Insectorum nonnullorum novorum (ex ordine dipterorum) descriptions. *Ann. Soc. ent. France* (1) 4: 681-585.
- Wiedemann, C. R. W., 1817. Neue Zweiflügler (Diptera Linn.) aus der Gegend um Kiel. [Wiedemann's] *Zool. Mag.* 1 (1): 61-68.
- Wiedemann, C. R. W., 1819. Brasilianische Zweiflügler. [Wiedemann's] *Zool. Mag.* 1 (3): 40-56.
- Wiedemann, C. R. W., 1821. *Diptera exotica*, 244 pp., 1 fig., 2 pls. Kiliae [=Kiel].
- Wiedemann, C. R. W., 1824. *Munus rectoris in Academia Christiana Albertina aditurus. Analecta entomologica ex Museo Regio Havniensi maxime congesta*, 60 pp., 1 pl. Kiliae [=Kiel].
- Wiedemann, C. R. W., 1828. *Aussereuropäische zweiflügelige Insekten* 1: xxxii + 608 pp., 7 pls. Hamm.
- Wiedemann, C. R. W., 1830. *Aussereuropäische zweiflügelige Insekten* 2: xii + 684 pp., 5 pls. Hamm.
- Wilcox, J., 1935. New asilid flies of the genus *Ablautus* with a key to the species. *Canadian Entomologist* 67: 222-227.
- Wilcox, J., 1936a. New *Ommatius* with a key to the species (Diptera, Asilidae). *Bull. Brooklyn ent. Soc.* 31 (4): 172-177, pl. 10.
- Wilcox, J., 1936b. Asilidae, new or otherwise, from the Southwest, with a key to the genus *Stichopogon* [sic]. *Pan-Pacific Entomologist* 12 (4): 201-212, 1 fig.
- Wilcox, J., 1937. Asilidae, new or otherwise, from the Southwest, with a key to the genus *Stictopogon* [sic]. *Pan-Pacific Entomologist* 13 (1-2): 37-45, 1 fig.
- Wilcox, J., 1941. New *Heteropogon* with a key to the species (Diptera, Asilidae). *Bull. Brooklyn ent. Soc.* 36 (2): 50-56.
- Wilcox, J., 1949. The genus *Itolia* Wilcox (Diptera: Asilidae). *Pan-Pacific Entomologist* (1948) 24: 191-193.
- Wilcox, J., 1959. The *clausa* group of *Cophura* Osten Sacken (Diptera: Asilidae). *Bull. Brooklyn ent. Soc.* 54 (4): 121-127, 1 fig.
- Wilcox, J., 1960. *Laphystia* Loew in North America (Diptera: Asilidae). *Ann. ent. Soc. America* 53: 328-346, pls. 1-2.
- Wilcox, J., 1961. The genus *Hodophylax* James (Diptera: Asilidae). *Bull. Brooklyn ent. Soc.* 56: 112-116.
- Wilcox, J., 1964. The genus *Metapogon* (Diptera: Asilidae). *Pan-Pacific Entomologist* 48 (1): 191-200.
- Wilcox, J., 1965a. *Proctacanthella* Bromley (Diptera: Asilidae). *Bull. Kansas ent. Soc.* 38 (2): 106-110, 1 fig.
- Wilcox, J., 1965b. New *Cophura*, with revised keys to the groups and to the Nearctic species (Diptera: Asilidae), *Ann. ent. Soc. America* 58 (6): 805-816, 1 fig.
- Wilcox, J., 1965c. New *Heteropogon* Loew, with a key to the species and description of a new genus (Diptera: Asilidae). *Bull. South California Acad. Sci.* 64 (4): 207-222.
- Wilcox, J., 1966a. New species and a key to the species of *Saropogon* Loew. *Pan-Pacific Entomologist* 42: 127-136.
- Wilcox, J., 1966b. New species of *Haplopogon* Loew with a key to the species (Diptera: Asilidae). *Bull. South California Acad. Sci.* 63 (2): 99-106.
- Wilcox, J., 1966c. *Efferia* Coquillett in America north of Mexico (Diptera: Asilidae). *Proc. California Acad. Sci.* 39 (2): 85-234, 197 figs.
- Wilcox, J., 1966d. New species and keys to the species of *Ablautus* Loew and *Omnablautus* Pritchard (Diptera: Asilidae). *Canadian Entomologist* 98: 673-682.
- Wilcox, J., 1971. The genera *Stenopogon* Loew and *Scleropogon* Loew in America north of Mexico (Diptera: Asilidae). *Occas. Papers California Acad. Sci.* 89: 1-134, 263 figs.
- Wilcox, J., 1972. The genus *Sintoria* Hull (Diptera: Asilidae). *Pan-Pacific Entomologist* 48 (1): 51-58.
- Wilcox, J. & C. H. Martin, 1936a. A review of the genus *Cyrtopogon* Loew in North America (Diptera—Asilidae). *Entomologica americana* (n. s.) 16 (1): 1-95, 5 pls.
- Wilcox, J. & C. H. Martin, 1936b. The genus *Nannocyrtopogon* (Diptera, Asilidae). *Ann. ent. Soc. America* 29 (3): 449-459, 1 pl.
- Wilcox, J. & C. H. Martin, 1941. The genus *Dioctria* Meigen in North America (Diptera – Asilidae). *Entomologica americana* 21: 1-22, pl. 1.
- Wilcox, J. & C. H. Martin, 1942. *Nannodiocetria* n. n. for *Neodiocetria* Wilcox & Martin (nec Ricardo). *Bull. Brooklyn ent. Soc.* 37: 35.
- Wilcox, J. & C. H. Martin, 1957. *Nannocyrtopogon* (Diptera – Asilidae). *Ann. ent. Soc. Amer.* 50 (4): 376-392, 2 figs.
- Williston, S. W., 1883. On the North American Asilidae (Dasypogoninae, Laphriinae), with a new genus of Syrphidae. *Trans. American ent. Soc. & Proc. Acad.*

Catalogue of the Neotropical Diptera. Asilidae

- nat. Sci. Philadelphia, Ent. Sect.* (1884) 11: 1-35.
Williston, S. W., 1885. On North American Asilidae (Part II).
Trans. American ent. Soc. and Proc. Acad. nat. Sci. Philadelphia, Ent. Sect. 12: 53-76.
Williston, S. W., 1886. Dipterological notes and descriptions.
Trans. American ent. Soc. and Proc. Acad. nat. Sci. Philadelphia, Ent. Sect. 13: 287-307.
Williston, S. W., 1889. Notes on Asilidae. *Psyche* 5: 225-259.
Williston, S. W., 1891. Catalogue of the described species
of South American Asilidae. *Trans. American ent. Soc.* 18: 67-91.
Williston, S. W., 1893a. New or little known Diptera. *Kansas Univ. Quart.* 2: 59-78.
Williston, S. W., 1893b. A list of species of Diptera from San
Domingo. *Canadian Entomologist* 25: 170-171.
Williston, S. W., 1894. On the genus *Erax*. *Ent. News* 5: 136-137.
Williston, S. W., 1895. Two remarkable new genera of Diptera.
Kansas Univ. Quart. 4: 107-109.
Williston, S. W., 1896a. On the Diptera of St. Vincent (West
Indies). *Trans. ent. Soc. London* 1896: 253-446.
Williston, S. W., 1896b. *Manual of the families and genera
of North American Diptera*. Ed. 2, 167 pp. New Haven,
Connecticut.
Williston, S. W., 1897. Diptera brasiliiana. Part IV. *Kansas Univ. Quart.* (A) 6: 1-12.
Williston, S. W., 1901. Supplement (part), pp. 249-264,
265-272, 273-296, 297-328, 329-332, pls. 4-5, pl. 6, figs.
1-6, in Godman, F. D. & O. Salvin, eds., *Biologia
centrali-americana. Zoologia-Insecta-Diptera* 1:
378 pp., 6 pls. London.
Williston, S. W., 1907. Dipterological notes. *J. New York ent. Soc.* 13: 1-2.
Williston, S. W., 1908. *Manual of North American Diptera*.
Ed. 3, 405 pp., 163 figs. New Haven, Connecticut.
Wolcott, G. N., 1948. The insects of Porto Rico. *J. Agric. Univ. Porto Rico* 32 (3): 417-748.
Woodley, N. E. & D. J. Hilburn, 1994. The Diptera of
Bermuda. *Contr. American ent. Inst.*, Ann Arbor 28 (2):
10 [Asilidae].
Wulp, F. M. van der, 1869. Nog iets over noord-
amerikaansche Diptera. *Tijdschr. Entr.* (2) 4 (12): 80-86.
Wulp, F. M. van der, 1870. Opmerkingen omtrent uitlandsche
Asiliden. *Tijdschr. Ent.* (2) 5 (13) ("1869") 13: 207-217, 1 pl.
Wulp, F. M. van der, 1872. Bijdragen tot de kennis der
Asiliden van der Oost-Indische Archipel. *Tijdschr. Ent.*
(2) 7 (15): 129-279, pls. 9-12.
Wulp, F. M. van der, 1882a. Remarks on certain American
Diptera in the Leyden Museum and description of nine
new species. *Notes Leyden Mus.* 4: 73-92.
Wulp, F. M. van der, 1882b. Amerikaansche Diptera. *Tijdschr. Ent.* 25: 77-163, pls. 9-10.
Wulp, F. M. van der, 1884. Nalezing over amerikaansche
Diptera. *Tijdschr. Ent.* 27: 140-142.
Wulp, F. M. van der, 1896. *Catalogue of the described
Diptera from South Asia*, iii + 219 pp. Hague.
Zack, R. S., 1984. List of types in the James entomological
collection. *Melanderia* 42: 1-41.
Zimsen, E., 1954. The insect types of C. R. W. Wiedemann
in the Zoological Museum in Copenhagen. *Spolia zool.
Mus. hauniensis* 14: 1-43.
Zimsen, E., 1964. *The type material of I. C. Fabricius*, 656
pp., 2 pls. Munksgaard, Copenhagen.

Index (Synonyms in *italics*)

- abana* Curran, 1934, Mallophora – 38
abana Curran, 1953, Ommatius – 47
abdominales Brown of authors, Cerotainiops, misident. – 82
abdominalis (Say), 1824 (Asilus), Osprioncerus – 127
abdominalis (Wiedemann), 1821 (Asilus), Nerax – 19
aberrans (Wiedemann), 1821 (Dasypogon), Diogmites – 64
Ablautatus, emend. – 112
Ablautus Loew, 1866 – 112
ablayoensis (Hengst & Artigas), 2003 (Philonerax),
 Lochmorhynchus, error – 32
Acanthocneminae Hermann, 1912 – 75
Acanthodelphia Bigot, 1867 – 54
acapulcae Martin, 1961, Lissoteles – 133
acapulcae Pritchard, 1943, Cophura – 63
acapulquensis (Cole & Pritchard), 1964 (Carreraia),
 Carreraomyia – 35
achaetus Scarbrough, 1993, Ommatius – 47
aciculatus Pritchard, 1938, Taracticus – 61
aconaguana Artigas, Lewis & Parra, 2005, Tillobroma – 131
Acronymes Williston, 1866 – 112
Acronymchini Artigas & Papavero, 1991 – 111
acrus (Curran), 1931 (Mallohorā), Megaphorus – 44
acuminatus Carrera, 1959, Eraxasilus – 26
acunai (Bromley), 1929 (Andrenosoma), Pilica – 83
acus Martin, 1957, Apachekolos – 106
acuta (Artigas), 1980 (Myaptex), Myaptexaria – 46
acuticornis Carrera, 1945, Hybozelodes – 94
acutulus Martin, 1975, Wilcoxius – 47
acutus Martin, 1975, Machimus – 9
acutus Scarbrough, 1990, Ommatius – 47
Aczelia Carrera, 1955 – 57
ada Curran, 1941, Mallophora – 40
adustus Loew, 1874, Saropogon – 59
aeaca Williston, 1901, Mallophora – 35
aeacides Loew, 1866, Osprioncerus – 127
aeacidinus (Williston), 1886 (Stenopogon), Osprioncerus – 127
aeacus (Wiedemann), 1828 (Dasypogon), Osprioncerus – 127
aedon (Walker), 1849 (Dasypogon), Senobasis – 79
aegra Martin, 1957, Leptogaster – 108
aemula Hull, 1962, Ommatius, error – 48
aenescens Hermann, 1912, Oidardis – 95
aestuans authors, not Linnaeus, Pogoniefferia – 23
aestuans-group of Hine, 1919, Erax – 19
aestuans-group of Wilcox, 1966, Efferia – 19
affinis (Bellardi), 1861 (Dasypogon), Diogmites – 69
affinis (Bellardi), 1861 (Discocephala), Holcocephala – 135
affinis (Bellardi), 1861 (Erax), Nerax – 19
affinis (Fabricius), 1805 (Laphria), Lampria – 99
affinis (Macquart), 1850 (Atomosia), Aphestia – 85
affinis Schiner, 1868, Mallophora – 43
agalla (Walker), 1849 (Discocephala), Holcocephala – 135
ajax Curran, 1941, Mallophora – 43
alamoensis Martin, 1968, Osprioncerus – 127
alba (Martin), 1975 (Systelogaster), Systologaster – 111
albalulus Martin, 1968, Stenopogon – 128
albescens (Schiner), 1868 (Erax), Cratolestes – 17
albescens (Schiner), 1868 (Erax), Lochmorhynchus – 33
Albibarbefferia Artigas & Papavero, 1997 – 14
albibarbis (Curran), 1930 (Protichisma), Cerotainia – 90
albibarbis-group of Wilcox, 1966 (Efferia) – 14
albibarbis Hull, 1962, Labromyia – 57
albibarbis (Macquart), 1838 (Erax), Albibarbefferia – 15
albicans (Carrera & d'Andretta), 1953 (Erax), Lochmorhynchus – 32
albiceps (Hermann), 1912 (Atonia), Atoniomyia – 88
albicincta Bromley, 1929, Mallophora – 38
albicollis Bigot, 1878, Dasyllis – 82
albidus Scarbrough & erez-Gelabert, 2006, Rhopalagaster – 100
albifacies (Hermann), 1912 (Atonia), Atoniomyia, error – 88
albifacies Williston, 1885, Promachus – 44
albifasciata (Back), 1904 (Osprioncerus), Prolepsis – 124
albifrons Back, 1904, Saropogon – 59
albifrons Walker, 1850, Mallophora – 36
albimanus (Carrera), 1949 (Mirolestes), Cyrtophrys – 75
albinigrus Artigas, 1981, Lochmorhynchus – 32
albipes Hermann, 1912, Hybozelodes – 94
albipilosus Macquart, 1846 (Asilus) – 138
albisetosus Wulp, 1882 (Epitriptus) – 56
albispinosus (Macquart), 1850 (Erax), Lochmorhynchus – 32
albitarsis (Macquart), 1846 (Dasypogon), Archilestris – 119
albitibia Macquart, 1848 (Laphria) – 101
albitibialis Macquart, 1847 (Laphria) – 101
?alboatra (Walker), 1851 (Discocephala), Holcocephala – 136
albolineatum (Macquart), 1850 (Laphria), Andrenosoma – 80
albolineatus (Macquart), 1850 (Laphria), Anarmostus – 7
albomacula Martin, 1966, Townsendia – 134
albomarginatus (Carrera), 1949 (Lastauroides), Neodiogmites – 74
albospinosus (Bellardi), 1861 (Asilus, ?Epitretus), Philonicus – 11
alcanor Walker, 1849, Laphria – 100
alecto Rondani, 1848, Mallophora – 41
aleto Martin & Papavero, 1970, Mallophora, error – 42
alexanderi (Carrera), 1949 (Lastauroides), Neodiogmites – 74
alexanderi Farr, 1965, Ommatius – 48
alexanderi Papavero, 1971, Acronymes – 112
alia Scarbrough & Perez-Gelabert, 2009, Efferia – 17
Alpiolaphria Carrera, 1951 – 96
Alexipogon Curran, 1934 – 133
Allopogon Schiner, 1886 – 67
almeida Carrera, 1946, Senobasis – 79
alphus Martin, 1975, Schildia – 110
alpuyea (Cole & Pritchard), 1964 (Carreraia), Carreraomyia – 35
alterus (Williston), 1901 (Asilus (Tolmerus)), Machimus – 9
alticola Carrera & Machado-Allison, 1968, Lastaurus – 73
Alvarenga Carrera, 1960 – 62
alvesi Carrera, 1949, Diogmites – 69
Alyssomyia Hull, 1962 – 118
amanda (Aldrich), 1905 (Laphria), Neophoneus, error – 101
amandus (Walker), 1849 (Laphria), Neophoneus – 101
amapa Artigas, Papavero & Serra, 1991. Dissmeryngodes – 92
amarynceus (Walker), 1849 (Asilus), Nerax – 21
amazona Carrera, 1955, Austenmyia – 58
amazonicus (Bromley), 1934 (Erax), Nerax – 19
ambigua (Macquart), 1846 (Erax), Triorla – 23
amblayoensis (Hengst & Artigas), 2003 (Philonerax),
 Lochmorhynchus – 32
Amblyonychus Hermann, 1921 – 33
americana Carrera, 1955, Apoxyria – 102
americanus Becker, 1919, Epipamponeurus – 8
americanus (Curran), 1930 (Damalis), Orhhodops – 138
americanus Macquart, 1846 (Lophonotus) – 56
amethystinus Carrera, 1955, Diogmites – 69
Amorimius, nom. nov. – 58
amphinome (Walker), 1849 (Asilus), Eccritosia – 53
amphissa (Walker), 1849 (Erax), Eraxasilus – 26
ampliatus Scarbrough, 2002, Ommatius – 48
Ampyx Walker, 1855 – 77
amula Curran 1928, Ommatius – 48
anacaona Scarbrough & Perez-Gelabert, 2006, Atomosia – 85
analisis Jaennicke, 1867, Nicocles – 65

- analis (Macquart), 1846 (Discocephala), Holcocephala – 135
 analis Macquart, 1838, Senobasis – 79
Anarmostus Loew, 1860 – 7
anceps (Hermann), 1912 (Pholidotus), Joartigasia – 98
anceps Osten Sacken, 1887, Promachus – 44
ancylocera (Schiner), 1868 (Atomosia), Atonomyia – 88
andrenoides Bromley, 1934, Aphestia – 85
Andrenosoma Rondani, 1856 – 80
Andrenosomatini Hull, 1962 – 80
Andrenosomatini Sabrosky, 1999 – 81
Andrenosomini Hull, 1962 – 80
angulata Artigas, Lewis & Parra, 2005, Tillobroma – 131
angulosus Scarbrough, 2002, Ommatius – 48
angusta (Kertész), 1909 (Dolichodes), Plesiomma, error – 126
angusta Scarbrough & Perez, 2009, Efferia – 17
angustatus Scarbrough, 2002, Ommatius – 69
angustipennis Loew, 1866, Diogmites – 69
angustiventris (Macquart), 1846 (Dasypogon), Diogmites – 69
angustum (Macquart, 1848 (Dasypogon), Plesiomma – 125
Anisopogon Loew, 1874 – 114
Annamyia Pritchard, 1941 – 62
annularis (Fabricius), 1805 (Dasypogon), Amblyonychus – 33
annulatum (Bigot), 1857 (Senobasis), Blepharepium – 66
annulatus (Say), 1823 (Leptogaster), Psilonyx – 110
annulipes (Macquart), 1838 (Atomosia), Aphestia – 84
annulipes (Macquart), 1846 (Erax), Lecania – 30
annulipes Walker, 1855, Leptogaster – 108
annulitarsis (Rondani), 1868 (Dasypogon), Caenarolia – 68
annuliventris Artigas & Angulo, 1980, Mallophora – 36
anomala (Bellardi), 1861 (Erax), Efferia – 17
anomala (Carrera), 1957 (Diogmites), Caenarolia – 67
anomala-group of Wilcox, 1966, Efferia – 17
anomalus-group of Hine, 1919, Erax – 17
anonyma Williston, 1901, Atomosia – 85
anonymus (Williston), 1901 (Asilus (Neoitamus)),
 Prolatiforceps – 12
antennata (Banks), 1920 (Atomosia), Atomosiella – 88
anthracinus (Loew), 1851 (Dasypogon), Lastaurus – 73
antica Curran, 1934, Mallophora – 43
anticus (Wiedemann), 1828 (Laphria), Dissmeryngodes – 92
antidomus (Walker), 1849 (Asilus), Taurhynchus – 55
antiphus Walker, 1849 (Asilus) – 138
antiqua Walker, 1855, Mallophora – 36
Apachekolos Martin, 1957 – 106
apatma Hull, 1962, Cophura, error – 63
aper (Walker), 1855 (Erax), Carinefferia – 16
Aphamartania Schiner, 1866 – 62
Aphestia Schiner, 1866 – 84
Aphractia Artigas, Papavero & Serra, 1991 – 85
apicalis (Bellardi), 1861 (Asilus), Ommatius – 48
apicalis (Bromley), 1934 (Pachychaeta), Lecania – 31
apicalis Enderlein, 1914, Leptogaster – 108
apicalis Schiner, 1868, Ommatius – 52
apicalis (Schiner), 1867 (Lochites), Senobasis – 79
?apicalis (Walker), 1854 (Discocephala), Holcocephala – 135
apicalis (Wiedemann), 1821 (Asilus), Nerax – 19
apicata Schiner, 1866, Apoxyria – 102
Apocleinae Papavero, 1973 – 7
Apocliina Lehr, 1969 – 7
Apolastauroides Artigas & Papavero, 1988 – 65
Apotinocerus Hull, 1962 – 32
apotma Pritchard, 1943, Cophura – 63
Apoxyria Schiner, 1866 – 102
appendiculata Macquart, 1846 (Atomosia) – 101
appendiculatum (Macquart), 1846 (Laphria), Andrenosoma – 80
aquilonius Martin, 1961, Lissoteles – 133
arachnoides Bigot, 1878, Pogonosoma – 84
arauguensis Kaletta, 1976, Townsendia – 134
Araiopogon Carrera, 1949 – 59
†Araripogon Grimaldi, 1990 – 138
Araucopogon Artigas & Papavero, 1988 – 58
Araujoa Artigas & Papavero, 1991 – 118
araujoi Carrera, 1952, Rhopalagaster – 100
arawak Farr, 1963, Psilonyx – 110
Archilestes Schiner, 1866 – 118
Archilestris Loew, 1874 – 119
Archilestrodes Artigas & Papavero, 1991 – 119
ardens Macquart, 1834, Mallophora – 36
ardens Macquart of Artigas & Angulo, 1980, Mallophora,
 misident. – 36
ardens (Wiedemann), 1828 (Dasypogon), Lastaurina – 72
arenicola James, 1937, Leptogaster – 108
arenicolus Martin, 1968, Parataracticus – 59
argentifascia Bromley, 1946, Diplosynapsis, error – 17
argentifacies James, 1953, Taracticus – 61
argentifacies (Williston), 1901 (Triclis), Perasis – 105
argentifascia Enderlein, 1914, Diplosynapsis – 17
argentifrons (Hine), 1911 (Erax), Efferia – 17
argentina (Wulp), 1882 (Laparus), Aczelia – 57
argentinae Martin, 1972, Leptopteromyia – 110
argentinae Martin, 1971, Leptopteromyia, nom. nud. – 110
argentinus Artigas & Papavero, 1990, Argyropogon – 133
argentipes Macquart, 1838, Mallophora – 36
argirogaster Tomasovic, 2002, Triorla, error – 24
argutus Fisher, 1977, Zabrops – 106
argyrata Curran, 1926, Townsendia – 134
argyrocincta (Schiner), 1867 (Saropogon), Caenarolia – 67
argyrogaster (Macquart), 1846 (Erax), Triorla – 24
argyropasta Hermann, 1912, Cerotainia – 90
argyrophora Schiner, 1868, Atomosia – 85
Argyropogon Artigas & Papavero, 1990 – 133
argyropus Schiner, 1868, Cerotainia – 90
argyropyga Hermann, 1912, Cerotainia – 90
aria Curran, 1941, Mallophora – 42
arida-group of Wilcox, 1966, Efferia – 16
aridus Curran, 1930, Saropogon – 60
aridus Scarbrough, 2002, Ommatius – 48
aristata Carrera, 1950, Glaphyropyga – 28
Aristofolia Ayala, 1978 – 30
arizonensis (Bromley), 1937 (Stenopogon), Ospricerus – 127
arizonensis (Schaeffer), 1916 (Lasiocnemus), Cophura – 63
arizonensis (Williston), 1892 (Senoprosopis), Philonicus – 11
armata Artigas & Angulo, 1980, Mallophora – 36
armata Hermann, 1912, Atomosia – 85
armata (Hine), 1918 (Erax), Albabarbefferia – 15
arnaudi Martin, 1968, Stenopogon – 128
arnaudi Wilcox, 1966, Ablautus – 113
arno Townsend, 1895, Proctacanthus – 54
arrayanensis Artigas, 1970, Dasycyrton – 113
Arthriticopus Enderlein, 1914 – 135
Asicya Lynch Arribgálzaga, 1880 – 102
Asiliidae Harris, 1833 – 7
Asilica Burmeister, 1829 – 7
Asilici Latreille, 1802 – 7
Asilidae Kirby & Spence, 1817 – 7
Asilidae Latreille, 1802 – 7
Asilides Leach, 1815 – 7
asiliformis (Wulp), 1882 (Hypenetes), Tillobroma – 131
Asilina Newman, 1834 – 7
Asilinae Latreille, 1802 – 7
Asilini Schinz, 1823 – 7
Asilitae Macquart, 1838 – 7
Asilites Newman, 1834 – 7
asiloides (Bigot), 1978 (Lochites), Caenarolia – 68

- Asilopsinae Cockerell, 1920 – 7
Asilus-group – 7
Aspidopyga Carrera, 1949 – 63
Astylum Rondani, 1850 – 78
Astylum Kertész, 1909, emend. – 78
atennatus Williston, 1891, *Cyrtophrys*, error – 75
Atomasinae Engel, 1929 – 84
Atomosia Macquart, 1838 – 85
Atomosiaria Lynch Arribálzaga, 1882 – 84
Atomosiella Wilcox, 1937 – 88
Atomosiinae G. H. Hardy, 1928 – 84
Atomosiini Lynch Arribálzaga, 1882 – 84
Atomosinae Hermann, 1912 – 84
Atonia Williston, 1889 – 88
Atonomyia Hermann, 1912 – 88
atra Macquart, 1834, *Mallophora* – 36
Atractia Macquart of Hermann, 1912, part, misident. – 94
Atractia Macquart, 1838 – 89
Atractocoma Artigas, 1970 – 45
atrata (Philippi), 1865 (*Dasypogon*), *Lamprozona* – 95
atratus (Bigot), 1878 (*Diogmites*), *Lastaurus* – 73
atriapex Carrera, 1943, *Diogmites* – 69
atriapex (Carrera & Papavero), 1962 (*Lastauroides*)
 Neodiogmites – 74
atribarbis Artigas, 1970, *Stizolestes* – 13
atrifrons Cole, 1924, *Holopogon* – 115
atripes Fabricius, 1805 (*Dasypogon*) – 138
atripes Wilcox, 1949, *Itolia* – 115
atrosus Scarbrough, 1997, *Ommatius* – 48
atrum Bromley, 1929, *Plesiomma* – 125
attenuata (Hull), 1958 (*Opopotes*), *Glyphyropyga* – 28
attenuatus (Loew), 1851 (*Dasypogon* (*Cyrtophrys*)),
 Cyrtophrys – 75
atypha Pritchard, 1943, *Cophura* – 63
audax McAtee, 1919, *Laphria* – 100
audouinii Macquart, 1838 (*Gonypes*) – 138
augusta (Kertész), 1909 (*Dolichodes*), *Plesiomma*, error – 125
aurantiaca Hermann, 1926, *Systelogaster*, nom. nud. – 111
aurarius Wiedemann, 1821 (*Dasypogon*) – 138
aurata (Enderlein), 1914 (*Laphria*), *Smeryngolaphria* – 97
aurata Schiner, 1868, *Cerotainia* – 90
aureolus Carrera & Papavero, 1962, *Diogmites* – 69
aureomaculatus (Bromley), 1932 (*Asilus*), *Stizolestes* – 13
auribarbis (Wiedemann), 1821 (*Asilus*), *Nerax* – 19
auricincta (Loew), 1851 (*Laphria* (*Automolina*)), *Lamprozona* – 94
auricinctum (Schiner), 1867 (*Senobasis*), *Blepharepium* – 66
aurifer Hermann, 1912, *Rhopalogaster* – 100
aurifex Osten Sacken, 1887, *Lampria* – 98
aurivestitus Hine, 1919 (Erax) – 56
aurolineatus (Macquart), 1846 (*Proctacanthus*), *Taurhynchus* – 55
auromystacea Macquart, 1855, *Mallophora* – 37
Austenmyia Carrera, 1955 – 58
australis Adisoemarto & Wood, 1975 (*Dioctria*), *Nannodioctria* – 117
austrinus Martin, 1961, *Lissoteles* – 133
Automolina Hermann, 1912 – 94
autumnalis Martin, 1961, *Lissoteles* – 133
aveledoi (Kaletta), 1978 (*Iucundus*), *Oidardis* – 95
avicenniae Farr, 1965, *Andrenosoma* – 80
avidus (Wulp), 1869 (*Asilus*), *Machims* – 9
†axelrodi Grimaldi, 1990, *Arariogon* – 138
ayala Kaletta, 1974, *Eumecosoma* – 93
ayalai Scarbrough, 2002, *Ommatius* – 48
Aymarasilus Artigas, 1974 – 119

bacillifera Carrera, 1949, *Cleptomyia* – 58
badia Martin, 1975, *Polacantha* – 11
badiapex (Bromley), 1928 (Erax), *Nerax* – 20

bahamiensis Scarbrough, 1996, *Leptogaster* – 108
baleta (Walker), 1849 (*Asilus*), *Lecania* – 31
baletus Walker, 1849 (*Asilus*) – 138
balmacedensis Artigas, 1970, *Lochyrus* – 8
barbata (Fabricius), 1787 (*Asilus*), *Eccritosia* – 53
barbata (Fabricius), 1805 (*Dasypogon*), *Pogoniefferia* – 23
barbatus-group of Hine, 1919 (Erax) – 14
barbiellinii (Curran), 1935 (*Promachina*), *Amblyonychus* – 35
barbiellinii Curran, 1935, *Atomosia* – 85
barbiellinii (Curran), 1935 (*Mirolestes*), *Cyrtophrys* – 75
barbiellinii Curran, 1934, *Eccritosia* – 53
barbiellinii Curran, 1934, *Ommatius* – 48
barbiellinii Curran, 1934, *Porasilius* – 23
barbipes (Wiedemann), 1819 (*Laphria*), *Mallophora* – 37
barbiventris (Rondani), 1850 (*Asilus*), *Taurhynchus* – 55
bardyllis (Walker), 1849 (*Asilus*), *Nerax* – 19
barrettoi (Carrera), 1949 (*Macrocolus*), *Amorimius* – 58
basalis (Curran), 1935 (*Allopongon*), *Caenarolia* – 67
basalis Curran, 1930, *Bathropsis* – 90
basalis Walker, 1855, *Leptogaster* – 108
basifascia Walker, 1855, *Proctacanthus* – 54
basingeri Pritchard, 1938, *Hodophylax* – 64
basingeri Wilcox & Martin, 1936, *Cyrtopogon* – 113
bassleri Curran, 1942, *Blepharepium* – 67
bassleri Curran, 1941, *Mallophora* – 37
bastardoanus Scarbrough, 2003, *Ommatius* – 48
batesi Bromley, 1931, *Andrenosoma* – 80
Bathropsis Hermann, 1912 – 90
Bathypogonini Artigas & Papavero, 1991 – 112
beameri Wilcox, 1936, *Ommatius* – 48
Beameromyia Martin, 1957 – 107
bechyneorum Ayala, 1982, *Holcocephala* – 135
beckeri Jaennicke, 1867, *Atomosia* – 86
belfragei (Hine), 1919 (Erax), *Efferia* – 17
bella Bromley, 1929, *Rhopalogaster* – 100
bella (Loew), 1897 (Blax), *Cophura* – 63
bella Schiner, 1867, *Cerotainia* – 90
bellardii (Bromley), 1919 (*Deromyia*), *Diogmites* – 69
bellardii (Schiner), 1868 (*Eristicus*), *Eicherax* – 25
bellardii Martin, 1965, *Promachus* – 44
bellengeri Farr, 1865, *Mallophora* – 37
bellingeri Martin & Papavero, 1970, *Mallophora*, error – 37
belzebul Schiner, 1868, *Mallophora* – 39
bengryi Farr, 1963, *Leptogaster* – 108
bequaerti Bromley, 1934, *Atomosia* – 86
?bergii Lynch Arribálzaga, 1880, *Mallophora* – 41
bevistylatus (Wulp), 1882 (*Proctacanthus*), *Apotinocerus* – 32
beviventris Kertész, 1909, *Aphamartania*, error – 62
bicincta (Loew), 1866 (*Dizonias*), *Prolepsis* – 124
bicincta Martin & Papavero, 1970, *Lampria*, error – 98
bicolor (Bellardi), 1861 (Erax), *Albibarbefferia* – 15
bicolor Bellardi, 1861, *Pseudorus* – 77
bicolor (Bigot), 1875 (*Emphysomera*), *Ommatius* – 48
bicolor (Engel), 1930 (*Macrocolus*), *Amorimius* – 58
bicolor (Jaennicke), 1867 (*Saropogon*), *Diogmites* – 69
bicolor (Macquart), 1848 (*Gonypes*), *Leptogaster* – 108
biezankoi (Carrera & Papavero), 1962 (*Lastauropsis*),
 Lastaurina – 72
bifascia Bromley, 1946, *Proctacanthus*, error – 54
bifasciatus Carrera, 1949, *Diogmites* – 69
bigoti Bellardi, 1861, *Atomosia* – 86
bigoti (Bellardi), 1861 (*Dasypogon*), *Diogmites* – 69
bigotii Lynch Arribálzaga, 1882, *Mallophora* – 37
bilineatus (Loew), 1866 (*Deromyia*), *Diogmites* – 69
bilineatus (Wulp), 1882 (Erax), *Nerax* – 19
bimaculata (Bellardi), 1861 (Erax), *Albibarbefferia* – 15
bimaculata Walker, 1855 (*Laphria*) – 101

- bimaculatus (Bromley), 1929 (Deromyia), Diogmites – 69
bipartitus Scarbrough, 1985, Ommatius – 48
bitincta Walker, 1860, Lampria – 98
Blacodes Loew, 1874 – 63
blaisdelli Wilcox, 1971, Stenopogon – 128
blantoni Martin, 1961, Lissoteles – 133
Blax Loew, 1872 – 63
Blepharepiini Papavero, 1973 – 65
Blepharepium Rondani, 1848 – 66
boharti Bromley, 1951, Stenopogon – 128
Bohartia Hull, 1958 – 118
bokermannii Artigas, Papavero & Costa, 1997, Protometer – 105
bolivari Ayala, 1983, Glaphyropyga – 28
bolivari Kaletta, 1986, Cerotainia – 90
bombimorpha (Rondani), 1950 (Dasipogon), Lastaurus – 74
bomboides (Wiedemann), 1821 (Asilus), Mallophora – 37
Bombomima Enderlein, 1914 – 99
bonariensis (Macquart), 1838 (Dasypogon), Blepharepium – 66
boraceae Carrera, 1958, Lecania – 31
boraceana Carrera, 1952, Senobasis – 79
borburatensis Ayala, 1981, Pteralbis – 12
Boropis Hull, 1962 – 125
borroii Schumann, 1973, Lochmorhynchus, error – 32
borrori Artigas, 1970, Lochmorhynchus – 32
brachyptera (Philippi), 1865 (Asilus), Myaptex – 46
brasiliæ Martin, 1971, Leptopteromyia – 110
brasiliensis Schiner, 1866, Aphestia – 85
brasiliensis Schiner, 1867, Cerotainia – 90
brasiliensis (Schiner), 1867 (Senoprosopis), Stenasilus – 13
brethesi (Gemignani), 1936 (Tolmerolestes), Prolepsis – 123
brevicornis (Macquart), 1838 (Laphria), Atomosia – 86
brevicornis (Philippi), 1865 (Clavator), Alyssomyia – 118
brevis Martin, 1975, Machimus – 9
brevis Martin, 1968, Ospricerus – 127
brevis Martin, 1975, Polacantha – 11
brevistylata (Williston), 1901 (Atonia), Atoniomyia – 88
breviusculus Loew, 1872, Stenopogon – 129
?*breviusculus* (Walker), 1855 (Trupanea), Amblyonychus – 34
breviventris (Macquart), 1848 (Dasypogon), Aphamartania – 62
breviventris Macquart, 1850, Mallophora – 39
?*breviventris* (Rondani), 1848 (Gonytes), Systologaster – 111
bromleyana Carrera, 1949, Senobasis – 79
bromleyanus (Carrera & d'Andretta), 1950 (Asilus), Machimus – 9
bromleyi Carrera, 1949, Diogmites – 69
bromleyi (Carrera & d'Andretta), 1950 (Dizonias), Prolepsis – 124
bromleyi Curran, 1930, Mallophora – 39
bromleyi (Curran), 1931 (Proctacanthus), Taurhynchus – 55
bromleyi D. E. Hardy, 1943, Nicocles – 65
bruneri Bromley, 1929, Mallophora – 41
brunnescens (Bromley), 1929 (Erax), Nerax – 19
brunneus (Fabricius), 1787 (Asilus), Diogmites – 69
bryanti Wilcox, 1966, Saropogon – 59
Brychomyia Hull, 1962 – 98
Buckellia Curran, 1925 – 63
bulbosa Artigas, 1970, Alyssomyia – 118
bullata Scarbrough & Perez-Gelabert, 2009, Efferia – 17
bullatus (Bromley), 1934 (Holopogon), Haplopogon – 135
bullatus (Bromley), 1929 (Asilus), Wilcoxius – 47
bullatus Scarbrough, 2002, Ommatius – 48
bullatus Wulp, 1882 (Holopogon) – 117
burmeisteri Roeder, 1900, Trichioscelis – 105
busiris Lynch Arribálzaga, 1881, Phonicocleptes – 75

Cacodaemon Schiner, 1866 – 122
Cacodaemonides Strand, 1927 – 122
caedens (Wiedemann), 1828 (Dasypogon), Plesioma – 125
Caenarolia Thomson, 1869 – 67

Caenorolia Williston, 1891, error – 67
caeruleiventris Macquart, 1846 (Asilus) – 139
caeruleum Scarbrough & Perez-Gelabert, 2006, Eumecosoma – 93
cajennense (Fabricius), 1787 (Asilus), Blepharepium – 66
?*calatina* (Walker), 1849 (Asilus), Lecania – 31
calcarata Williston, 1901, Aphestia, error – 85
calceata Schiner, 1867, Aphestia – 85
calceolata (Bigot), 1878 (Lasiocnemus), Systologaster – 111
calderense Artigas, 1970, Rhadinosoma – 46
calida (Fabricius), 1787 (Asilus), Mallophora – 38
caliente (Wilcox), 1966 (Efferia), Carinefferia – 16
californicus Wilcox, 1935, Ablautus – 113
calla Pritchard, 1943, Cophura – 63
callida Curran, 1934, Mallophora, error – 38
Callinicus Loew, 1872 – 113
calogastra (Philippi), 1865 (Laphria), Andrenosoma – 81
calopyga (Schiner), 1868 (Cylindrophora), Theromyia – 65
calverti Hine, 1917, Eumecosoma – 93
?*caminarium* (Wiedemann), 1828 (Dasypogon), Plesioma – 126
campestris Curran, 1941, Mallophora – 37
camposi Curran, 1931, Andrenosoma – 81
camposi Curran, 1934, Cerotainia – 90
camposi Curran, 1931, Psilocurus – 105
camposi (Curran), 1934 (Proctacanthus), Taurhynchus – 55
campsonianus (Curran), 1931 (Erax), Nerax – 19
cana (Hine), 1916 (Erax), Pogoniofferia – 22
cancerae Martin, 1975, Machimus – 9
candens (Walker), 1851 (Trupanea), Mallophora – 38
candida Coquillett, 1893, Efferia – 17
candidus (Macquart), 1846 (Dasypogon), Stichopogon – 134
cannella (Bromley), 1934 (Erax), Pogoniofferia – 22
canus (Hine), 1919 (Erax), Pogonefferia – 22
capillatus (Williston), 1901 (Asilus (Neoitamus)), Prolatiforceps – 12
capnoptera (Wiedemann), 1828 (Dasypogon), Archilestris – 119
capnopterus Martin & Papavero, 1970, Archilestris, error – 119
capronae Maryin, 1961, Lissoteles – 133
captans (Walker), 1851 (Trupanea), Promachus – 45
caputitis Scarbrough & Perez-Gelabert, 1995, Wilcoxius – 47
?*caracasae* (Martin), 1975 (Pachychaeta), Lecania – 31
carbonaria (Wiedemann), 1828 (Asilus), Threnia – 14
carbonarius (Philippi), 1865 (Dasypogon), Saropogon – 59
Carebaricus Artigas & Papavero, 1991 – 112
carinata (Bellardi), 1861 (Erax), Carinefferia – 16
carinatus-group of Hine, 1919, Erax – 16
carinatus Wilcox, 1964, Metapogon – 116
Carinefferia Artigas & Papavero, 1997 – 16
carmina Kaletta, 1974, Eumecosoma – 93
carrascoi (Artigas), 1974 (Scylaticus), Scylaticodes – 130
carrerai Artigas & Papavero, 1988, Neodiogmites – 74
carrerai Ayala, 1982, Glaphyropyga – 28
carrerai (Bromley), 1951 (Doryclus), Pseudorus – 78
carrerai (Hull), 1958 (Catostola), Ctenodontina – 30
carrerai Hull, 1958, Lycosimyia – 95
Carreraia Cole & Pritchard, 1964 – 35
Carreraomyia Cole, 1969 – 35
castaneipes Bigot, 1878, Lamprozona – 95
castaneus (Macquart), 1838 (Dasypogon), Diogmites – 69
castanipes Bigot, 1879, Lamprozona, emend. – 95
castigans (Walker), 1851 (Dasypogon), Caenarolia – 68
catastygnum Papavero, 1971, Microsylum – 121
Catonomya Hull, 1962 – 92
Catostola Hull, 1958 – 30
catulus Osten Sacken, 1887, Stichopogon – 134
caudatus Hine, 1941, Proctacanthus – 54
caudatus Williston, 1901, Psilocurus – 105
caudex (Walker), 1849 (Asilus), Nerax – 19

- cayennensis* (Gmelin), 1792 (*Asilus*), *Blepharepium*, error – 66
caymanensis Scarbrough, 1988, *Efferia* – 18
cazieri (Curran), 1953 (*Erax*), *Nerax* – 19
?cellata (Schiner), 1868 (*Erax*), *Diplosynapsis* – 17
Centrolaphria Enderlein, 1914 – 92
ceppicus Say, 1829 (*Dasypogon*) – 139
cerasinus Martin, 1975, *Machimus* – 9
Ceratotaenia Lynch Arribálzaga, 1880, emend. – 90
Ceraturgus Wiedemann of authors, misident. – 61
cerdai Tomasovic, 2022, *Porasilus* – 23
Cerotainia Schiner, 1866 – 90
Cerotoninops Curran, 1930 – 82
Cerozodus Bigot, 1857 – 30
cerverai Bromley, 1929, *Atomosia* – 86
chalycoprocta (Loew), 1866 (*Sphageus*), *Prolepsis* – 122
chalybea Hermann, 1912, *Pronomopsis* – 77
chalybeiventris (Loew), 1851 (*Dasypogon* (*Saropogon*)),
 Saropogon – 60
chalybeum Williston, 1885, *Andrenosoma* – 81
chapadensis (Bromley), 1928 (*erax*), *Nerax* – 19
cheysomela Cresson Jr., 1926, *Mallophora*, error – 38
chilechicoensis (Artigas), 1970 (*Philonerax*), *Lochmorhynchus* – 33
chilensis (Brèthes), 1925 (*Atomosia*), *Lamprozona* – 95
chilensis (Hermann), 1912 (*Automolina*), *Lamprozona* – 95
chilensis Macquart, 1850 (*Dasypogon*) – 139
chilensis (Macquart), 1850 (*Dasypogon*), *Scylaticodes* – 131
Chilesus Bromley, 1932 – 7
chiliensis (Macquart), 1850 (*Erax*), *Cratolestes* – 17
chiloensis (artigas), 1970 (*Nesiotes*), *Neotes* – 29
choapensis (artigas), 1970 (*Araiopogon*), *Saropogon* – 59
chrysauge (Osten Sacken), 1887 (*Asilus*), *Wyliea* – 14, 1907 – 113
Chrysoceria Williston, 1907 – 113
chrysomela Bromley, 1925, *Mallophora* – 38
chrysops Martin, 1957, *Beameromyia* – 107
chrysorhea Hull, 1958, *Helolaphyctis* – 113
Chrysotrichlis Artigas, Papavero & Costa, 1997 – 102
cienagaensis (Hengst & Artigas), 2003 (*Philonerax*),
 Lochmorhynchus – 32
ciguaya Scarbrough & Perez-Gelabert, 2006, *Atomosia* – 86
ciliata Walker, 1851, *Mallophora* – 41
cilicrura Rondani, 1850, *Mallophora* – 37
cilipes Walker, 1857, *Lampria* – 98
cinctum (Bellardi), 1851 (*Laphria*), *Andrenosoma* – 81
cinctus Bellardi, 1861, *Promachus* – 45
cinerascens (Bellardi), 1861 (*Erax*), *Albibarbeffera* – 15
cinerascens Walker, 1855, *Mallophora* – 40
cinerescens Papavero & Ibáñez Bernal, 2001 (*Erax*),
 Albibarbeffera, error – 15
cinereum (Bellardi), 1861 (*Laphria*), *Andrenosoma* – 81
cinereum Bigot, 1878 (*Bathypogon*) – 139
cinereus (Philippi), 1865 (*Erax*), *Lochmorhynchus* – 33
cingulata Artigas & Angulo, 1980, *Mallophora* – 38
cingulata (Bellardi), 1861 (*Erax*), *Albibarbeffera* – 15
cingulipes Wakker, 1849, *Leptogaster* – 108
cinnamomeus Scarbrough, 1984, *Ommatius* – 48
cinthiae Vieira, Castro & Bravo, 2004, *Ommatius* – 48
circumdata Bellardi, 1861, *Lampria* – 98
circumflava Artigas & Angulo, 1980, *Mallophora* – 38
claripennis (Schiner), 1868 (*Bathypogon*), *Creolestes* – 120
claripennis (Schiner), 1967 (*Lochites*), *Senobasis* – 79
clarkei Hull, 1958, *Zabrotica* – 132
clasicella (Carrera), 1960 (*Atractia*), *Hybozelodes* – 94
clasicella (Macquart), 1850 (*Laphria*), *Pilica* – 83
clava Scarbrough & Perez-Gelabert, 2009, *Efferia* – 19
clavata (Macquart), 18387 (*Asilus*), *Lecania* – 31
Clavator Philippi, 1865 – 131
clavigera (Rondani), 1850 (*Astylium*), *Senobasis* – 79
clavigera Rondani of Carrera, 1952, *Senobasis*, misident. – 79
clavigeroides Papavero, 1975, *Senobasis* – 79
clavijoi Kalletta, 1986, *Cerotainia* – 90
clavipes Curran, 1941, *Mallophora* – 38
clavipes (Fabricius), 1805 (*Laphria*), *Lampria* – 98
clavipes (Johnson), 1894 (*Leptogaster*), *Apachekolos* – 106
clavistylus (Rondani), 1848 (*Gonytes*), *Leptogaster* – 108
clavitarsis Curran, 1941, *Mallophora* – 38
Cleptomyia Carrera, 1949 – 58
Cnodalomyia Hull, 1962 – 7
coachellus Wilcox, 1966, *Ablautus* – 113
coarcatum (Perty), 1833 (*Laphria*), *Blepharepium* – 66
coarcatum Perty of Snow, 1896; Aldrich, 1905; Martin, 1956,
 Blepharepium, misident. – 67
Cochleariocera Artigas, Papavero & Costa, 1997 – 103
cockerellorum (James), 1953 (*Erax*), *Nerax* – 19
coerulea Williston, 1901, *Laphria* – 100
coeruleiventris Thomson, 1869, *Mallophora* – 36
coffeatus (Wiedemann), 1819 (*Dasypogon*), *Diogmites* – 70
colalao Lamas M., 1973, *Prolepsis* – 122
coleus Martins, 1975, *Machimus* – 9
colimae Martin, 1968, *Stenopogon* – 129
colombiae Macquart, 1838 (*Asilus*) – 139
colombiae Martin, 1971, *Leptopteromyia* – 110
columbiae Williston, 1891 (*Asilus*), error – 139
columbiana (Enderlein), 1914 (*Centrolaphria*), *Dissmeryngodes* – 92
columbina Schiner, 1868 (*Laphystia*) – 106
comata (Bellardi), 1861 (*Erax*), *Carineffera* – 16
comatus (Hermann), 1912 (*Atractia*), *Hybozelodes* – 94
combustus Loew, 1874, *Saropogon* – 59
combustus Loew of Back, 1909, *Saropogon*, misident. – 59
commiles (Walker), 1851 (*Erax*), *Nerax* – 20
complanatus Scarbrough, 1993, *Ommatius* – 48
complicata (James), 1953 (*Pachychaeta*), *Lecania* – 31
componens Walker, 1860, *Laphria* – 99
composita (Hine), 1918 (*Asilus*), *Polacantha* – 11
concepcionensis (Bromley), 1932 (*Asilus*), *Menexenus* – 10
concinnata (Williston), 1901 (*Erax*), *Carineffera* – 16
concinnata Williston, 1901, *Leptogaster* – 108
concolor Walker, 1851 (*Erax*), *Triorla* – 24
conjugens (Hermann), 1912 (*Atractia*), *Hybozelodes* – 94
connexus (Wiedemann), 1828 (*Asilus*), *Eichoichemus* – 26
consanguineus (Loew), 1866 (*Stenopogon*), *Ospricerus* – 127
conspectus Tomasovic, 2001, *Pseudorus* – 77
constrictus Scarbrough, 2002, *Ommatius* – 48
contraria Walker, 1851, *Mallophora* – 43
contusa (Wiedemann), 1828 (*Laphria*), *Pilica* – 83
conus Scarbrough, 2002, *Ommatius* – 48
copelloi Gemignani, 1930, *Mallophora* – 41
Cophura Osten Sacken, 1887 – 63
cophuroides Carrera, 1949, *Aspidopyga* – 63
coprates Walker, 1848, *Proctacanthus* – 54
copulata (Wiedemann), 1819 (*Asilus*), *Lecania* – 31
coquilletti Back, 1909, *Saropogon* – 59
coquilletti (Hine), 1919 (*Erax*), *Pogonioeffera* – 22
coquilletti Hine, 1911, *Proctacanthus* – 54
coquilletti Wilcox, 1935, *Ablautus* – 113
coquimbensis Artigas, 1970, *Dasycurton* – 113
cora Curran, 1941, *Mallophora* – 37
cora Pritchard, 1943, *Cophura* – 63
corallium Martin, 1966, *Andrenosoma* – 81
corallogaster (Bigot), 1878 (*Laphria*), *Lampria* – 98
?coriacea (Wiedemann), 1821 (*Dasypogon*), *Holococephala* – 135
coronata Schiner, 1867, *Atractia* – 89
corsair Bromley, 1951, *Senobasis* – 79
cordesi Artigas & Angulo, 1980, *Mallophora* – 38
coscaronorum Artigas, Papavero & Costa, 1997, *Gymnotrichis* – 103

- costai Artigas, Papavero & Serra, 1991, *Cyphomyiactia* – 92
 costalis Lynch Arribálzaga, 1880 (*Dasytigon*) – 139
 costaricensis Lamas M., 1973, *Prolepsis* – 122
 costatus Rondani, 1850, *Ommatius* – 48
 costatus (Schiner), 1868 (*Erax*), *Eicherax* – 25
coxalis Curran, 1930, *Atomosia* – 86
crabroniformis (Schiner), 1867 (*Cacodaemon*), *Prolepsis* – 122
crassitarsis (Macquart), 1838 (*Dasytigon*), *Lastaurus* – 73
crassitarsis (Macquart), 1846 (*Megapoda*), *Pseudorus* – 78
crassitarsus (Bromley), 1946 (*Dasytigon*), *Lastaurus*, error – 73
crassum Bromley, 1929, *Andrenosoma* – 81
crassus (Bromley), 1932 (*Asilus*), *Lochyrrus* – 8
Cratolestes Hull, 1962 – 16
Cratopoda Hull, 1962 – 8
craverii (Bellardi), 1861 (*Dasytigon*), *Diogmites* – 70
craverii Bellardi, 1861, *Mallophora* – 40
craverii Bellardi, 1861, *Proctacanthus* – 54
crenus Martin, 1975, *Wilcoxius* – 47
Creolestes Hull, 1962 – 119
cressoni (Hine), 1919 (*Erax*), *Carinefferia* – 16
cribatus (Hengst & Artigas), 2003 (*Philonerax*), *Lochmotryphon* – 33
cribratus (Hull), 1962 (*Lonchodogonus*), *Lochmorhynchus* – 33
crinita Martin, 1957, *Apachekolos* – 106
critesi (Artigas), 1970 (*Hypenetes*), *Tillobroma* – 131
crocea Williston, 1901, *Leptogaster* – 108
croceiventris (Wiedemann), 1821 (*Laphria*), *Dasyllis* – 82
crocuscopa Artigas & Angulo, 1980, *Mallophora* – 38
cruciger Hermann, 1921 (*Scylaticus*) – 132
cruentum (McAtee), 1919 (), *Andrenosoma* – 81
cruentus (Lynch Arribálzaga), 1880 (*Proctacanthus*), *Taurhynchus* – 55
cruralis Rondani, 1863, *Mallophora* – 40
Cryptomerynx Enderlein, 1914 – 96
Ctenodontina Enderlein, 1914 – 30
ctesicles (Walker), 1851 (*Asilus*), *Lecania* – 31
cuantlensis (Bellardi), 1861 (*Dasytigon*), *Diogmites* – 70
cubana Artigas & Papavero, 1997, *Mallophora* – 38
cubanus Scarbrough, 1985, *Ommatius* – 48
cubensis (Bigot), 1857 (*Gonypes*), *Beameromyia* – 107
cubensis (Bromley), 1929 (*Erax*), *Nerax* – 20
culiciformis (Walker), 1855 (*Asilus*), *Eicherax* – 25
cunctabundum Papavero & Bernardi, 1973, *Blepharepium* – 67
cuneigaster (Artigas), 1970 (*Scylaticus*), *Scylaticodes* – 131
curicoensis Artigas, 1970, *Pritchardia* – 122
curiosa Curran, 1941, *Mallophora* – 37
currani Bromley, 1931, *Andrenosoma* – 81
currani (Bromley), 1951 (*Erax*), *Triorla* – 25
currani Martin, 1959, *Holopogon* – 115
curupaoensis (Kaletta), 1978 (*Voluptarius*), *Oidardis* – 95
curvicosta Carrera, 1958, *Holcocephala* – 136
cuyanus Lynch Arribálzaga, 1880 (*Asilus*) – 139
cyanea Macquart, 1834, *Megapoda* – 77
cyaneiventris (Macquart), 1846 (*Megapoda*), *Pseudorus* – 78
cyanescens (Bigot), 1878 (*Cormansis*), *Lamprozona* – 95
cyanescens Rondani, 1848, *Atomosia* – 86
cyaniventris (Bromley), 1934 (*Andrenosoma*), *Pilica* – 83
cyanogaster (Loew), 1851 (*Dasytigon* (*Saropogon*)), *Araucopogon* – 58
cyanogaster (Loew), 1851 (*Dasytigon* (*Saropogon*)), *Saropogon* – 60
Cylicomera Lynch Arribálzaga, 1881 – 120
Cylindrophora Philippi, 1865 – 65
cymafera (Artigas), 1983 (*Apoxytria*), *Cymbipyga* – 103
Cymbipyga Artigas, Papavero & Costa, 1997 – 103
Cyphomyiactia Arrtigas, Papavero & Serra, 1991 – 92
Cyphotomyia Williston, 1889 – 90
cyrtophora (Hermann), 1912 (*Nusa*), *Pilica* – 83
Cyrtophriina Papavero, 1975 – 75
Cyrtophrys Loew, 1851 – 75
Cyrtopogon Loew, 1847 – 114
Cyrtopogonini Artigas & Papavero, 1991 – 112
Cystoprosopa Hull, 1962 – 125
dallasi Gemignani, 1931, *Mallophora* – 42
Damalini Hull, 1962 – 135
dammersi Wilcox, 1965, *Cophura* – 63
dana Curran, 1941, *Mallophora* – 37
dandrettae Carrera, 1949, *Pseudorus* – 77
dandrettae Carrera of Oldroyd, 1964, *Pseudorus*, misident. – 78
danforthi Curran, 1935, *Atomosia* – 86
danforthi Curran, 1951, *Proctacanthus* – 54
daphne Pritchard, 1943, *Cophura* – 63
Dapsilochetus Hull, 1962 – 125
daraps (Walker), 1849 (*Asilus*), *Taurhynchus* – 55
darlingtoni Curran, 1951, *Proctacanthus* – 54
dascyllus (Walker), 1849 (*Asilus*), *pogoniefferia* – 23
Dasipogonia Raguès, 1908 – 5
Dasipogonina Rondani, 1863 – 57
Dasipogoninae Karl, 1835 – 57
Dasycyrton Philippi, 1865 – 113
Dasyllis Loew, 1851 – 82
Dasypecus Philippi, 1865 – 114
Dasypoginæ Wulp, 1896 – 57
Dasypogomnia Raguès, 1908 – 57
Dasypogonidae Bigot, 1852 – 57
Dasypogonidii Bigot, 1853 – 57
Dasypogonina Loew, 1857 – 57
Dasypogoninae Macquart, 1838 – 57
Dasypogonini Macquart, 1838 – 57
Dasypogonitae Macquart, 1838 – 57
Dasypogonites Macquart, 1850 – 57
dasypus (Wiedemann), 1828 (*Laphria*), *Atomosia* – 86
Dasythricaria Lynch Arribálzaga, 1882 – 96
dasythrix Hermann, 1912, *Cerotainia* – 90
Dasythrix Loew, 1851 – 96
Dasytrichini Artigas, Papavero & Pimentel, 1988 – 96
Dasytrichini Lynch Arribálzaga, 1882 – 97
Daulopogon Loew, 1874 – 133
davidsoni (Artigas), 1970 (*Hypenetes*), *Tillobroma* – 131
davisi Wilcox, 1966, *Efferia* – 18
debilis Hermann, 1912, *Cerotainia* – 90
dejectus (Williston), 1901 (*Holopogon*), *Heteropogon* – 114
delgadoi Kaletta, 1978, *Bathropsis* – 90
deltoidea (Bellardi), 1861 (*Discocephala*), *Holcocephala* – 136
demifasciata (Macquart), 1850 (*Erax*), *Triorla* – 24
dentatus Scarbrough, 2002, *Ommatius* – 48
dentipes Fabricius, 1805 (*Laphria*) – 101
depressus Scarbrough, 2002, *Ommatius* – 48
Deromyia Philippi, 1865 – 76
Deromyia Philippi of authors, misident. – 69
deserti (Wilcox), 1966 (*Efferia*), *Pogonioefferia* – 22
destitutus Scarbrough, 2002, *Ommatius* – 49
dichroa Wiedemann, 1828 (*Laphria*) – 101
dichroma Kertész, 1909, *Eumecosoma*, emend. – 93
dicrana Scarbrough, 1997, *Beameromyia* – 107
Dicranus Loew, 1851 – 120
dicromum Bigot, 1878, *Eumecosoma* – 93
didymus Scarbrough, 1993, *Ommatius* – 49
digitata (Artigas), 1970 (*Hypenetes*), *Tillobroma* – 131
digna Pritchard, 1941, *Aphamartania* – 62
dignus Scarbrough, 2000, *Ommatius* – 49
dilata Martin, 1966, *Townsendia* – 134
dilectus (Walker), 1855 (*Erax*), *Triorla* – 24

- dimidiata* (Curran), 1935 (Allopogon), Caenarolia – 67
dimidiata Hermann, 1926, Holcocephala – 136
dimidiatus (Macquart), 1847 (Dasypogon), Taracticus – 61
dimidiatus Oldroyd, 1964, Pseudorus – 77
dimidiatus Scarbrough, 1985, Ommatius – 49
dina (Curran), 1934 (Proctacanthus), Taurhynchus – 55
Diochiini Arnett, 1985 – 117
Dioctriinae Enderlein, 1936 – 117
Dioctriini Hendel, 1936 – 117
Dioctrini Lehr, 1977 – 117
Diocrodes Coquillett, 1904 – 61
Diogmites Loew, 1866 – 68
diogmitiformis Artigas & Papavero, 1971. Taperigna – 124
Diplosynapsis Enderlein, 1914 – 17
Discocephala Macquart, 1838 – 135
disjunctus (Williston), 1901 (Erax), Eicherax – 25
dispar (Bromley), 1937 (Stenopogon), Scleropogon – 128
dispar Coquillett, 1902, Saropogon – 59
dispar (Hermann), 1912 (Atractia), Hybozelodes – 94
dispar (Walker), 1855 (Atomosia), Dissmeryngodes – 92
Dissmeryngodes Hermann, 1912 – 92
dissona Lamas M., 1973, Cylicomera – 120
distendens (Wiedemann), 1838 (Asilus), Pseudorus – 77
distends Notario, Michela, Fiorentino & Castresana, 2000,
 Pseudorus, error – 78
distincta Schiner, 1867, Leptogaster – 108
distinguendus (F. Lynch Arribálzaga), 1881 (Scylaticus),
 Euthrixius – 130
diversus Williston, 1901, Ospriocerus – 127
dives (Wiedemann), 1828 (Laphria), Lampria – 98
divisa (Walker), 1860 (Discocephala), Holcocephala – 136
divisus (Coquillett), 1902 (Pyncopogon), Heteropogon – 114
Dizonias Loew, 1861 – 122
doellojuradoi Gemignani, 1930, Mallophora – 36
dolabridiformis Scarbrough, 2002, Ommatius – 49
Dolichodes Macquart, 1838 – 125
dolichogaster (Williston), 1901 (Erax), Nerax – 21
dolichomerus (Williston), 1901 (Asilus (Heligmoneura)),
 Prolatiforceps – 12
dolosus Hull, 1962, Graptostylus – 114
dominicanus Curran, 1951, Proctacanthus – 54
dorothyae (Martin), 1975 (Furcilla), Scarbroughia – 46
dorothyae Martin, 1962, Heteropogon – 114
dorothyae Martin, 1968, Stenopogon – 129
dorsalis Macquart, 1848 (Dasypogon) – 139
dorsalis (Williston), 1901 (Leptogaster), Beameromyia – 107
Doryclus Jaennicke, 1867 – 77
drabicola Cole, 1916, Lasiopogon – 133
drakei (Prithard, 1935 (Buckellia), Cophura – 63
dryas Fisher & Hespenheide, 1982, Glaphyropyga – 28
dubia Bigot, 1878, Cerotainia – 91
dubius (Bellardi), 1861 (Dasypogon), Diogmites – 70
duillius (Walker), 1849 (Dasypogon), Diogmites – 70
duncani (Bromley), 1937 (Stenopogon), Scleropogon – 128
duncani (Wilcox), 1966 (Efferia), Albibarbeffera – 15
duncani Wilcox, 1941, Heteropogon – 114
durangoensis (Cole), 1964 (Mallophorina), Megaphorus – 44
dureti Artigas & Angulo, 1980, Mallophora – 38

eburneus Hull, 1962 (Heteropogon), error – 139
eburnus Walker, 1849 (Dasypogon) – 139
Eccritosia Schiner, 1866 – 53
Echinotropis Martin, 1975 – 11
Echthodopini Adisoemarto & Wood, 1975 – 118
Ecoritosia Bromley, 1951, error – 53
Efferia Coquillett of authors, misident – 19
Efferia Coquillett, 1893 – 17

Efferia, aestuans-group of Wilcox, 1966 – 19
Efferia, albobarbis-group of Wilcox, 1966 – 15
Efferia, anomala-group of Wilcox, 1966 – 17
Efferia, arida-group of Wilcox, 1966 – 16
Efferia-group – 14
ehrenbergi (Wilcox), 1966 (Efferia), Pogonioefferia – 22
Eicherax Bigot, 1857 – 25
Eicherax-group – 25
Eichoichemus Bigot, 1857 – 26
Eichoichemus-group – 26
elegans (Bromley), 1934 (Andrenosoma), Pilica – 84
elegans Hermann, 1912, Strombocodia – 96
elizabethae Martin, 1968, Stenopogon – 129
elotensis (Martin), 1966 (Sphageus), Prolepsis – 123
elusivus Scarbrough, 2003, Ommatius – 49
emarginata Artigas, 1970, Cratopoda – 8
emarginatus Scarbrough, 1985, Ommatius – 49
emeralda Hull, 1962, Sintoria – 117
emiliae Carrera, 1960, Mallophora – 38
Emphysomera Schiner of authors, misident. – 47
Enarmostus Walker, 1855, error – 113
Enigmomorpheus Hull, 1962, error – 121
Enigmomorphini Hull, 1962 – 118
Enigmomorphus Hermann, 1912 – 121
Epipamponeurus Becker, 1919 – 8
equestris (Wiedemann), 1828 (Dasypogon), Caenarolia – 68
Erax Scopoli of authors, misident. – 17, 19
erax (Bromley), 1934 (Andrenosoma), Pilica – 83
Erax, aestuans-group f Hine, 1919 – 19
Erax, animalus-group of Hine, 1919 – 17
Erax, barbatus-group of Hine, 1919 – 15
Eraxasilus Carrera, 1959 – 26
eraxoides (Curran), 1935 (Promachbina), Eichoichemus – 26
Erecmoneminae Hermann, 1912, part – 80, 84, 96, 97, 100
Eremomyia Artigas, 1970 – 32
Eretomyia Artigas, 1971 – 32
Eristicus Loew, 1848 – 25
eritrichus (Philippi), 1865 (Asilus), Stizolestes – 13
erythrogaster (Wiedemann), 1828 (Laphria), Pilica – 83
erythropus Schiner, 1867, Ommatius – 49
erythropyga (Wiedemann), 1828 (Laphria), Andrenosoma – 81
erythrura Hermann, 1912, Dasyllis – 82
Euarmostus Walker, 1851 – 113
Eumecosoma Schiner, 1866 – 93
eupator (Walker), 1851 (Asilus), Eicherax – 25
eupoda (Bigot), 1878 (Cormansis), Atomosia – 87
Eurhabdus Aldrich, 1923 – 107
eurylabis (Wiedemann), 1828 (Asilus), Nerax – 20
Euthrix Philippi, 1865 – 130
Euthrixius Artigas, 1971 – 130
Eutrichodes Hermann, 1920 – 103
evae Artigas, Papavero & Costa, 1997, Protometer – 105
evansi Martin, 1968, Ospriocerus – 127
exacta Scarbrough & Perez-Gelabert, 2009, Eifferia – 18
examinans (Walker), 1851 (Dasypogon), Diogmites – 69
excellens Enderlein, 1914, Archilestris – 119
exilis Curran, 1928, Ommatius – 49
eximea (Hine), 1919 (Erax), Albibarbeffera, error – 15
eximia (Bellardi), 1861 (Erax), Albibarbeffera – 15
exquisita (Osten Sacken), 1887 (Proctacanthus),
 Proctacanthella – 54

facialis (Curran), 1935 (Mirolestes), Cyrtophrys – 76
fairchildi Curran, 1942, Mallophora – 40
fallax (Macquart), 1846 (Dasypogon), Lastaurus – 74
falcatus Scarbrough, 2002, Ommatius – 49
farri Hull, 1962, Phellopteron – 100

- fascia Martin, 1966, Itolia – 115
fascialis (Carrera), 1949 (Mirolestes), Cyrtophrys – 76
fasciata Lynch Arribálzaga, 1880, Asicya – 102
fasciata Walker, 1851, Mallophora – 43
fasciatus (Macquart), 1834 (Dasypogon), Diogmites – 70
fascipennis (Hermann), 1912 (Rhopalaster), Ichneumolaphria – 98
fascipennis (Macquart), 1834 (Laphria), Dasyllis – 82
fascipennis Macquart, 1850, Mallophora – 43
fascipennis (Schiner), 1867 (Euscelidia), Systologaster – 111
fascipennis Philippi, 1865 and Williston, 1891, Deromyia, error – 76
fasciventris Hermann, 1926, Holcocephala – 137
fasciventris (Macquart), 1950 (Dasypogon), Stichopogon – 134
fautricoides Curran, 1930, Mallophora – 38
fautrix Osten Sacken, 1887, Mallophora – 38
fax (Lynch Arribálzaga), 1881 (Tolmerolestes), Prolepsis – 123
feigei Kalletta, 1983, Cophura – 63
felis Bromley, 1931, Stenopogon – 129
feminea Curramn, 1930, Cerotainia – 91
?femorata Hermann, 1926, Holcocephala – 136
femorata Macquart, 1838, Lecania – 31
femorata Roeder, 1900, Tricloscelis – 105
fenestellus Martin, 1975, Prolatiforceps – 12
fenestrata (Macquart), 1838 (Senobasis), Prolepsis – 123
fenestratus Hermann, 1921, Acronyches – 112
fenestratus Bigot, 1878, Lastaurus – 73
fernandezi Ayala, 1982, Holcocephala – 136
fernandezi Scrbrough, 2002, Omnitius – 49
fernandezii Kalletta, 1976, Lissoteles – 133
ferruginea Walker, 1855, Leptogaster – 108
ferrugineum (Macquart), 1838 (Dolichodes), Plesioma – 125
ferrugineus (Lynch Arribálzaga), 1880 (Allopogon), Diogmites – 70
fervens Wiedemann, 1830, Leptogaster – 108
fervidus (Curran), 1934 (Proctacanthus), Taurhynchus – 55
fiebrigii Bezzi, 1909, Townsendia – 134
fimbriata Hermann, 1926, Holcocephala – 136
 \dagger *fimbriatus* Scarbrough & Poinar, 1992, Ommatius – 49
 \dagger *fimbrillus* Scarbrough, 2000, Ommatius – 49
fisheri Martin, 1967, Holopogon – 115
fitzroyi Artigas, 1974, Zoticus – 30
flamipennis (Bromley), 1931 (Andrenosoma), Pilica – 84
flaventis Scarbrough & Perez-Gelabert, 2005, Apachekolos – 107
flavescens James, 1853 (Eicherax) – 56
flavescens Scarbrough, 2003, Ommatius – 49
flavianalis (Macquart), 1848 (Erax), Eichoichemus – 27
flavicollis Curran, 1941, Mallophora, nom. nud. – 36
flavida (Wiedemann), 1828 (Asilus), Lecania – 31
flavidus (Macquart), 1838 (Erax), Lecania – 31
flavifasciatus (Macquart), 1838 (Trupanea), Amblyonychus – 34
flavipellitus Enderlein, 1914, Lastaurus – 73
flavipennis Macquart, 1846 (Dasypogon) – 139
flavipennis (Macquart), 1846 (Dasypogon), Aphamartania – 62
flavipennis (Macquart), 1846 Proctacanthus), Taurhynchus – 55
flavipennis Wiedemann, 1828 (Dasypogon) – 139
flavipes (Bigot), 1880 (Phoneus), Efferia – 18
flavipes Coquillett, 1904, Ablautus – 113
flavipes Hermann, 1912, Cerotainia – 91
flavipes Hermann, 1926, Holcocephala – 136
flavipes Hull, 1962, Laphygmolestes – 104
flavitibia (Aldrich), 1905 (Neophoneus), Efferia, error – 18
flavitibius (Bigot), 1880 (Phoneus), Efferia, emend. – 18
flaviventris Engel, 1930, Holcocephala, error – 137
flavofasciatus (Wiedemann), 1828 (Asilus), Nerax – 20
flavotibia (Bigot), 1878 (Phoneus), Efferia, emend. – 18
flavotibialis Martin, 1968, Stenopogon – 129
flavotibius (Bigot), 1880 (Phoneus), Efferia, emend. – 18
flexus Scarbrough, 2000, Ommatius – 49
flukei Carrera, 1952, Senobasis – 79
fluviatilis (Carrera), 1960 (Atractia), Lycosimyia – 95
forbesi (Curran), 1931 (Erax), Nerax – 20
forfex Osten Sacken, 1887, Promachus – 45
formidolosa (Walker), 1860 (Laphria), Pilica – 83
fortis (Walker), 1855 (Erax), Efferia – 18
fragilis Back, 1909, Stichopogon – 134
fragilis (Carrera), 1944 (Shannomyioleptus), Schildia – 110
fraterna Lynch Arribálzaga, 1881, Cylicomera – 120
fraternus Bigot, 1878, Saropogon – 60
fraternus Macquart, 1846 (Dasypogon) – 139
frauendorfi Schiner, 1866, Aphamartania – 62
fredericoi Carrera, 1952, Atomasia – 86
freycineti Macquart, 1838, Mallophora – 36
frezieri Artigas, 1970, Lochyrus – 9
fritzi Artigas & Angulo, 1980, Mallophora – 39
frontalis Curran, 1930, Atomasia – 86
frosti Bromley, 1951, Senobasis – 79
frustus flavidus (Cole), 1964 (Mallophorina), Megaphorus – 44
fucosa (Artigas), 1970 (Hypenetes), Tillobroma – 131
fucosoides Artigas, Lewis & Parra, 2005, Tillobroma – 131
fulgida Schiner, 1868, Lampria – 99
fuliginosum (Wiedemann), 1821 (Dasypogon), Plesioma – 125
fuliginosus (Bellardi), 1861 (Asilus, ?Mochtherus), Philonicus – 11
fulva Banks, 1911, Mallophora – 41
fulvianalis Macquart, 1850, Mallophora – 39
fulvibarbs (Macquart), 1848 (Erax), Efferia – 18
fulvicornis (Macquart), 1850 (Dasypogon), Saropogon – 60
fulvicornis (Macquart), 1846 (Dasypogon), Tillobroma – 131
fulvigaster Bigot, 1878, Microstylum – 121
fulvipes Carrera, 1946, Atoniomya – 89
fulvipes (Hermann), 1912 (Atractia), Hybozelodes – 94
fulvipes (Philippi), 1865 (Deromyia), Neoderomyia – 59
fulvipes (Philippi), 1965 (Deromyia), Saropogon – 60
fulvis (Bromley), 1946 (Senobasis), Diogmites, error – 70
fulvithorax Fabricius, 1805, Laphria – 100
fulvithorax (Macquart), 1838 (Erax), Eicherax – 25
fulviventris Macquart, 1850, Mallophora – 39
fulvus (Bigot), 1878 (Dasypogon), Diogmites – 70
fumiflamma Walker, 1851, Prolepsis – 123
funebris Artigas, Papavero & Pimentel, 1988, Pilica – 83
funebris Lamas M., 1973, Prolepsis – 123
funestum Loew, 1861, Plesioma – 126
furax (Williston), 1885 (Erax), Albabarbefferia – 15
Furcilla Martin, 1975 – 46
fusca Bromley, 1934, Mallophora – 41
fuscanipennis (Macquart), 1850 (Erax), Nerax – 20
fuscipennis Bellardi, 1862, Ommatius – 49
fuscipennis (Blanchard), 1852 (Gonypus), Deromyia – 76
fuscipennis (Macquart), 1834 (Dasypogon), Blepharepium – 67
fuscipennis (Macquart), 1846 (Trupanea), Promachus – 45
fuscipennis (Macquart), 1847 (Erax), Eraxasilus – 26
fuscipennis Macquart of Bellardi, 1861, Promachus, misident. – 44
fuscolimbatus Bigot, 1878, Stenopogon – 129
fucus (Wiedemann), 1828 (Erax), Nerax – 20

gabrieli Cole & Wilcox, 1938, Lasiopogon – 133
galactodes Loew, 1866, Microstylum – 121
galbinus Martin, 1968, Stenopogon – 129
gamaxus Walker, 1851 (Asilus) – 139
gammaxus Hull, 1962 Asilus, error – 139
garciai Lamas M., 1971, Porasilus – 23
gavius (Walker), 1851 (Asilus), Eichoichemus – 27
gayi (Macquart), 1838 (Dasypogon), Saropogon – 60
gayi (Macquart), 1838 (Asilus), Stizolestes – 13
gayi Macquart of Hull, 1962, Cratopoda, misident. – 8
geijskesi Curran, 1942, Mallophora – 40
geijskesi Papavero & Bernardi, 1974, Archilestris – 119

- gelascens* (Walker), 1860 (Dasypogon), Stichopogon – 134
gellergrimmi Papavero, Cophura, **nom. nov.** – 63
geminatus Bromley, 1932, Chilesus – 6
geminus Scarbrough & Perez-Gelabert, 2006, Ommatius – 49
geniculata Macquart, 1838, Mallophora – 43
geniculata (Wiedemann), 1821 (Laphria), Atomosia – 86
geniculatus (Bigot), 1878 (Ceraturgus), Taracticus – 61
genitalis (Bromley), 1934 (Pachychaeta), Lecania – 31
geosarginus Papavero, 1971, Acronyches – 112
gerion (Walker), 1849 (Asilus), Eraxasilus – 26
germainii Bigot, 1857, Lycomya – 9
gertschi (Bromley), 1951 (Asilus), Machimus – 9
gerulewiczii Kalleta, 1986, Cerotainia – 91
getzendaneri Wilcox, 1959, Cophura – 63
gibba (Curran), 1930 (Eumecosoma), Oidardis – 95
gibber (Williston), 1883 (Cyrtopogon), Metapogon – 116
gibbosa Hermann, 1912, Oidardis – 95
gibbosus Philippi, 1865, Dasycyrton – 113
giganteus Hine, 1911, Promachus – 45
giganteus (Hull), 1962 (Megonyx), Acronyches – 112
gladiatus Scarbrough, 2002, Ommatius – 49
Glaphyropyga-group – 28
Glaphyropyga Schiner, 1866 – 28
glauca Enderlein, 1914 (Laphria) – 101
goniostigma (Bellardi), 1861 (Dasypogon), Diogmites – 70
Gonytes Latreille, 1805 – 108
gorayebi Artigas, Papavero & Pimentel, 1988, Smeryngolaphria – 97
gossei (Farr), 1965 (Efferia), Nerax – 20
gracilis (Bigot), 1878 (Dasypogon), Caenarolia – 68
gracilis (Macquart), 1838 (Trupanea), Amblyonychus – 34
gracilis Martin, 1966, Townsendia – 134
gracilis Philippi, 1865, Deromyia – 76
gracilis Williston, 1907, Leptopteromyia – 110
gracillima (Walker), 1855 (Leptogaster), Schildia – 110
gracipes Artigas & Angulo, 1980, Mallophora – 39
Grajahua Artigas & Papavero, 1991 – 130
graminicola Farr, 1963, Beameromyia – 107
grandis (Hine), 1919 (Erax), Albibarbefferia – 15
grandis (Macquart), 1846 (Dasypogon), Neodiogmites – 74
Graptostylus Hull, 1962 – 114
griseus (Guérin-Méneville), 1830 (Asilus), Lochmorhynchus – 32
grossa Carrera, 1946, Atoniomyia – 89
guatemalae Martin, 1975, Schildia – 111
guentherii (Lynch Arribálzaga), 1882 (Doryclus), Pseudorus – 78
guerrerensis Pritchard, 1938, Taracticus – 61
guianicus (Curran), 1934 (Proctacanthus), Taurhynchus – 55
guildianus (Williston), 1885 (Mallophora), Megaphorus – 44
guimaraesi Artigas & Papavero, 1991, Archilestroides – 119
gurupi Artigas, Papavero & Pimentel, 1988, Smeryngolaphria – 97
guttatus Martin, 1975, Machimus – 9
gwenaee Scarbrough, 1984, Ommatius – 49
Gymnotriclis Artigas, Papavero & Costa, 1997 – 103
gyrophora (Schiner), 1868 (Lochites), Senobasis – 79
gyrophora Schiner of Carrera, 1949, Senobasis, misident. – 80
- Hadrokolos Martin, 1959 – 114
haemorhoa (Wiedemann), 1830 (Laphria), Dasyllis – 82
haemorrhoea (Zimsen), 1964 (Laphria), Plesiomma, error – 126
haemorrhoum (Fabricius), 1805 (Laphria), Plesiomma – 126
hagno (Walker), 1849 (Asilus), Taurhynchus – 55
haigi Wilcox, 1971, Scleropogon – 128
haitensis (Macquart), 1848 (Erax), Efferia – 18
haitensis Scarbrough, 1984, Ommatius – 49
haloesus (Walker), 1849 (Erax), Nerax – 20
halterata Enderlein, 1914, Diplosynapsis – 17
hamus Martin, 1968, Stenopogon – 129
hanebrinki Scarbrough & Rutkauskas, 1983, Ommatius – 49
- Haplopogon Engel, 1930 – 135
hebes (Walker), 1855 (Asilus), Eraxasilus – 26
Helcocephala Loew, 1874, error – 135
helix (Bromley), 1935 (Asilus), Cratopoda – 8
Helolaphytis Hermann, 1920 – 103
hemivitrea Artigas & Angulo, 1980, Mallophora – 39
herdonius Walker, 1851 (Asilus) – 139
hermanni Bezzi, 1910, Lissoteles – 133
hermanni Carrera, 1949, Pseudorus – 78
hermanni Hull, 1962, Myaptex – 46
heros Bromley, 1931, Andrenosoma – 81
hertiventris Artigas & Angulo, 1980, Mallophora, error – 42
heteroneurus Philippi, 1865, Dasypecus – 114
Heteropogon Loew, 1847 – 114
?heteroptera (Macquart), 1846 (Erax), Albibarbefferia – 15
Hexamerita Speiser, 1920 – 103
heydenii (Jaennicke), 1867 (Dasypogon), Diogmites – 70
?heydenii (Wiedemann), 1828 (Asilus), Amblyonychus – 34
hiemalis Martin, 1968, Stenopogon – 129
hierroi Scarbrough, 2003, Ommatius – 49
hilarii (Macquart), 1838 (Asilus), Lecania – 31
himantocera (Wiedemann), 1828 (Asilus), Glaphyropyga – 28
hirsutum Hermann, 1912, Eumecosoma – 93
hirtipes (Macquart), 1838 (Dasypogon), Pritchardia – 122
hirtiventris Macquart, 1850, Mallophora – 42
hirtosus (Bromley), 1946 (Dasypogon), Neodiogmites, error – 74
hirtuosus (Schiner), 1868 (Bathypogon), Creolestes – 120
hirtuosus (Wiedemann), 1821 (Dasypogon), Neodiogmites – 74
hispaniolae Scarbrough, 1984, Ommatius – 49
hispidella (Hermann), 1912 (Atonia), Atoniomyia – 89
hispidus Scarbrough, 1984, Ommatius – 49
histrio (Wiedemann), 1828 (Leptogaster), Psilonyx – 110
Hodites Hull, 1962 – 93
Hodophylax James, 1933 – 64
Holcocephala Jaennicke, 1867 – 135
holcocephalinus Papavero, 1975, Pseudorus – 78
Holocephala Williston, 1891, error – 135
Holopogon Loew, 1847 – 115
holosericeus Schiner, 1867, Ommatius – 49
homopoda (Bellardi), 1862 (Laphria), Lampria – 99
hondurana James, 1953, Atomosia – 86
Hoplistomerinae Enderlein, 1936 – 102
horni (Bromley), 1925 (Promachus), Amblyonychus – 34
huasquina Artigas, Lewis & Parra, 2005, Tillobroma – 131
huatajata Lamas M., 1973, Prolepsis, error – 123
huatayata Lamas M., 1972, Prolepsis – 123
hubbelli (James), 1953 (Erax), Nerax – 20
humatus Scarbrough, 1993, Ommatius – 50
humilis (Bellardi), 1861 (?Dasypogon), Cophura – 63
humilis (Bellardi), 1861 (Asilus), Machimus – 9
humilis Williston, 1901, Cophura – 64
hyacinthina Scarbrough, 1996, Leptogaster – 108
hyacinthinus Bigot, 1878, Saropogon – 60
hyalipennis (Macquart), 1838 (Erax), Proctophoroides – 27
Hybozelodes Hermann, 1912 – 94
Hypenetes Loew of authors, misident. – 131
Hypomelas (Loew), 1866 (Diogmites), Saropogon – 60
- ialapensis* James, 1953, Diogmites, error – 70
icarius Carrera, 1960, Alvarenga – 62
ichneumon (Osten Sacken), 1887 (Laphria), Lampria – 99
Ichneumolaphria Carrera, 1951 – 97
igneum Bromley, 1929, Andrenosoma – 81
igualae Pritchard, 1943, Cophura – 63
imbuda (Curran), 1934 (Erax), Nerax – 20
imisericorde (Artigas), 1970 (Megametopon), Megalometopon – 29
imitator Carrera, 1953, Diogmites – 70

- imitator Hermann, 1921, *Acronyches* – 112
 imitator Lynch Arribálzaga, 1883 (*Asilus*) – 139
impendens (Wiedemann), 1828 (*Asilus*), *Stenasilus* – 13
imperialis Forbes, 1988, *Efferia* – 18
inamatus Walker, 1860 (*Asilus*) – 139
inauratus Aldrich, 1905 (*Asilus*), error – 139
inca Curran, 1941, *Mallophora* – 39
inca (Martin), 1975 (*Pachychaeta*), *Lecania* – 31
inca Curran, 1942, *Blepharepium* – 66
incanipes Artigas & Angulo, 1980, *Mallophora* – 39
incerta (Becker), 1919 (*Metapogon*), *Cophura* – 64
incisura Scarbrough & Perez-Gelabert, 2009, *Efferia* – 18
incisuralis (Macquart), 1838 (*Trupanea*), *Amblyonychus* – 34
incisuralis Macquart, 1838, *Atomosia* – 87
incisuralis Scarbrough & Page, *in* Scarbrough & Page, 2005, Beamerymyia – 107
inclusus (Walker), 1851 (*Dasypogon*), *Diogmites* – 70
incognita (Forbes), 1987 (*Efferia*), *Albibarbefferia* – 15
incomptus (Philippi), 1865 (*Asilus*), *Stizolestes* – 13
incurvatus Scarbrough, 1993, *Ommatius* – 50
indecisa Lamas M., 1973, *Prolepsis* – 123
indecorum Loew, 1866, *Plesiomma* – 126
indigena Scarbrough & Perez-Gelabert, 2006, *Holcocephala* – 136
indistinctus (Bromley), 1937 (*Stenopogon*), *Scleropogon* – 128
indistinctus Bromley of Carrera & d'Andretta, 1950, *Stenopogon*, misident. – 128
infernalis (Wiedemann), 1821 (*Asilus*), *Mallophora* – 39
inflata (Hine), 1911 (*Erax*), *Pogonioefferia* – 22
inflatum (Hull), 1962 (*Plesiomma (Dapsilochetus)*), *Plesiomma* – 125
inflatus Scarbrough, 2003, *Ommatius* – 50
infractus Scarbrough, 1985, *Ommatius* – 51
infumata (Lynch Arribálzaga), 1880 (*Allopogon*), *Aczelia* – 57
infuscatus (Bellardi), 1861 (*Asilus*), *Prolatiforceps* – 12
infuscatus Scarbrough, 1990, *Ommatius* – 50
inornata (Loew), 1851 (*Laphria (Dasythrix)*), *Dasythrix* – 97
inornata (Rondani), 1848 (*Discocephala*), *Holcocephala* – 136
inserens Walker, 1851, *Blepharepium* – 66
insertus (Walker), 1851 (*Dasypogon*), *Blepharepium* – 67
insulara Martin, 1957, Beamerymyia – 107
insularia Martin, 1968, Beamerymyia, error – 107
intactus (Wiedemann), 1828 (*Dasypogon*), *Diogmites* – 70
?interlineata (Walker), 1860 (*Discocephala*), *Holcocephala* – 136
integerimus Scarbrough, 1990, *Ommatius* – 50
intermedius Lamas M., 1971, *Porasilus* – 23
interrupta (Macquart), 1834 (*Asilus*), *Triorla* – 24
inti Artigas, 1974, *Aymarasilus* – 119
intima Williston, 1901, *Leptogaster* – 109
invariatus (Walker), 1851 (*Asilus*), *Nerax* – 19
invasus Scarbrough & Perez-Gelabert, 2005, *Apachekolos* – 107
iopterus (Wiedemann), 1828 (*Asilus*), *Anasmostus* – 7
iracema Artigas, Papavero & Serra, 1991, *Dissmeryngodes* – 92
Isopogonini G. H. Hardy, 1948 – 62
Itolia Wilcox, 1936 – 115
Iucundus Kaletta, 1978 – 95
Ivettea Artigas & Papavero, 1991 – 115
ivetteae Artigas, 1970, *Nomomyia* – 29
jagua Scarbrough & Perez-Gelabert, 2006, *Atomosia* – 86
jaimacensis Hull, 1962, *Cerotainia*, error – 91
jalaeosus (Gowdey), 1926 (*Erax*), *Nerax*, error – 19
jalapensis (Bellardi), 1861 (*Dasypogon*), *Diogmites* – 70
jaliscoensis Williston, 1901, *Dicranus* – 120
jaloesus (Gowdey), 1926 (*Erax*), *Nerax*, error – 20
jamaicensis Farr, 1973, *Eurhabdus* – 108
jamaicensis Farr, 1963, *Leptogaster* – 109
jamaicensis Farr, 1965, *Ommatius* – 50
jamaicensis Farr, 1962, *Schildia* – 111
jamaicensis Johnson, 1919, *Cerotainia* – 91
jamesi Pritchard, 1935, *Proctacanthella* – 54
janiceae Fisher, 1977, *Zabrops* – 106
jaujensis Artigas, 1970, *Leptochelina* – 121
jemima Curran, 1941, *Mallophora* – 37
jemina Martin & Papavero, 1970, *Mallophora*, error – 37
jimagua Scarbrough & Perez-Gelabert, 2006, *Atomosia* – 86
joartigasia Martínez & Martínez, 1975 – 98
jolyi Ayala, 1981, *Pteralbis* – 12
jolyi Kaletta, 1986, *Cerotainia* – 91
Josmayala Kaletta, 1978 – 93
jubatus (Coquillett), 1904 (*Scleropogon*), *Stenopogon* – 129
jungens Schiner, 1867, *Plesiomma* – 125
juventus Scarbrough & Perez-Gelaberet, 2005, *Wilcoxius* – 47
kalettae Ayala, 1978, *Eichoichemus* – 27
kamakusa Artigas & Papavero, 1988, *Apolastaurooides* – 65
kamerlacheri Schiner, 1867, *Leptogaster* – 109
kamerlocheri Williston, 1891, *Leptogaster*, error – 109
keiseri (Carrera & Papavero), 1965 (*Alyssomyia*), *Creolestes* – 119
keisseri Artigas, 1970, *Creolestes*, error – 119
kelleri Carrera, 1952, *Threnia* – 14
knutsoni Papavero, 1971, *Aphamartania* – 62
labiata (Fabricius), 1805 (*Laphria*), *Megapoda* – 76
labidiphorus (Tomasovic), 2002 (*Efferia*), *Nerax* – 21
labidophorus (Wiedemann), 1828 (*Erax*), *Nerax* – 21
Labromyia Hull, 1962 – 57
lades (Walker), 1849: 403 (*Asilus*), *Nerax* – 21
lagunae Wilcox, 1972, *Sintoria* – 117
lampon (Walker), 1849 (*Asilus*), *Mallophora* – 42
Lampria Macquart, 1838 – 98
Lamprozona Loew, 1851 – 94
lancea Tomasovic, 2002, *Tipulogaster* – 111
landbecki (Philippi), 1865 (*Dasypogon*), *Obelophorus* – 124
langei Carrera, 1947, *Phonicocleptes* – 75
lanei (Carrera), 1949 (*Lastaurax*), *Neodiogmites* – 74
lanei Carrera, 1949, *Senobasis* – 79
lanosa (Artigas), 1970 (*Dasycyrton*), *Raulcortesia* – 117
Laphira Raguès, 1909 – 80
Laphria Meigen, 1803 – 99
Laphridae Bigot, 1857 – 80
laphriicornis Enderlein, 1914, *Cryptomerynx* – 96
Laphriida Gistel, 1848 – 80
Laphriina Rondani, 1856 – 80
Laphriinae Macquart, 1838 – 80
Laphriinae Schiner, 1860 – 81, 101
Laphriini Macquart, 1838 – 97
Laphrina Rondani, 1856 – 80
Laphritae Macquart, 1838 – 80
Laphrites Walker, 1855 – 80
Laphritidae Bigot, 1852 – 80
Laphygmolestes Hull, 1962 – 104
Laphystia Loew, 1847 – 104
Laphystiinae Hendel, 1936 – 102, 106
Laphystiinae Papavero, 1973 – 102
Laphystiini Hendel, 1936 – 102
lapila (Ayala), 1978 (*Aristofolia*), *Lecania* – 31
Lapria Meigen, 1800 – 99
lascivus (Wiedemann), 1928: 474 (*Asilus*), *Nerax* – 21
Lasiopogon Loew, 1847 – 133
Lastauracini Sabrosky, 1999, error – 65
Lastaurax Carrera, 1949 – 74
Lastaurina Curran, 1934 – 72
Lastaurini Papavero, 1973 – 65
Lastauroides Carrera, 1949 – 74
Lastauronia Carrera, 1949 – 72

- Lastauopsis* Carrera, 1949 – 73
Lastaurus Loew, 1851 – 73
lateralis (Macquart), 1838 (Erax), Triorla – 24
lateralis (Walker), 1860 (Trupanea), Amblyonychus – 34
laterepunctata (Hermann), 1912 (Atonia), Atoniomyia – 89
lateripunctata Hull, 1962, Atoniomyia, error – 89
lateropunctata Carrera & d'Andretta, Atoniomyia, error – 89
laticeps Bromley, 1929, Cerotainia – 91
laticornis (Walker), 1855 Laphria, Pilica – 83
laticrus Scarbrough & Perez-Gelabert, 2006, Ommatius – 50
latiforceps (Bromley), 1928 (Erax), Proctophoroides – 27
latipennis (Loew), 1866 (Stenopogon), Osprioncerus – 127
latipes (Wulp), 1870 (Doryclus), Pseudorus – 78
latruncula (Williston), 1885 (Erax), Carinefferia – 16
latus (Coquillett), 1904 (Holopogon), Haplopogon – 135
?latus (Philippi), 1865 (Dasypogon), Dasypodus – 114
lautus (Back), 1909 (Holopogon), Haplopogon, error – 135
lautus (Bromley), 1934 (Holopogon), Haplopogon, error – 135
lautus Pritchard, 1941, Haplopogon, error – 135
lax (Kirby), 1882 (Tolmerolestes), Prolepsis – 123
lebasii Macquart, 1838 (Asilus) – 139
lebasii Macquart, 1838 (Dasypogon) – 139
Lecania-group – 30
Lecania Macquart, 1838 – 30
leechi Wilcox, 1964, Metapogon – 116
Leinendra Carrera, 1945 – 8
lenkoi Papavero, 1975, Senobasis – 79
leonides (Walker), 1851 (Asilus), Taurhynchus – 56
leonina Hermann, 1912, Cerotainia – 91
leoninus Artigas, 1970, Lochmorhynchus – 33
Leptochelina Artigas, 1970 – 121
leptogaster Loew, 1866, Plesiomma – 126
Leptogaster Meigen, 1803 – 108
leptogaster (Perty), 1833 (Laphria), Plesiomma – 126
Leptogasterinae G. H. Hardy, 1926 – 106
Leptogastrinae Schiner, 1862 – 106
Leptogastridae Martin, 1968 – 106
Leptoharpacticus Lynch Arribálzaga, 1880 – 28
Leptopteromyia Williston, 1907 – 109
lerneri Curran, 1953, Leptogaster – 109
lerneri Curran, 1951, Proctacanthus – 54
lesbius Lamas M., 1971, Porasilius – 23
leschenaulti Macquart, 1838, Mallophora – 39
lestes (Williston), 1901 (asilus), Martintella – 46
lestomyiformis Wilcox & Martin, 1936, Nannocyrtopogon – 116
Lestophonax Hull, 1962 – 8
leucophaea Lynch Arribálzaga, 1880, Dasythrix – 97
leucopogon (Wiedemann), 1828 (Asilus), Taurhynchus – 56
leucoptera Artigas, Lewis & Parra, 2005, Tillobroma – 132
leucopyga Artigas & Angulo, 1980, Mallophora – 39
leucopyga (Wiedemann), 1828 (Asilus), Lecania – 31
leucotricha Artigas, Lewis & Parra, 2005, Tillobroma – 132
leucotricha Carrera & d'Andretta, 1953, Mallophora – 39
lewisi Farr, 1965, Andrenosoma – 81
limbata Macquart, 1834 (Laphria) – 101
limbativentris Lynch Arribálzaga, 1880, Atomosia, emend. – 86
limbiventris Tomson, 1869, Atomosia – 86
lindigii (Schiner), 1868 (Dasypogon), Diogmites – 70
lineata Curran, 1930, Atomosia – 86
lineata Hermann, 1912, Rhopalagaster – 100
lineata Scarbrough, 1996, Leptogaster – 109
lineatum (Fabricius), 1781 (Asilus), Plesiomma – 126
lineola (Bromley), 1934 (Deromyia), Diogmites – 71
lineolatus Scarbrough, 1988, Ommatius – 50
Lissoteles Bezzii, 1910 – 133
litoralis (Curran), 1930 (Deromyia), Diogmites – 71
lizbethae Ayala, 1978, Eichoichemus – 27
Lochites Schiner, 1866 – 78
Lochitomyia Brèthes, 1925 – 78
Lochmorhynchus Engel, 1930 – 32
Lochmorhynchus-group – 32
Lochyrys Artigas, 1970 – 8
loewi Artigas & Papavero, 1993, Plesiomma – 126
loewi (Bromley), 1929 (Asilus), Wilcoxius – 47
Loewiella Williston, 1896 – 63
loewii (Bellardi), 1862 (Erax), Nerax – 21
Lonchodogonus Hull, 1962 – 32
longicornis (Hermann), 1912 (Atractia), Aphractia – 85
longicornis (Wiedemann), 1828 (Laphria), Rhopalagaster – 100
longimana (Hermann), 1912 (Protichisma), Cerotainia – 91
longipennis (Bellardi), 1861 (Discocephala), Holcocephala – 136
longipennis Schiner, 1868, Threnia – 14
longipennis Thomson, 1869, Caenarolia – 68
longipes (Johnson), 1894 (Leptogaster), Apachekolos – 106
longipes (Macquart), 1838 (Dasypogon), Phonicocleptes – 75
longissimus Tomasovic, 2002, Eurhabdus – 108
longiterebratus (Macquart), 1850 (Erax), Lochmorhynchus – 33
longiungulata (Macquart), 1838 (Dasypogon), Caenarolia – 68
longiungulatus (Macquart), 1850 (Dasypogon), Dicranus – 121
longiusculus (Walker), 1855 (Asilus), Eraxasilus – 26
longiventris Schiner, 1867, Plesiomma – 126
longulus (Loew), 1866 (Stenopogon), Osprioncerus – 127
Lonquimayus Artigas & Papavero, 1991 – 116
lopesi Artigas & Papavero, 1991, Grajahuá – 130
lopesi Carrera, 1949, Senobasis – 79
lopesi Carrera & Papavero, 1965, Pritchardia – 122
lopesi Martin, 1971, Leptopteromyia – 110
Lophoceraea Hermann, 1912 – 94
lowei Papavero & Ibáñez Bernal, 2001 (Eraxc), error – 21
lucasi (Bellardi), 1861 (Dasypogon), Prolepsis – 124
lucida Curran, 1941, Mallophora – 37
lucidatus Scarbrough, 1997, Ommatius – 50
lucidus (Hermann), 1912 (Atractia), Hybozelodes – 94
lucifer (Wiedemann), 1828 (Dasypogon), Prolepsis – 123
luctuosus (Macquart), 1838 (Asilus), Eraxasilus – 26
luctuosus Macquart, 1838 (Dasypogon) – 139
Iudius (Coquillett), 1893 (Anisopogon), Heteropogon – 114
lugens (Philippi), 1865 (Dasypogon), Scylaticodes – 131
lugens Schiner, 1868, Threnia – 14
lugubre (Jaennicke), 1867 (Dioctria), Plesiomma – 126
lugubris Lynch Arribálzaga, 1880, Mallophora – 39
lugubris (Macquart), 1846 (Dasypoon), Lastaurus – 73
lugubris Williston, 1901, Scleropogon – 128
lunatus Scarbrough, 2002, Ommatius – 50
lupus (Bromley), 1931 (Andrenosoma), Pilica – 83
lurettae Martin, 1978, Machimus – 9
luridum Rondani, 1849, Blepharepium – 67
luteipes Hermann, 1926, Holcocephala – 136
lutipes (Wiedemann), 1828 (Asilus), Stenasilus – 13
lutzi Curran, 1931, Cophura – 64
Lycomya Bigot, 1857 – 9
Lycomyia, emend. – 9
Lycomyia Lynch Arribálzaga, 1882 – 7
Iycorius (Walker), 1851 (Asilus), Eichoichemus – 27
Lycosimyia Hull, 1958 – 95
lynchi Carrera, 1949, Blepharepium – 66
lynchi Gemignani, 1930, Mallophora – 36
Lynchia Williston, 1889 – 65
lynchii (Brèthes), 1901 (Myolestes), Cyrtophrys – 76
lynchii (Williston), 1889 (Cyphotomyia), Cerotainia – 91
Macahyba Carrera, 1947 – 104
Machimina Lehr, 1969 – 7
Machimini Lehr, 1977 – 7

- Machimus Loew, 1847 – 9
macquarti Bellardi, 1861, Atomosia – 86
macquartii (Perty), 1833 (Laphria), Lampria – 99
?*macquartii* (Rondani), 1848 (Mallophora), Amblyonychus – 34
macquarti Rondani, 1850, Mallophora – 40
macquarti Rondani of Arigas & Angulo, 1980, Mallophora,
 misident. – 38
Macrocolus Engel, 1930 – 58
Macronix Bigot, 1857 – 120
macropygialis (Williston), 1901 (Leptogaster), Psilonyx – 110
?*macrotelus* (Walker), 1837 (Asilus), Taurhynchus – 56
macrum Loew, 1861, Plesiomma – 126
macula Martin, 1957, Beameromyia – 107
macula (Rondani), 1848 (Discocephala), Holcocephala – 136
macularis (Wiedemann), 1821 (Asilus), Eicherax – 25
maculata Wilcox, 1936, Itolia – 115
maculatus (Macquart), 1838 (Erax), Triorla – 24
maculatus Curran, 1934, Diogmites – 71
maculipennis (Macquart), 1838 (Dasypogon), Aphamartania – 62
maculipennis (Macquart), 1855 (Senobasis), Blepharepium – 67
maculipennis (Macquart), 1846 (Laphria), Smeryngolaphria – 97
maculosus Scarbrough & Perez-Gelabert, 2006, Ommatius – 50
maestrae Broimley, 1929, Atomosia – 86
magellanica (Artigas), 1970 (Hypenetes), Tillobroma – 132
magnicauda (Curran), 1934 (Leptogaster), Psilonyx – 110
magnifica (Walker), 1854 (Dasypogon), Archilestris – 119
magnus Bellardi, 1861, Promachus – 45
magnus Scarbrough & Perez-Gelabert, 2005, Apachekolos – 107
Mallophora-group – 33
Mallophora Macquart, 1838 – 35
mallophorides (Walker), 1851 (Morimna), Lastaurus – 73
Mallophorina Curran, 1934 – 44
mallophoroides Hull, 1962, Lestophonax – 8
mallophoroides Kertész, 1909, Lastaurus, error – 73
maren Pritchard, 1941. Annamyia – 62
marga Pritchard, 1941, Aphamartania – 62
marginalis Bromley, 1946, Ommatius, error – 50
marginalis Williston, 1901, Laphria – 99
marginata (Bellardi), 1861 (Erax), Albibarbefferia – 15
marginata Hermann, 1912, Cerotainia – 91
marginatus (Osten Sacken), 1887 (Atractia), Hybozelodes – 94
marginellus (Fabricius), 1781 (Asilus), Ommatius – 50
mariae Scarbrough, 2000, Ommatius – 50
martensis Artigas & Papavero, 1989, Meliponomima – 137
Martinella Artigas & Papavero, 1995 – 46
martini Farr, 1963, Leptogaster – 109
martini Fisher, 1992, Ctenodontina – 30
martini Lamas M., 1973, Prolepsis – 123
martini Papavero, 1975, Pseudorus – 78
Martinia Hull, 1962 – 104
Martinomyia Özdiiken, 2006 – 104
martinorum (Artigas & Papavero), 1988 (Macrocolus),
 Amorimius – 58
Martintella Artigas, 1996 – 46
matilei Papavero, 1971, Alvarenga – 62
matteii Ayala, 1982, Holcocephala – 136
maya Carrera & d'Andretta, 1953, Ctenodontina – 30
maya Martin, 1968, Acronyches – 112
maya (Martin), 1957 (Pachychaeta), Ctenodontina – 30
mayi (Edwards), 1932 (Asilus), Stizolestes – 13
mcalpinei Martin, 1975, Machimus – 9
meastylus Williston, 1891 (Asilus), error – 139
media Clements & Bennett, 1969, Mallophora – 40
medianus (Wiedemann), 1828 (Asilus), Nerax – 21
medinai Artigas, 1970, Dasycyrton – 113
megacephalus (Bellardi), 1861 (Asilus), Ommatius – 50
megachile (Coquillett), 1892 (Mallophora), Megaphorus – 44
Megalometopon Artigas & Papavero, 1995 – 29
Megametopon Artigas, 1970 – 29
Megaphorus Bigot, 1857 – 44
Megapoda Macquart, 1834 – 76
Megapodina Papavero, 1975 – 75
Megapodinae Hull, 1962 – 75
Megapodini Carrera, 1949 – 75
Megapodininae Hull, 1962 – 75
megastylus Philippi, 1865 (Asilus) – 139
Megonyx Hull, 1962 – 112
melaleucus (Schiner), 1868 (Dasypogon), Neodiogmites – 74
melaleucus (Wiedemann), 1828 (Asilus), Eichoichemus – 27
melana Martin, 1957, Beameromyia – 107
melanacrus Wiedemann, 1828 (Asilus) – 139
melanocerus (Williston), 1901 (Asilus (Heligmoneura)),
 Prolatiforceps – 12
melanogaster Wiedemann, 1821, Laphria – 99
melanogaster (Wiedemann), 1821 (Dasypogon), Neodiogmites – 74
melanopogon Hermann, 1912, Atomosia – 87
melanosoma Scarbrough & Knutson, 1989, Cerotainia – 91
melanura (Wiedemann), 1828 (Laphria), Smeryngolaphria – 97
Meliponomima Artigas & Papavero, 1989 – 137
melirosoma (Carrera & Papavero), 1962 (Araiopogon), Saropogon – 60
mellina (wiedemann), 1828 (Asilus), Lecania – 31
mellipes Bromley, 1934, Saropogon – 60
mellipes Wiedemann, 1828 (Asilus) – 140
mellipes (Wiedemann), 1828 (Laphria), Eumecosoma – 93
membranosus Scarbrough, 1985, Ommatius – 50
memnon Osten Sacken, 1887, Diogmites – 71
mendax Curran, 1934, Senobasis – 79
Menexenus Artigas, 1970 – 10
meruuna Papavero, 1971, Acronyches – 112
mesoxanthum (Wiedemann), 1828 (Laphria), Andrenosoma – 81
metallescens Hermann, 1912, Atomosia – 87
metalescens Kaletta, 1974, umecosoma, error – 93
metallescens Schiner, 1868, Eumecosoma – 93
metallica Bromley, 1929, Atomosia – 87
Metapogon Coquillett, 1904 – 116
mexicana Carrera & d'Andretta, 1950, Mallophora – 43
mexicana Macquart, 1847, Lampria – 99
mexicana Williston, 1901, Aphestia – 85
mexicanae Martin, 1971, Leptopteromyia – 110
mexicanum Martin, 1960, Microstylum – 121
mexicanus (Bromley), 1951 (Doryclus), Pseudorus – 78
mexicanus (Cole), 1923 (Stenopogon), Proctacanthella – 54
mexicanus (Hine), 1919 (Erax), Nerax – 21
mexicanus Macquart, 1846 (Dasypogon) – 140
?*mexicanus* (Macquart), 1846 (Asilus), Machimus – 9
micans (Philippi), 1865 (Dasypogon), Hexameritia – 103
micans Schiner, 1867, Proctacanthus – 55
mickii Hermann, 1912, Atoniomyia, eror – 89
micronyx Tomasovic, 2002, Rhopalogaster – 100
micropygialis Williston, 1901, Leptogaster – 109
Microstylum Macquart, 1838 – 121
microthorax Aldrich, 1923, Schildia – 111
midas Osten Sacken, 1887 (Asilus); Williston, 1893 (Asilus);
 Aldrich, 1905 (Asilus), Wyliea, error or emend. – 14
mikii (Williston), 1886 (Atomosia), Atoniomyia – 89
midgon (Hull), 1962 (Nerax), Lecania, error – 31
miles (Wiedemann), 1828 (Dasypogon), Caenarolia – 68
mimus (Osten Sacken), 1877 (Ablautatus), Ablautus – 113
minima Curran, 1930, Cerotainia – 91
minor (Macquart), 1847 (Erax), Eicherax – 25
minos (Bromley), 1931 (Andrenosoma), Pilica – 83
minos Osten Sacken, 1887, Ospriccerus – 127
minos (Wiedemann), 1824 (Asilus), Mallophora – 40
minuscula (Artigas), 1970 (Dasycyrton), Ivettea – 116

- minusculus Hine, 1911, Promachus – 45
 minuta (Bellardi), 1861 (Discocephala), Holcocephala – 136
 ?minuta (Bellardi), 1861 (Discocephala), Holcocephala – 137
 minuta Williston, 1895, Townsendia – 134
 ?minutus (Wiedemann), 1828 (Laphria), Hybozelodes – 94
 mirandai (Carrera), 1951 (Alipiolaphria), Cryptomerynx – 96
Mirolestes Curran, 1935 – 75
 misera Artigas, 1973, Alyssomyia – 118
 misera (Walker), 1854 (Dasypogon), Tocantinia – 61
mitescens (Walker), 1851 (Trupanea), Amblyonychus – 35
 mixtus (Carrera), 1949 (Lastauroides), Neodiogmites – 74
 mochica Lamas M., 1973, Ctenodontina – 30
Mochtherus Loew, 1849 – 10
 modellus (Bromley), 1932 (Asilus), Stizolestes – 13
modesta (Bigot), 1878 (Senobasis), Blepharepium – 67
modesta Hermann, 1920, Helolaphycitis, nom. nud. – 103
modesta Philippi, 1865 (Laphria) – 101
modestus (Carrera), 1949 (Lastauroides), Neodiogmites – 74
 mogiana Carrera, 1955, Holcocephala – 136
 molle Bromley, 1934, Eumecosoma – 93
 mollis (Hermann), 1912 (Atonia), Atonomyia – 89
 moloch (Hull), 1962 (Martinia), Martinomyia – 104
 monensis Scarbrough, 1984, Ommatius – 50
monki Bromley, 1934, Osprioncerus – 127
montensis Scarbrough & Perez-Gelabert, 2009, Efferia – 18
monticola Ayala, 1982, Holcocephala – 136
Morimna Walker, 1851 – 73
morusum Loew, 1872, Microstylum – 122
mrazi Hradský, 1968, Phellopteron – 100
mucida Osten Sacken, 1887, Atomosia – 87
mucidus (Walker), 1837 (Asilus), Lochmorhynchus – 33
mucius (Walker), 1849 (Asilus), Leptocephalacticus – 28
multicincta Walker, 1851, Leptogaster – 109
mundata (Wiedemann), 1838 (Dasypogon), Senobasis – 80
murina (Philippi), 1865 (Erax), Nomomyia – 29
murina (Philippi), 1865 (Cylindrophora), Theromyia – 65
?mustela Hermann, 1912, Neophoneus – 101
mutabilis Loew, 1851, Lastaurus – 73
Myaptex-group – 45
Myaptex Hull, 1962 – 46
Myaptexaria Artigas & Papavero, 1995 – 46
mydas (Brauer), 1885 (Asilus), Wyliea – 14
mygdon (Walker), 1851 (Erax), Lecania – 31
Myiolestes Kertész, 1909 – 75
Myiothera Williston, 1889 – 65
Myolestes Brèthes, 1904 – 75
mystaceus (Macquart), 1938 (Proctacanthus), Taurhynchus – 56
nahuelbutae Artigas, Lewis & Parra, 2005, Tillobroma – 132
nana (Pritchard), 1941 (Aphamartania), Theromyia – 65
Nannocyrtopogon Wilcox & Martin, 1936 – 116
Nannodiocria Wilcox & Martin, 1942 – 117
narrius Scarbrough, 2002, Ommatius – 50
neariacensis (Martin & Papavero), 1970 (Machimus ((Tolmerus)),
 Prolatiforceps, error – 12
nearno Martin, 1962, Proctacanthus – 55
Neantonia Bromley, 1935 – 88
nebulosus Scarbrough, 2008, Ommatius – 50
necans (Wiedemann), 1828 (Dasypogon), Caenarolia – 68
negrabarbis Artigas, 1970, Creolestes, error – 120
negriventris Artigas, 1970, Deromyia, error – 76
nemacula Martin, 1966, Townsendia – 134
Neoderomyia Artrigas, 1971 – 59
Neodiocria Wilcox & Martin, 1941 – 117
Neodiogmites Carrera, 1949 – 74
Neoeristicus Osten Sacken, 1878 – 25
Neomochtherus Osten Sacken, 1878 – 10
Neophoneini Artigas, Papavero & Pimentel, 1988 – 100
Neopogon Bezzi, 1910 – 134
Neophoneus Williston, 1889 – 101
neoternatus (Bromley), 1931 (Deromyia), Diogmites – 71
Neotes Artigas & Papavero, 1995 – 29
neotropicus Curran, 1928, Ommatius – 50
neowillistoni (Bromley), 1935 (Erax), Eichoichemus – 27
nephressa Pritchard, 1943, Cophura – 64
Nerax Hull, 1962 – 19
neriacensis (Bromley), 1951 (Asilus), Prolatiforceps – 12
nervosa (Macquart), 1838 (Erax), Triorla – 24
Nesiotes Artigas, 1970 – 29
neta Curran, 1941, Mallophora – 43
neusae Artigas, Papavero & Costa, 1997, Cochleariocera – 103
Nicocles Jaennicke, 1867 – 64
nigellus Hermann, 1912, Hybozelodes – 94
nigellus Scarbrough, 1984, Ommatius – 50
niger (Carrera), 1949 (Lastauroides), Neodiogmites – 74
niger Martin, 1955, Parataracticus – 59
nigerrimus Carrera, 1955, Dicranus – 121
nigra Bigot, 1878, Aphestia – 85
nigra Bigot, 1878, Cerotainia – 91
nigra (Hull), 1962 (Othioniomyia (Sphagolestes)), Oidardis – 96
nigra Wiedemann, 1828, Leptogaster – 109
nigra Williston, 1885, Mallophora – 36
nigrescens (Blanchard), 1852 (Leptogaster), Deromyia, error – 76
nigrialbus Martin, 1970, Promachus – 45
nigribarbis (Philippi), 1865 (Clavator), Creolestes – 120
nigricala (Curran), 1925 (Deromyia), Diogmites – 71
nigrifemorata Macquart, 1838, Mallophora – 40
nigrimanus Curran, 1951, Proctacanthus – 55
nigrimystaceus (Macquart), 1847 (Erax), Eicherax – 25
nigrimystaceus Williston, 1901, Taracticus – 61
nigrinasutus Bigot, 1878, Saropogon – 60
nigrinus Macquart, 1848 (Asilus) – 140
nigrinus (Wiedemann), 1821 (Asilus), Nerax – 21
nigripennis (Bellardi), 1861 (Atomosia), Cerotainia – 91
nigripennis (Macquart), 1847 (Dasypogon), Diogmites – 71
nigripes (Bellardi), 1861 (Dasypogon), Diogmites – 71
nigripes (Bellardi), 1981 (Erax), Eicherax – 25
nigripes (Fabricius), 1787 (Asilus), Amblyonychus – 34
nigripes Fabricius of Wiedemann, 1828 (Asilus), Amblyonychus,
 misident. – 35
nigripes Hine, 1911, Promachus – 45
?nigripes (Macquart), 1838 (Trupanea), Amblyonychus – 34
nigripes (Macquart), 1838 (Atomosia), Dissmeryngodes – 92
nigripes (Macquart), 1850 (Erax), Nerax – 2
nigripes Williston, 1901, Taracticus – 61
nigrita (Fabricius), 1805 (Diocria), Holcocephala – 136
nigritarsis (Fabricius), 1805 (Dasypogon), Mallophora – 40
nigritarsis Fabricius of Macquart, 1838, Mallophora, misident. – 38
nigritarsis (Hine), 1919 (Erax), Nerax – 21
nigritarsis (Macquart), 1846 (Dasypogon), Diogmites – 71
nigritulus Coquillett, 1904, Stenopogon – 129
nigriventris (Bigot), 1878 (Lochites), Deromyia – 76
?nigriventris Jaennicke, 1867, Mallophora – 42
nigriventris (Philippi), 1865 (Asilus), Stizolestes – 13
nigriventris Philippi of Hull, 1962 (Asilus), Stizolestes,
 misident. – 14
nigroaenea Walker, 1851, Atomosia – 87
nigrocaudatus (Williston), 1901 (Asilus (Neoitamus)),
 Prolatiforceps – 12
nigrofemoratus Hine, 1911, Proctacanthus – 55
nigrolimbatus Martin, 1968, Stenopogon – 129
nigrostriata Engel, 1930 (Heligmoneura) – 140
nigroverticellus Bromley, 1937, Stenopogon – 129
nigrum Bromley, 1931, Andrenosoma – 81

- nigrum* Macquart, 1838, Plesiomma – 126
niphardis Hermann, 1912, Rhopalagaster – 100
nitens (Coquillett), 1904 (Stenopogon (Ospricerus)),
 Ospricerus – 127
nitida Hull, 1858, Helolaphycitis – 103
nitida (Wiedemann), 1830 (Dasypogon), Holcocephala – 136
nitidigaster Macquart, 1850 (Dasypogon) – 132
nitidula Curran, 1941, Mallophora, nom. nud. – 36
nitidula Lynch Arribálzaga, 1882, Rhatimomyia – 96
nitidus Wilcox, 1966, Saropogon – 60
niveibarba (Aldrich), 1905 (Epitriptus), Machimus, error – 10
niveibarbus (Bellardi), 1861 (Asilus), Machimus – 10
nivosa Artigas, 1970, Atractocoma – 45
nobilis Osten Sacken, 1887, Promachus – 45
nodicornis (Wiedemann), 1828 (Asilus), Cerozodus – 30
nodosipes (Enderlein), 1914 (Arthriticopus), Holcocephala – 136
Nomomyia Artigas, 1970 – 29
nordestina Carrera, 1947, Macahyba – 104
norma Curran, 1928, Ommatius – 50
notata (Bigot), 1878 (:Diogmites), Senobasis – 80
Nothopogon Artigas & Papavero, 1991 – 116
notialis Martin, 1967, Hadrokolos – 114
notocinereatus Artigas, 1970 (Dasycyrton), Lonquimayus – 116
nubeculosa Bigot, 1878, Leptogaster – 109
nuda Hermann, 1912, Atomosia – 87
numitor (Osten Sacken), 1887 (Laphria), Smeryngolaphria – 97
Nyssoprosopa Hull, 1962 – 30
- oaxacensis* (Martrin), 1968 (Stenopogon), Scleropogon – 128
Obelophorus Schiner, 1868 – 124
Oberon Carrera & Papavero, 1962 – 59
obscuripennis Enderlein, 1914, Holcocephala – 137
obscuripes Loew, 1862, Leptogaster – 109
obscurus Carrera, 1949, Diogmites – 71
obscurus (Macquart), 1838 (Erax), Nerax – 21
obtusa (Engel), 1930 (Hypenetes), Tillobroma – 132
obtusa Hull, 1962, Cnadalomyia – 8
occidens (Walker), 1851 (Dasypogon), Blepharepium – 66
occidentale (Philippi), 1865 (Asilus), Megalometopogon – 29
occidentalis (Hine), 1909 (Asilus), Machimus – 10
occidentalis Williston, 1901 (Damalis), Orrhodops – 138
ocellata Martin, 1975, Schildia – 111
ochrealis Kertész, 1909 (Stichopogon), error – 140
ocrealis Rondani, 1863 (Philammosius) – 140
oculata (Fabricius), 1805 (Dioctria), Holcocephala – 137
Oidardis Hermann, 1912 – 95
Oklahomensis Pritchrd, 1935, Promachus – 45
olbus (Walker), 1849 (Laphria), Pilica – 83
Ommatiinae Ricardo, 1929 – 47
Ommatiini Hardy, 1927 – 47
Ommatini Hull, 1962 – 47
Ommatius Wiedemann, 1821 – 47
Ommatius-group – 47
omus Pritchard, 1942, Cerotainiops – 82
opaca Coquillett, 1904, Laphystia – 104
Opopotes Hull, 1958 – 28
opposita Walker, 1851, Mallophora – 43
orcina (Wiedemann), 1828 (Asilus), Mallophora – 41
oregonae Martin, 1975, Neomochtherus – 10
orenoquensis Bigot, 1876, Ommatius – 51
oreophilus Farr, 1965, Ommatius – 51
oriens Martin, 1975, Machimus – 10
ornata (Wiedemann), 1819 (Laphria), Senobasis – 80
ornatipes James, 1953, Cerotainia – 91
Orrhodops Hull, 1954 – 137
ortegai Martin, 1968, Stenopogon – 129
Orthoneuromyia Williston, 1893 – 105
- Ospricerus* Loew, 1866 – 127
Othoniomyia Hermann, 1912 – 95
ovatus (Martin), 1967 (Promachus (Amblyonychus)),
 Amblyonychus – 34
ovatus Scarbrough, 2002, Ommatius – 51
- pachychaeta* (Bromley), 1928 (Erax), Efferia – 45
Pachychaeta Bigot, 1857 – 30
painteri Bromley, 1934, Promachus – 45
painteri Martin, 1975, Machimus – 10
painteri Pritchard, 1943, Cophura – 64
painteriorum Martin, 1968, Ospricerus – 127
pampina Aryigas, 1970, Alyssomyia – 118
pamponerooides (Edwards), 1932 (Asilus), Stizolestes – 14
Panamasilus Curran, 1930 – 97
panamensis (Curran), 1931 (Cophura), Aphamartania – 62
panamensis Curran, 1930, Atomosia – 87
panamensis (Curran), 1942 (Doryclus), Pseudorus – 78
panamensis (Curran), 1942 (Panamasilus), Smeryngolaphria,
 error – 97
papaveroi Artigas, 1970 (Dasycyrton), Lonquimayus – 116
papaveroi Artigas & Angulo, 1980, Mallophora – 41
paradoxus Hermann, 1912, Enigmomorphus – 121
parastriola Pamplona & Aires, 1999, Triorla – 24
parasylveirii Artigas & Ángulo, 1980, Mallophora – 41
Parataracticus Cole, 1924 – 59
Paratractia Hull, 1962 – 85
pardalina Hermann, 1926, Holcocephala – 137
parkeri (Wilcox), 1966 (Efferia), Pogonioeffera – 22
parphorus (Walker), 1851 (Erax), Eicherax – 25
parva (Bigot), 1857 (Dasypogon), Atomosia – 87
parva (Martin), 1973 (Systelogaster), Systologaster – 111
parvula (Bellardi), 1861 (Erax), Carineffera – 16
parvula Bigot, 1878, Lampria – 99
parvulus Schaeffer, 1916, Ommatius – 51
parvus (Bigot), 1878 (Bathypogon), Creolestes – 120
parvus Bigot, 1875, Ommatius – 51
parvus Carrera, 1948, Diogmites – 70
parvus Rondani, 1850 (Dasipogon) – 140
parvus (Walker), 1855 (Asilus), Nerax – 21
patagonensis (Artigas), 1970 (Efferia), Lochmorhynchus, error – 33
patagoniensis (Macquart), 1850 (Erax), Lochmorhynchus – 33
paurosomus Pritchard, 1935, Heteropogon – 115
pavidus Williston, 1901 (Erax) – 56
pectinata Carrera, 1955, Holcocephala – 137
pectinatipes Enderlein, 1914, Ctenodontina – 30
Pediophonus Lynch Arribálzaga, 1883 – 57
pegnai Artigas, 1970, Theromyia – 65
pennatus (Hermann), 1912 Lophoceraea, Hybozelodes – 94
pennipes Hermann, 1926, Holcocephala – 137
peralta (Wilcox), 1966 (Efferia), Albibarbeffera – 15
Perasis Hermann, 1905 – 105
peraugustimus Scarbrough, 1990, Ommatius – 51
peregrinus (Carrera & Machado-Allison), 1963 (Neoitamus),
 Pteralbis – 12
peregrinus Osten Sacken, 1887, Ommatius – 51
perfecta Curran, 1934, Tricloscelis – 106
?perfectus (Walker), 1851 (Trupanea), Amblyonychus – 34
pernambucana Artigas & Papavero, 1991, Araujoa – 118
perniger (Schiner), 1868 (Erax), Cratolestes – 17
perniger Schiner, 1868, Saropogon – 60
perplexus (Back), 1909 (Deromyia), Diogmites – 71
perrumpaea Park, 1968 and Papavero & Ibáñez Bernal, 2001 (Erax),
 Triorla, error – 24
perrumpens (Walker), 1860 (Asilus), Triorla – 24
peruae Martin, 1971, Leptopteromyia – 110
peruviana Hermann, 1912, Bathropsis – 90

- peruviana Hermann, 1926, Holcocephala – 137
peticus (Walker), 1849 (Asilus), Eraxasilus – 26
petila (Martin), 1975 (Furcilla), Scarbroughia – 47
petila Martin, 1975, Polacantha – 11
petilus (Martin), 1968 (Stenopogon), Scleropogon – 128
phalma Walker, 1849 (Dasypogon) – 140
phalna Martin & Papavero, 1970 (?Heteropogon), error – 140
Phelini Papavero, 1973, error – 124
Phellini G. H. Hardy, 1926 – 124
Phellopteron Hull, 1962 – 100
Philammosius Rondani, 1856 – 113
philippii (Schiner), 1868 (Scylaticus), Euthrixius – 130
Philonerax Bromley, 1932 – 32
Philonicus Loew, 1849 – 10
Philonotus Neihours, 1886 – 11
phoenicogaster (Hermann), 1912 (Nusa), Pilica – 84
phoenicurus (Loew), 1866 (Dizonias), Prolepsis – 124
Pholidotus Hermann, 1912 – 98
Phoneus Macquart, 1838 – 101
Phonicocleptes Lynch Arribálzaga, 1881 – 75
pica Macquart, 1850, Mallophora – 41
picea Scarbrough & Perez-Gelabert, 2009, Efferia – 18
piceus Walker, 1851, Pseudorus – 78
piceus Walker of Schiner, 1866 and Hermann, 1912, Pseudorus,
 misident. – 78
picta Carrera, 1955, Cophura – 64
picticornis Loew, 1866, Scleropogon – 128
pictipennis Hermann, 1912, Smeryngolaphria – 97
pictipennis (Schiner), 1868 (Erax), Nerax – 2
pictitarsis (Bigot), 1878 (?Laparus), Callinicus – 113
pictus Hermann, 1912, Hybozelodes – 94
pictus (Philippi), 1865 (Dasypogon), Araucopogon – 58
pilatei (Johnson), 1903 (Dizonias), Prolepsis – 124
Pilica Curran, 1931 – 83
piliferous Scarbrough, 1985, Ommatius – 51
pilipes Macquart, 1834 (Laphria) – 101
pilipes Thomson, 1869, Atomosia – 87
pilosa (Hine), 1919 (Erax), Efferia – 18
pilosa (Wilcox), 1937 (Promachina), Promachella – 44
pilosa Martin, 1966, Itolia – 115
pilosipes Lynch Arribálzaga, 1880, Atomosia, error – 87
pilosulus (Bigot), 1875 (Emphysomera), Ommatius – 51
pilosulus (Bromley), 1929 (Erax), Nerax – 21
pilosus Curran, 1928, Ommatius, error – 51
pinguis (Hermann), 1912 Atonia), Atoniomya – 89
pipliens Curran, 1941, Mallophora – 40
placida (Wulp), 1882 (Deromyia), Caenarolia – 68
plagiata (Walker), 1848 (Thereva), Stichopogon – 134
Planeteolestes Lynch Arribálzaga, 1879 – 66
Planus Scarbrough & Perez-Gelabert, 2005, Wilcoxius – 47
platycerus Hermann, 1912, Hybozelodes – 94
plaumannii (Bromley), 1951 (Doryclus), Pseudorus – 78
plazai Kalletta, 1986, Bathropsis – 90
plebeius (Osten Sacken), 1887 (Neomochtherus), Philonicus – 11
plebius Martin, 1965, Philonicus, error – 11
Plesioma Schiner, 1866, error – 125
Plesiomma Macquart, 1838 – 125
Plesiommatus Artigas & Papavero, 1991 – 125
pleuralis (Curran), 1935 (Mirolestes), Cyrtophys – 76
pleuriticum (Wiedemann), 1828 (Laphria), Eumecosoma – 93
plinthopyga (Wiedemann), 1821 (Asilus), Eccritosia – 53
plutactites Papavero, 1971, Acronyches – 112
?pluto (Lynch Arribálzaga), 1881 (Tolmerolestes), Prolepsis – 123
pluto (Wiedemann), 1828 (Asilus), Mallophora – 41
podexargenteus Enderlein, 1914, Townsendia – 134
poecilolamprus (James), 1953 (Erax), Nerax – 21
poecilopus Philippi, 1865 (Asilus) – 140
pogonias-group of Wilcox, 1966 (Efferia), Pogoniefferia – 22
pogonias (Wiedemann), 1821 (Asilus), Pogoniefferia – 22
Pogoniefferia Artigas & Papavero, 1997 – 22
Pogonosoma Rondani, 1865 – 84
Polacantha Martin, 1975 – 11
pollens Osten Sacken, 1878, Microstylum – 122
pollinifera Carrera, 1945, Glaphyropyga – 28
pollinosa Curran, 1930, Cophura – 64
pollinosa (Hull), 1962 (Nyssoprosopa), Lecania – 31
Porasilus Curran, 1934 – 23
porpinqua Kertész, 1909, Cerotainia, error – 91
portoricensis (Hine), 1919 (Erax), Efferia – 18
potamon (Walker), 1851 (Erax), Eraxasilus – 26
praelongus Scarbrough & Perez-Gelabert, 2006, Ommatius – 51
praepotens (Macquart), 1846 (Laphria), Dasyllis – 82
praestigiatus Scarbrough, 1990, Ommatius – 51
pretiosus Banks, 1911, Ommatius – 51
pritchardi Carrera, 1943, Aphamartania – 62
pritchardi Hull, 1960, Cerotainiops – 82
Pritchardia Stuardo Ortiz, 1946 – 122
Proctacantha Scudder, 1884, emend. – 54
Proctacanthella Bromley, 1934 – 54
Proctacanthus-group – 52
Proctacanthus Macquart, 1838 – 54
Proctophoroides Artigas & Papavero, 1995 – 27
Proctophorus Schiner, 1866 – 26
Prolatiforceps Martin, 1975 – 12
Prolepsis Walker, 1851 – 122
prolifica (Osten Sacken), 1887 (Erax), Carinefferia – 16
prolongatus Scarbrough, 1985, Ommatius – 51
Promachella Cole & Pritchard, 1964 – 44
Promachina Bromley, 1934 (July) – 33
Promachina Bromley, 1934 (March) – 33
Promachina Lehr, 1969 – 33
Promachus Loew, 1848 – 44
Pronomopsis Hermann, 1912 – 77
propinquua Schiner, 1868, Cerotainia – 91
propinquus (Bromley), 1928 (Erax), Echoichemus – 27
Protichisma Hermann, 1912 – 90
Protometer Artigas, Papavero & Costa, 1997 – 105
proxima Walker, 1855 (Laphria) – 101
prudens (Pritchard), 1935 (Mallophorina), Megaphorus – 44
pruinosis Carrera, 1959, Eraxasilus – 26
Prytaninae Hermann, 1920 – 103
Pseudarchilestes Bigot, 1889 – 119
Pseudoarchilestes Bigot, 1890 – 119
pseudojalapensis (Bellardi), 1862 (Dasypogon), Diogmites – 71
pseudojalapensis (Kertész), 1909 (Deromyia) Diogmites,
 emend. – 71
pseudopluto Lamas M., 1973, Prolepsis – 123
pseudorubripes Lamas M., 1972, Pronomopsis – 77
Pseudorus Walker, 1851 – 77
Pseudoryculus Carrera, 1953 – 77
Psilocurus Loew, 1874 – 105
psilogaster (Wiedemann), 1828 (Laphria), Atractia – 89
Psilonyx Aldrich, 1923 – 110
psylogaster Carrera & Vulcano, 1961, Atractia, error – 89
Pteralbis Ayala, 1981 – 12
pubescens Bromley, 1929, Atomosia – 87
pubescens Curran, 1934, Leptogaster – 109
puella (Bromley), 1932 (Hypenetes), Pritchardia – 122
puella (Wiedemann), 1828 (Laphria), Atomosia – 87
puellus Bromley, 1934, Psilocurus – 105
pulchella Curran, 1925, Mallophora – 40
pulchella Williston, 1901, Cophura – 64
pulchellus (Bellardi), 1861 (Promachus), Amblyonychus – 34
pusillus (Engel), 1885 (Emphysomera), Ommatius – 51

- pulcher Williston, 1901, *Holopogon* – 115
 pulcherimus Williston, 1901, *Saropogon* – 61
 pulchripes (Bromley), 1928 (Erax), *Wygodasilius* – 27
pullata Burmeister, 1861, *Discocephala*, nom. nud. – 138
 pulverius Scarbrough, 1967, *Ommatius* – 51
 pulverulenta Schiner, 1867, *Atractia* – 89
pumillus Papavero & Ibáñez Bernal, 2001, *Ommatius*, error – 51
 pumilus Macquart, 1838 (*Dasypogon*) – 140
 pumilus (Macquart), 1850 (Erax), *Eichoichemus* – 27
 pumilus Macquart, 1847, *Ommatius* – 51
pumilus Macquart of Bellardi, 1861, *Ommatius*, misident. – 51
pumilus (Walker), 1855 (Erax), *Albibarbefferia* – 15
punctata (Bromley), 1934 (*Andrenosoma*), *Pilica* – 84
punctifera Hermann, 1921, *Atomosia* – 87
punctipennis Coquillett, 1904 (*Metapogon*) – 116
punctipennis Macquart, 1838 (*Dasypogon*) – 140
punctipennis (Philippi), 1865 (*Clavator*), *Tillobroma* – 132
punctissima Hull, 1962, *Hodites* – 93
punicea Martin, 1957, *Beameromyia* – 107
puntarenensis Artigas, 1970, *Lochmorhynchus* – 33
purpurascens (Walker), 1856 (*Laphria*), *Andrenosoma* – 81
purpurea Artigas, Lewis & Parra, 2005, *Tillobroma* – 132
purpurea (Walker), 1850 (*Trupanea*), *Mallophora* – 37
purus Curran, 1930, *Saropogon* – 60
pusilla (Macquart), 1838 (*Laphria*), *Lampria* – 99
pusilla Macquart, 1838, *Mallophora* – 41
pusilla Wiedemann, 1828 (*Laphria*) – 101
Pycnopogon Loew, 1847 – 114
pygophorum Schiner, 1868, *Andrenosoma* – 81
Pygostylus Loew, 1866 – 64
pyrrhaeca (Wiedemann), 1828 (*Laphria*), *Andrenosoma* – 81
pyrrhogonus (Wiedemann), 1828 (*Asilus*), *Nerax* – 21
pyrrhomystax (Wiedemann), 1828 (*Asilus*), *Eichoichemus* – 27
pyrrhopygia (Wiedemann), 1828 (*Laphria*), *Pilica* – 84
pyrura Rondani, 1868, *Mallophora* – 42
- quadratus* Bellardi, 1861, *Promachus* – 44
quadratus Scarbrough, 2002, *Ommatius* – 51
quadrimaculata (Bellardi), 1861 (Erax), *Albibarbefferia* – 15
quadrimaculatum Bromley, 1929, *Andrenosoma* – 81
quadrimaculata (Bellardi), 1861 (*Dasypogon*), *Prolepsis* – 124
quadrinotata (Bigot), 1878 (*Cacodaemon*), *Prolepsis* – 123
quaterna Scarbrough, 1997, *Beameromyia* – 107
quatuorlineatus (Macquart), 1838 (*Trupanea*), *Amblyonychus* – 34
quinquemaculata Artigas, 1970, *Alyssomyia* – 118
- rabelloi* Carrera, 1952, *Threnia* – 14
ramoni Jaennicke, 1867, *Leptogaster* – 109
ramsdeni (Bromley), 1929 (*Asilus*), *Wilcoxius* – 47
rapax (Osten Sacken), 1887 (Erax), *Pogonioefferia* – 23
rapax (Walker), 1851 (*Dasypogon*), *Neodiogmites* – 74
rarus Martin, 1968, *Acronyches* – 112
Raulcortesia Artigas & Papavero, 1991 – 117
rava (Coquillett), 1893 (Efferia), *Pogonioefferia* – 23
Regasilus Curran, 1931 – 29
reinhardi Bromley, 1951, *Psilocurus* – 105
rejectus Williston, 1901, *Heteropogon* – 115
remus Tomasovic, 2002, *Diplosynapsis* – 17
renatoi Ayala, 1983, *Glyphyropyga* – 28
reticulatus (Fabricius), 1805 (*Laphria*), *Diogmites* – 71
rex Bromley, 1925, *Mallophora* – 37
Rhadinosoma Artigas, 1970 – 46
Rhatimomyia Lynch Arribálzaga, 1882 – 96
Rhatitomyia Williston, 1891, emend. – 96
Rhipidocephala Hermann, 1926 – 135
rhombungulata Carrera, 1949, *Senobasis* – 80
rhopalocera (Lynch Arribálzaga), 1882 (*Ceratotaenia*),
- Cerotainia* – 91
Rhopalogaster Macquart, 1834 – 100
riali Vieira, Castro & Bavo, 2005, *Ommatius* – 51
rica Curran, 1935, *Atomosia* – 87
ricnotes (Engel), 1930 (*Neoeristicus*), *Eicherax* – 25
rionegrensis (Lamas), 1971 (*Stenopogon*), *Carebaricus* – 112
robusta Bromley, 1951, *Proctacanthella* – 54
robusta Hermann, 1908 (*Laphystia*) – 106
robusta (Wiedemann), 1828 (*Asilus*), *Mallophora* – 41
robustus Carrera, 1949, *Lastaurus* – 73
roederi Williston, 1896, *Leptogaster* – 109
rosalesi Carrera & Machado-Allison, 1963, *Atomosia* – 87
rosariana (Carrera), 1959 (*Tolmerolestes*), *Prolepsis* – 123
rossi Martin, 1968, *Stenopogon* – 128
rossi Wilcox, 1972, *Sintoria* – 117
rubescens (Bellardi), 1861 (*Dasypogon*), *Diogmites* – 71
rubescens Bigot, 1878 (*Laphria*) – 102
rubida (Hermann), 1912 (*Atractia*), *Aphractia* – 85
rubida (Wiedemann), 1821, *Leptogaster* – 109
rubidapex (Hermann), 1912 (*Nusa*), *Pilica* – 84
rubidiventris (Macquart), 1850 (Erax), *Triorla* – 24
rubidum (Williston), 1901 (*Nusa*), *Andrenosoma* – 81
rubiginipennis Macquart, 1838 (*Dasypogon*) – 140
rubra Carrera, 1945, *Leinendera* – 8
rubricornis (Macquart), 1838 (*Proctacanthus*), *Taurhynchus* – 56
rubricornis (Philippi), 1865 (*Clavator*), *Creolestes* – 120
rubripes (Bigot), 1878 (*Scylaticus*), *Euthrixius* – 130
rubripes (Carrera & Papavero), 1962 (*Macrocolus*), *Amorimius* – 58
rubripes Hermann, 1912, *Pronomopsis* – 77
rubripes (Lynch Arribálzaga), 1881 (*Tolmerolestes*), *Prolepsis* – 123
rubriventris (Hermann), 1912 (*Pholidotus*), *Joartigasia* – 98
rubriventris (Macquart), 1850 (*Proctacanthus*), *Eccritosia* – 53
rubriventris (Macquart), 1834 (*Laphria*), *Lampria* – 99
rufa (Bromley), 1931 (*Andrenosoma*), *Dasyllis* – 83
rubrofasciata Lynch Arribálzaga, 1881, *Cylicomera* – 120
rufescens (Philippi), 1865 (*Clavator*), *Creolestes* – 120
rufibarbis (Macquart), 1838 (Erax), *Pogonioefferia* – 23
rufibasis Bigot, 1878, *Diogmites* – 71
ruficauda Bigot, 1878 (*Scylaticus*) – 132
ruficauda Curran, 1928, *Ommatius* – 51
ruficauda (Wiedemann), 1828 (*Asilus*), *Mallophora* – 42
ruficauda (Williston), 1885 (*Laphria*), *Andrenosoma* – 81
ruficaudis (Hermann), 1912 (*Pholidotus*), *Joartigasia* – 98
rufimanus (Perty), 1833 (*Dasypogon*), *Megapoda* – 76
rufina (Wiedemann), 1829 (*Asilus*), *Lecania* – 31
rufipalpis Macquart, 1838 (*Asilus*) – 140
rufipalpis (Macquart), 1838 (*Dasypogon*), *Diogmites* – 71
rufipennis (Macquart), 1838 (*Dasypogon*), *Prolepsis* – 123
rufipennis (Macquart), 1847 (*Ceraturgus*), *Taracticus* – 61
rufipennis (Wiedemann), 1828 (*Laphria*), *Pilica* – 83
rufipes (Macquart), 1838 (Erax), *Nerax* – 21
rufipes (Philippi), 1865 (*Dasypogon*), *Euthrixius* – 130
rufipes Macquart, 1847, *Atomosia* – 87
rufipes Macquart, 1838, *Lecania* – 31
rufipes Macquart, 1850, *Mallophora* – 42
rufithorax (Macquart), 1846 (Erax), *Nerax* – 22
rufipes Macquart, 1846, *Ommatius* – 51
rufitarsis Williston, 1891, *Diogmites*, error – 71
rufithorax (Wiedemann), 1828 (*Dasypogon*), *Holcocephala* – 137
rufitibia (Macquart), 1848 (Erax), *Nerax* – 22
rufitibia (Macquart), 1848 (Erax), *Efferia*, part – 18
rufiventris (Blanchard), 1852 (*Laphria*), *Andrenosoma* – 81
rufiventris Bromley, 1946, *Aphamartania*, error – 62
rufiventris Macquart, 1838, *Mallophora* – 42
rufiventris Macquart, 1838, *Proctacanthus* – 55
rufiventris (Roeder), 1887 (*Megapoda*), *Pseudorus* – 78
rufiventris Walker, 1854 (*Dasypogon*) – 140

- rufiventris Wiedemann, 1821 (*Dasypogon*) – 140
russelli Scarbrough, 1984, *Ommatius* – 51
ritulus (Wiedemann), 1821 (*Dasypogon*), *Dicranus* – 121
- saccas* Walker, 1849, *Ommatius* – 51
sackeni Hine, 1911, *Promachus* – 45
sagax (Williston), 1901 (*Erax*), *Albibarbefferia* – 16
sagitata (Hengst & Artigas), 2003 (*Philonerax*), *Lochmorhynchus*, nom. nud. – 32
sallaei (Aldrich), 1905 (*Deromyia*), *Diogmites*, error – 71
sallei (Bellardi), 1861 (*Dasypogon*), *Diogmites* – 71
saltei Bromley, 1929, *Plesiomma* – 126
saltei (Curran), 1934 (*Proctacanthus*), *Taurhynchus* – 56
saltei (Curran), 1931 (*Laphystia*), *Triclioscelis* – 106
?sandaraca (Martin), 1966 (*Sphageus*), *Prolepsis* – 123
sarae Rieda, 1998, *Cerotainia* – 91
Saropogon Williston, 1891, error – 59
sarcophaga (Hermann), 1912 (*Nusa*), *Andrenosoma* – 82
Saropogon Loew, 1847 – 59
Saropogonini Hardy, 1926 – 57
Saroponini Martin & Papavero, 1970, error – 57
satanas (Wiedemann), 1828 (*Dasypogon*), *Prolepsis* – 123
satyrus Lamas M., 1971, *Porasilus* – 23
scalarata (Hermann), 1912 (*Atonia*), *Atoniomyia* – 89
scalaris (Hermann), 1908 (*Laphystia*), *Martinomyia* – 104
scapularis (Bigot), 1878 (*Leptogaster*), *Apachekolos* – 107
scapularis Bigot, 1878, *Lampria* – 99
Scarbroughia Papavero, nom. nov. – 46
schachti Geller-Grimm, 1997, *Ichneumolaphria* – 98
schadei Bromley, 1951 (*Erax*) – 56
Schildia Aldrich, 1923 – 110
schineri (Artigas), 1970 (*Hypenetes*), *Tillobroma* – 132
schineri Martin, 1965, *Ommatius* – 52
schlingeri Artigas, 1982, *Menexenus* – 10
schnusei (Bezzi), 1910 (*Neopogon*), *Stichopogon* – 134
schnusei (Hermann), 1908 (*Laphystia*), *Martinomyia* – 104
schrottkyi Bezzi, 1910, *Dicranus* – 121
schwarzii Curran, 1934, *Mallophora* – 40
Scleropogon Loew, 1866 – 128
scopifer Schiner, 1868, *Ommatius* – 52
scopifera (Schiner), 1868 (*Discocephala*), *Holcocephala* – 137
scopifera (Wiedemann), 1828 (*Asilus*), *Mallophora* – 42
scopifera Wiedemann of Macquart, 1838; Walker, 1855; Rondani, 1850; Bigot, 1857; Jaennicke, 1867 – 40
scopipeda Rondani, 1863, *Mallophora* – 42
scopitarsis Rondani, 1863, *Mallophora* – 42
?scoriacea (Wiedemann), 1828 (*Laphria*), *Atomosia* – 87
scutellaris Lynch Arribálzaga, 1880, *Mallophora* – 43
Scylaticina Artigas & Papavero, 1991 – 130
Scylaticodes Artigas & Papavero, 1991 – 130
seabrai Carrera, 1960, *Smeryngolaphria* – 97
Seabramyia Carrera, 1958 – 138
secabile Walker of Carrera, 1949, *Blepharepium*, part, misident. – 67
secabile Walker of Wilcox, 1937; Linsley, 1960, *Blepharepium*, misident. – 67
secabilis (Walker), 1860 (*Dasypogon*), *Blepharepium* – 66
selene Curran, 1935, *Atomosia* – 88
semirufa (Wiedemann), 1828 (*Dasypogon*), *Cystoprosopa* – 125
semiustus Coquillett, 1904, *Saropogon* – 60
semiviolacea Rondani, 1848, *Mallophora* – 37
senecta Adisoemarto & Wood, 1975, *Bohartia* – 118
senectus (Wulp), 1882 (*Proctacanthus*), *Lochmorhynchus* – 33
senex Osten Sacken, 1887, *Saropogon* – 60
senilis (Wiedemann), 1828 (*Asilus*), *Porasilus* – 23
Senobasiina Papavero, 1975 – 75
Senobasina Sabrosky, 1999 – 75
Senobasis Macquart, 1838 – 78
- Senobasis* Macquart of authors, misident. – 66
Senoprosopis Macquart of authors, misident. – 12
sepia (Hull), 1962 (*Cystoprosopa*), *Plesiomma* – 125
sericans Walker, 1957 (*Asilus*) – 140
sericans Walker, 1860 (*Atomosia*) – 102
sericeus (Philippi), 1865 (*Dasypogon*), *Lamprozona* – 95
serrajiboiensis Vieira, Castro & Bravo, 2004, *Ommatius* – 52
serrula Scarbrough & Perez-Gelabert, 2009, *Efferia* – 18
servillea Williston, 1891 (*Asilus*), error – 140
servillei Macquart, 1834 (*Asilus*) – 140
servillei (Macquart), 1838 (*Phoneus*), *Neophneus* – 101
setiferous Scarbrough, 1988, *Ommatius* – 52
setigera (Hermann), 1912 (*Atonia*), *Atoniomyia* – 89
setosa Hermann, 1912, *Atomosia* – 88
setosifemur (Enderlein), 1914 (*Tapinostylus*), *Glaphyropyga* – 28
setosus Bezark, 1984, *Stenopogon* – 129
?sexmaculata (Walker), 1855 (*Asilus*), *Lecania* – 32
sexpunctatum (Williston), 1901 (*Nusa*), *Andrenosoma* – 82
sexualis Curran, 1934, *Mallophora* – 53
Shannomyioleptus Carrera, 1944 – 110
shropshirei Curran, 1930, *Eumecosoma* – 93
sicyon (Walker), 1849 (*Asilus*), *Triorla* – 24
silacea (Martin), 1973 (*Systelogaster*), *Systologaster* – 111
silaceus Martin, 1968, *Stenopogon* – 129
sillersi Hull, 1963, *Laphystia* – 104
simile Scarbrough & Perez-Gelabert, 2006, *Plesiomma* – 126
similis Bigot, 1857, *Atomosia* – 88
similis Scarbrough & Perez-Gelabert, 2005, *Wilcoxius* – 47
similis Williston, 1901, *Taracticus* – 61
simplex (Macquart), 1848 (*Erax*), *Eicherax* – 26
simulans Scarbrough, 2002, *Ommatius* – 52
sinalensis Martin, 1968, *Ospriocerus* – 127
singularis Macquart, 1838 (*Erax*) – 56
singularis Macquart, 1838, *Mallophora* – 42
Sintoria Hull, 1962 – 117
sinuosa Martin, 1975, *Polacantha* – 11
sinuosa Scarbrough & Perez-Gelabert, 2009, *Efferia* – 18
Smeryngolaphria Hermann, 1912 – 97
snowii Bezzi, 1910, *Stichopogon* – 134
soccata Thomson, 1869, *Mallophora* – 42
soccata Thomson of Lynch Arribálzaga, 1880, *Mallophora*, misident. – 37
socculata Rondani, 1848, *Mallophora* – 40
sodalis Osten Sacken, 1887, *Cophura* – 64
sodalis Osten Sacken of Williston, 1901, *Cophura*, misident. – 63
sola Scarbrough & Perez-Gelabert, 2006, *Cerotainia* – 91
solus Bromley, 1951, *Saropogon* – 60
sonora Wilcox, 1960, *Laphystia* – 104
sonorense Papavero & Bernardi, 1973, *Blepharepium* – 67
sonorensis (Forbes), 1987 (*Efferia*), *Albibarbefferia* – 16
soror Bigot, 1878, *Atomosia* – 87
sororius Scarbrough & Perez-Gelabert, 2005, *Eurhabdus* – 108
sp. Artigas & Papavero, 1995, *Ctenodontina* – 30
sp. Artigas & Papavero, 1995, *Leptoharpacticus* – 28
sp. Artigas & Papavero, 1995, *Lochmorhynchus* – 33
sp. Farr, 1963, *Eurhabdus* – 108
†sp. Grimaldi, 1990, *Arariopogon* – 138
sp. Papavero, 1975, *Cyrtophrys* – 76
†sp. Poinar, 1992, *Pilica* – 84
†sp. Poinar, 1992, *Wilcoxia* – 117
sp. Scarbrough & Perez-Gelabert, 2006, *Antoniomyia* – 89
sp. Scarbrough & Perez-Gelabert, 2006, *Pilica* – 84
sp. C Scarbrough & Perez-Gelabert, 2009, *Eifferia* – 18
sp. near dasypus Artigas, Papavero & Serra, 1991, *Atomosia* – 88
spathulatus (Bellardi), 1861 (*Dasypogon*), *Ospriocerus* – 127
spatulatus Curran, 1928, *Ommatius* – 52
spatulatus Pritchard, 1935, *Heteropogon* – 115

- speciosa* Curran, 1934, Mallophora – 39
speciosus (Philipp), 1865 (Erax), Eccritosia – 53
spectabilis (Philippi), 1865 (Asilus), Cratolestes – 17
spectans (Walker), 1851 (Dasypogon), Phonicocleptes – 75
Sphageus Loew, 1861 – 122
Sphagolestes Hull, 1962 – 95
spiculatus (Hull), 1962 (Catonomyia), Dissmeryngodes – 92
spinipes Artigas, Lewis & Parra, 2005, Tillobroma – 133
spinipes (Fabricius), 1805 (Laphria), Lampria – 99
spinipes Hermann, 1926, Holcocephala – 137
spinitarsis Bromley, 1951, Leptogaster – 109
spinosa Tomasovic, 2002, Triorla – 24
spinosus Scarbrough, 1993, Ommatius – 52
spinulosa Scarbrough & Perez-Gelabert, 2009, Efferia – 19
spissibarbis Macquart, 1846 (Trupanea) – 56
spitzi Carrera, 1949, Caenarolia – 68
splendens (Macquart), 1834 (Laphria), Lampria – 99
splendens (Philippi), 1865 (Dasypogon), Hexameritia – 104
splendens (Williston), 1901 (Erax), Pogonoefferia – 23
spp. Artigas & Papavero, 1995, Lecania – 32
squamipes Cole, 1924, Ablautus – 113
staurophora (Schiner), 1868 (Lochites), Senobasis – 79
staurophorum Schiner, 1868, Eumecosoma – 93
Stenasilus Carrera, 1960 – 12
Stenobasis Agassiz, 1846, emend. – 78
Stenobasis Kertész, 1909, emend. – 78
Stichopogon Wilcox, 1936, error – 134
Stenopogon Loew, 1847 – 128
Stenopogoninae Hull, 1962 – 112
Stenopogoninae Paramonov, 1966 – 111
Stenopogonini Hull, 1962 – 111, 127
Stichopogon Loew, 1847 – 134
Stichopogoninae Artigas & Papavero, 1988 – 133
Stichopogoninae G. H. Hardy, 1930 – 133
Stichopogonini G. H. Hardy, 1930 – 133
Stictopogon Wilcox, 1937, error – 134
stigmatus (Carrera & d'Andretta), 1950 (Erax), Nerax – 22
Stilopogon Costa, 1883 – 134
stimicon (Walker), 1851 (Erax), Triorla – 24
Stizolestes Hull, 1962 – 13
stonei Bromley, 1937, Stenopogon – 129
stramineus Scarbrough, 1984, Ommatius – 52
strigarius Curran, 1931, Regasilus – 29
strigitibia Curran, 1931 (Dysmachus) – 57
striola (Fabricius), 1805 (Dasypogon), Triorla – 24
Strobilopygius Hull, 1956 – 122
Strombocodia Hermann, 1912 – 96
stylata (Fabricius), 1773 (Asilus), Efferia – 19
stylata Pritchard, 1938, Holcocephala – 137
subappendiculatus (Macquart), 1838 (Erax), Nerax – 22
subchalybeus (Bromley), 1928 (Erax), Nerax – 22
subcontractum (Walker), 1856 (Dasypogon), Blepharepium – 67
subheros Bromley, 1931, Andrenosoma – 82
subialapensis, Diogmites, error – 71
submaculus Martin, 1975, Machimus – 10
substitalis (Walker), 1851 (Trupanea), Amblyonychus – 34
†subtus Scarbrough & Poinar, 1992, Ommatius – 52
subviolacea Rondani, 1850, Mallophora, error – 37
sucinopedis Artigas, 1970, Dasycyrton – 114
sundra Pritchard, 1943, Cophura – 64
superbus Carrera, 1953, Diogmites – 71
surufus Martin, 1968, Stenopogon – 129
surumu Papavero & Bernardi, 1973, Blepharepium – 67
suspiciosa Scarbrough & Perez-Gelabert, 2009, Efferia – 19
sylveirii Macquart, 1838, Mallophora – 42
symmachus Loew, 1872, Diogmites – 72
Systelogaster Hermann, 1926 – 111
Systologaster, nom. nov. – 111
tachirensis Ayala, 1983, Glaphyropyga – 28
taeniatus (Bellardi), 1861 (Asilus, Philonicus), Philonicus – 11
tagax (Williston), 1883 (Triclis), Zabrops – 106
taina Scarbrough & Perez-Gelabert, 2006, Proctacanthella – 54
talabrensis Artigas, 1964, Pronomopsis – 77
tanbarki Wilcox, 1965, Cophura – 64
tandapiensis Scarbrough, 2002, Ommatius – 52
tanpadiensis Scarbrough, 2002, Ommatius, error – 52
Taperigna Artigas & Papavero, 1991 – 124
taperignae Artigas, Papavero & Pimentel, 1988, Smeryngolaphria – 97
Tapinostylus Enderlein, 1914 – 28
Taracticus Loew, 1872 – 61
tarsale Curran, 1930, Eumecosoma – 93
tatius Walker, 1851 (Asilus) – 140
tau Oten Sacken, 1887, Diogmites – 72
tauauana Artigas & Papavero, 1988, Neodiogmites – 74
Taurhynchus Artigas & Papavero, 1995 – 55
tenebrosus Carrera, 1949, Neodiogmites – 74
tenebrosus (Coquillett), 1904 (Stenopogon), Ospriocerus – 128
tenebrosus (Williston), 1901 (Asilus), Machimus – 10
tenebrosus Williston of Williston, 1901, Machimus, misident. – 9
tener (Bigot), 1878 (Holopogon), Lonquimayus – 116
tenuis Curran, 1930, Atomasia – 88
tenuis (Walker), 1851 (Dioctria), Cyrtophrys – 75
tenuis (Wiedemann), 1828 (Asilus), Stenasilus – 13
tenuiventris Macquart, 1866 (Asilus) – 140
tepui Scarbrough, 2008, Ommatius – 52
tequilae Martin, 1968, Ospriocerus – 128
terebratus (Macquart), 1850 (Dasypogon), Obelophorus – 124
teresita Lamas M., 1972, Diogmites – 72
ternatus (Loew), 1866 (Deromyia), Diogmites – 72
tertialis (Bromley), 132 (Hypenetes), Pritchardia – 122
tertiavitrea Artigas & Angulo, 1980, Mallophora – 43
tessellata (Wiedemann), 1828 (Dasypogon), Caenarolia – 68
testacea Loew of Williston, 1901, Leptogaster, misident. – 109
testaceipes Macquart, 1850, Mallophora – 36
testaceitarsis Macquart, 1954, Mallophora – 36
testaceum (Fabricius), 1805 (Dasypogon), Plesiomma – 126
testaceum Macquart, 1838, Plesiomma – 125
texana (Bromley), 1919 (Erax), Efferia – 18
texanus (Bromley), 1931 (Stenopogon), Scleropogon – 128
therevinus Rondani, 1850 (Asilus) – 140
therimachus (Walker), 1851 (Asilus), Threnia – 14
Theromyia Williston, 1891 – 65
thologaster Fisher, 1977, Zabrops – 106
thompsoni Artigas & Angulo, 1980, Mallophora – 43
thoracica Fabricius, 1805, Laphria – 100
Threnia Schiner, 1866 – 14
tiarense (Kaletta), 1978 (Josmayala), Eumecosoma – 93
tiarensis Ayala, 1983, Glaphyropyga – 28
tibialis Curran, 1934, Senobasis – 79
tibialis (Hengst & Artigas), 2003 (Philonerax), Lochmorhynchus, nom. nud. – 32
tibialis Macquart, 1846, Atomasia – 88
tibialis (Macquart), 1838 (Erax), Nerax – 22
tibialis Macquart, 1838, Mallophora – 43
tibialis Macquart, 1846, Mallophora – 40
tibialis (Macquart), 1850 (Proctacanthus), Taurhynchus – 56
?tibialis (Rondani), 1850 (Asilus), Eraxasilus – 26
tijucana Carrera, 1958, Seabramyia – 138
Tillobroma Hull, 1962 – 131
Tillobromatini Artigas & Papavero, 1991 – 130
Tillobromatini Sabrosky, 1999 – 130
Tillobromini Artigas & Papavero, 1991 – 130
timberlakei Wilcox, 1949, Itolia – 115

- Tipulogaster Cockerell, 1913 – 111
tisiphones Rondani, 1848, Mallophora – 41
tissiphones Schiner, 1866, Mallophora, error – 42
titan (Bromley), 1934 (Erax), Nerax – 22
titan (Carrera), 1959 (Promachina), Amblyonychus – 35
titanus Carrera, 1958, Leptogaster – 109
Tocantinia Carrera, 1955 – 61
toconaoensis Artigas, 1970, Zoticus – 30
tolandi Wilcox, 1959, Cophura – 64
tolandi Wilcox, 1961, Hodophylax – 64
tolandi Wilcox, 1965, Proctacanthella – 54
Tolmerolestes Lynch Arribálzaga, 1881 – 122
tolmides (Walker), 1849 (Laphria), Lampria – 99
tolteca Curran, 1941, Mallophora – 40
tornowii Brèthes, Leptogaster – 109
tortola (Curran), 1928 (Erax), Nerax – 22
Townsendia Williston, 1895 – 134
tractus Scarbrough, 2006, Ommatius – 52
transatlantica Schiner, 1868 (Laphria) – 102
transiens (Walker), 1949 (Dasypogon), Lastaurus – 73
trapezoidalis (Bellardi), 1861 (Promachus), Amblyonychus – 35
travassosi (Carrera), 1949 (Lastauronia), Lastaurina – 72
tregualemensis Artigas, Lewis & Parra, 2005, Tillobroma – 132
triangularis Artigas & Papavero, 1991, Nothopogon – 116
triangularis (Hermann), 1912 (Othoniomyia), Oidardis – 96
triangularis Scarbrough, 2002, Ommatius – 52
triangulata Martin, 1968, Leptogaster, error – 109
triangulata Martin, 1966, Townsendia – 135
tricella (Bromley), 1951 (Erax), Efferia – 18
tricellus Wilcox, 1964, Metapogon – 116
?trichina Tomasovic, 2002, Triorla – 25
trichonotus (Wiedemann), 1828 (Asilus), Amblyonychus – 35
trichostica Williston, 1901, Mallophora – 37
Tricloscelis Roeder, 1900 – 105
tricolor (Bellardi), 1861 (Erax), Albibarbeffera – 15
tricolor (Bellardi), 1861 (Dasypogon), Diogmites – 72
tricolor Carrera & Machado-Allison, 1968, Lastaurus – 73
tricolor (Philippi), 1865 (Dasypogon), Scylaticodes – 131
tricolor (Schiner), 1868 (Triclis), Hexameritia – 104
tricolor (Walker), 1850 (Mallophora), Eccritosia – 53
tridens Martin, 1975, Polacantha – 11
trifarius Loew, 1866, Ablautus – 113
trifasciatus (Say), 1823 (Dasypogon), Stichopogon – 134
Trigomiminae Lehr, 1988 – 135
Trigonomiminae Papavero, 1973 – 135
Trigonomiminae Enderlein, 1914 – 135
Trigonomimini Enderlein, 1914 – 135
triligata Walker, 1861 (Laphria) – 102
Triorla Parks, 1968 – 23
tripartita (Walker), 1854 (Dasypogon), Cleptomyia – 58
tristis (Walker), 1851 (Dasypogon), Prolepsis – 123
tristis Wiedemann, 1828 (Asilus) – 140
triton (Osten Sacken), 1887 (Erax), Pogonieffera – 23
triungulata Williston, 1901, Leptogaster – 109
tropica Curran, 1934, Leptogaster – 109
tropidus Scarbrough, 2002, Ommatius – 52
truncatus Martin, 1975, Machimus – 9
truncus Martin, 1975, Wilcoxius – 47
Trupanea Macquart, 1838 – 44
truquii Bellardi, 1861, Leptogaster – 108
truquii (Bellardi), 1861 (Asilus, Mochterus), Philonicus – 11
truquii Bellardi, 1861, Promachus – 45
truquii (Bellardi), 1861 (Dasypogon), Stenopogon – 129
truquii Bellardi of Williston, 1901, Stenopogon, misident. – 129
trux McAtee, 1919, Laphria – 100
tsacasi Artigas & Angulo, 1980, Mallophora – 43
tsacasi Papavero, 1971, Aczelia – 57
- Tsacasia* Artigas & Papavero, 1995 – 29
tucma Lynch Arribálzaga, 1880, Dicranus – 121
tuberculata (Coquillett), 1904 (Erax), Efferia – 18
tucumana Artigas & Papavero, 1991, Scylaticina – 130
tucumanensis Scarbrough, 2002, Ommatius – 52
tumbrensis (Artigas), 1980 (Eremomyia), Lochmorhynchus – 33
tumidus Scarbrough & Perez-Gelabert, 1995, Wilcoxius – 47
tuxpanguus (Bellardi), 1862 (Asilus), Philonicus – 11
- umbrina* Artigas & Angulo, 1980, Mallophora, nom. nud. – 38
uncatus Scarbrough, 1993, Ommatius – 52
unguiculatus Scarbrough, 2002, Ommatius – 52
unicolor (Bellardi), 1861 (Erax), Carineffera – 16
unicolor Hermann, 1912, Cerotainia – 92
unicolor Macquart, 1838, Atomesia – 88
uruguayensis Lynch Arribálzaga, 1882, Holcocephala – 137
- valdivianus* (Philippi), 1865 (Asilus), Stizolestes – 13
valentinei (Artigas), 1970 (Hypenetes), Tillobroma – 132
vallestris Ayala, 1982, Holcocephala – 137
vanduzeei Cole, 12923, Lissoteles – 133
vandykei Wilcox & Martin, 1936, Cyrtopogon – 113
varipennis (Walker), 1855 (Ampyx), Pseudorus – 78
varipes (Banks), 1920 (Laphria), Andrenosoma – 82
varipes (Schiner), 1867 (Senoprosopis), Stenasilus – 13
vauriei (Curran), 1933 (Erax), Nerax – 22
vegeta Lynch Arribálzaga, 1882, Mallophora – 41
velox (Wiedemann), 1828 (Asilus), Nerax – 22
velutinus (Carrera & Papavero), 1962 (Oberon), Saropogon – 61
venezuelanus Kaletta, 1976, Stichopogon – 134
venezuelensis Carrera & Machado-Allison, 1963, Glaphyropyga – 28
venezuelensis Macquart, 1846 (Laphria) – 102
ventralis Coquillett, 1898, Osprioncerus – 127
ventralis Macquart, 1838, Mallophora – 36
ventralis Martin, 1975, Machimus – 10
venustula Lynch Arribálzaga, 1880, Atomesia – 88
venustus (Bromley), 1929 (Leptogaster), Psilonyx – 110
venustus Carrera & d'Andretta, 1950, Promachus – 45
venustus (Philippi), 1865 (Dasypogon), Euthrixius – 130
veredus Artigas, 1970, Menexenus – 10
vertebrata (Bromley), 1940 (Erax), Albibarbeffera – 16
vetusta (Walker), 1837 (Asilus), Eccritosia – 53
vexillaria (artigas), 1970 (Myapex), Myaptexaria – 46
?vicina (Macquart), 1838 (Dioctria), Holcocephala – 137
viduata (Wiedemann), 1819 (Laphria), Atoniomysia – 89
villosus (Bellardi), 1861 (Erax), Triorla – 23
villosus (Carrera), 1949 (Lastauropsis), Lastaurus – 73
villosus Scarbrough, 1985, Ommatius – 52
violacea Macquart, 1846 (Laphria) – 102
violaceithorax (Lynch Arribálzaga), 1880 (Ceratotaenia), Cerotainia – 92
violaceus Williston, 1901, Holopogon – 115
virescens (Bellardi), 1861 (Dasypogon), Diogmites – 72
virginensis Scarbrough, 1977, Beameromyia – 107
virginianus (Wulp), 1882 (Proctacanths), Pogonieffera – 23
virilis (Artigas), 1970 (Myapex), Myaptexaria – 46
virilis (Wiedemann), 1828 (Erax), Lecania – 32
vitreus Bigot, 1875, Ommatius – 52
vitripennis (Bellardi), 1861 (Ceratiturgus), Taracticus – 61
vitripennis Schiner, 1867, Leptogaster – 109
vittata (Walker), 1837 (Xylophagus), Holcocephala – 137
vittata (Wiedemann), 1828 (Dasypogon), Caenarolia – 68
vittatus (Lynch Arribálzaga), 1880 (Proctacanthus), Taurhynchus – 56
vittatus (Olivier), 1789 (Asilus), Proctacanthus – 55
vivax (Hermann), 1912 (Atractia), Aphractia – 85

Catalogue of the Neotropical Diptera. Asilidae

- vivus Scarbrough, 1997, *Ommatius* – 52
 volubilis Scarbrough & Perez-Gelabert, 2005, *Apachekolos* – 107
Voluptarius Kaletta, 1978 – 95
vorax Curran, 1942, *Blepharepium* – 66
vorax Curran, 1934, *Leptogaster* – 109
vorax Curran, 1941, *Mallophora* – 37
vulgaris Carrerag, 1947, *Diogmites* – 72
vultus Scarbrough & Perez-Gelabert, 2005, *Apachekolos* – 107
wagneri Artigas & Papavero, 1995, *Tsacasia* – 29
wenzeli Papavero & Bernardi, 1974, *Archilestris* – 119
westcotti Martin, 1968, *Acronymes* – 112
weyenberghi (Wulp), 1882 (*Deromyia*), *Caenarolia* – 68
weyrauchi Carrera, 1952, *Senobasis* – 80
wiedemanni (Schiner), 1868 (*Promachus*), *Amblyonychus* – 35
wilcoxi Bromley, 1937, *Stenopogon* – 129
wilcoxi Kaletta, 1983, *Cophura* – 63
wilcoxi Papavero, 1971, *Saropogon* – 61
wilcoxi (Pritchard), 1935 (*Buckellia*), *Cophura* – 64
wilcoxi Pritchard, 1942, *Cerotainiops* – 82
Wilcoxia James, 1941 – 117
Wilcoxius Martin, 1975 – 47
wilhelmi Artigas & Angulo, 1980, *Mallophora* – 43
williamsi Wilcox, 1971, *Stenopogon* – 129
willinkinorum Artigas, Papavero & Costa, 1997, *Chrysotriclis* – 103
willistoni (Bromley), 1928 (*Erax*), *Eichoichemus* – 28
willistoni (Cole), 1964 (*Mallophora*), *Megaphorus* – 44
willistoni Curran, 1930, *Cerotainia* – 92
willistoni Curran, 1928, *Ommatius* – 52
willistoni Hermann, 1921, *Acronymes* – 112
willistoni Hermann, 1926, *Leptopteromyia*, nom. nud. – 110
willistoni (Hine), 1919 (*Erax*), *Carinefferia* – 16
willistoni Martin, 1962, *Heteropogon* – 115
willistoni Pritchard, 1943, *Cophura* – 64
willistoniana Enderlein, 1914, *Laphria* – 100
winthemi (Wiedemann), 1821 (*Dasytopogon*), *Diogmites* – 72
wirthi Artigas, 1970, *Cratolestes* – 17
wirthi Paramonov, 1964, *Eccritosia* – 53
woodleyi Scarbrough & Perez-Gelabert, 2009, *Eifferia* – 19
Wygodasilius Artigas & Papavero, 1995 – 27
wygodzinskyi Carrera, 1949, *Diogmites* – 72
Wyliea Martin, 1975 – 14
xanthocera (Williston), 1901 (*Asilus* (*Heligmoneura*)),
 Polacantha – 11
xanthocnemum (Wiedemann), 1828 (*Laphria*), *Andrenosoma* – 82
xanthopogon (Burmeister), 1861 (*Asilus* (*Proctacanthus*)),
 Eccritosia – 53
xanthoptera (Wiedemann), 1828 (*Laphria*), *Cerotainia* – 92
xanthopterus (Wiedemann), 1828 (*Asilus*), *Taurhynchus* – 56
xanthopus (Wiedemann), 1828 (*Laphria*), *Atomosia* – 88
Xenomyzini G. G. Hardy, 1948 – 135
xochimilcae Martin, 1968, *Stenopogon* – 129
xylocopides Walker, 1855; Schiner, 1866; Williston, 1891,
 Mallophora, error – 36
xylocopides Walker, 1851, *Mallophora* – 36
xylota (Curran), 1942 (*Panamasilus*), *Smeryngolaphria* – 97
yepezi (Carrera & Machado-Allison), 1962 (*Promachina*),
Amblyonychus – 35
youngi Martin, 1968, *Ospricerus* – 128
yuma (Wilcox), 1966 (*Efferia*), *Pogonoefferia* – 23
yurabia Scarbrough & Perez-Gelabert, 2006, *Atomosia* – 88
Zabrops Hull, 1958 – 106
Zabrotica Hull, 1958 – 132
zamon (Townsend), 1895 (*Proctacanthus*), *Eccritosia* – 53
zandra Pritchard, 1943, *Cophura* – 64
zanutoi Artigas, Papavero & Pimentel, 1988, *Pilica* – 84
zebra Martin, 1968, *Stenopogon* – 129
zephyreus Aldrich, 1923, *Eurhabdus* – 108
zephyrus Scarbrough & Page, 2005, *Psilonyx* – 110
zetterstedti (Jaennicke), 1865 (*Erax*), *Triorla* – 24
zikani Carrera, 1951, *Ichneumolaphria* – 98
zita Curran, 1941, *Mallophora* – 43
zonatus Cole & Wilcox, 1938, *Lasipogon* – 133
zona Martin, 1975, *Schildia* – 111
Zoticus Artigas, 1970 – 30
zottai Gemignani, 1930, *Mallophora* – 39